

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/51, G06F 17/50	A2	(11) International Publication Number: WO 97/26277 (43) International Publication Date: 24 July 1997 (24.07.97)
(21) International Application Number: PCT/US97/01071 (22) International Filing Date: 22 January 1997 (22.01.97) (30) Priority Data: 08/589,552 22 January 1996 (22.01.96) US (71) Applicants: CREATIVE BIOMOLECULES, INC. [US/US]; 45 South Street, Hopkinton, MA 01748 (US). BRANDEIS UNIVERSITY [US/US]; 45 South Street, Waltham, MA 02254-9110 (US). (72) Inventors: KECK, Peter; 50 Dolan Road, Millbury, MA 01527 (US). GRIFFITH, Diana, L.; 15 Woodridge Circle, Weston, MA 02193 (US). CARLSON, William, D.; 40 Black Oak Road, Weston, MA 02193 (US). RUEGER, David, C.; 19 Downey Street, Hopkinton, MA 01748 (US). SAMPATH, Kuber, T.; 6 Spring Street, Medway, MA 02053 (US). (74) Agent: GREENHALGH, Duncan, A.; Testa, Hurwitz & Thibault, L.L.P., High Street Tower, 125 High Street, Boston, MA 02110 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHODS AND COMPOSITIONS FOR PRODUCING MORPHOGEN ANALOGS		
(57) Abstract The invention disclosed herein provides methods and compositions for the computer-assisted design of morphogen analogs. Practice of the invention is enabled by the use of at least a portion of the atomic co-ordinates defining the three-dimensional structure of human osteogenic protein-1 (hOP-1) as a starting point in the design of the morphogen analogs. In addition, the invention provides methods for producing morphogen analogs of interest, and methods for testing whether the resulting analogs mimic or agonize human OP-1-like biological activity. The invention also provides a family of morphogen analogs produced by such methods.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

**METHODS AND COMPOSITIONS FOR
PRODUCING MORPHOGEN ANALOGS**

Related Applications

This application is a continuation-in-part of copending application U.S.S.N. 08/589,552,
5 filed January 22, 1996, the disclosure of which is incorporated herein by reference.

Field of the Invention

The present invention relates generally to methods and compositions for designing,
identifying, and producing compounds useful as tissue morphogenic protein analogs. More
specifically, the invention relates to structure-based methods and compositions useful in
10 designing, identifying, and producing molecules which act as functional mimetics of the tissue
morphogenic protein osteogenic protein-1 (OP-1).

Background of the Invention

Cell differentiation is the central characteristic of tissue morphogenesis which initiates
during embryogenesis, and continues to various degrees throughout the life of an organism in
15 adult tissue repair and regeneration mechanisms. The degree of morphogenesis in adult tissue
varies among different tissues and is related, among other things, to the degree of cell turnover in
a given tissue.

The cellular and molecular events which govern the stimulus for differentiation of cells is
an area of intensive research. In the medical and veterinary fields, it is anticipated that discovery
20 of the factor or factors which control cell differentiation and tissue morphogenesis will advance
significantly the ability to repair and regenerate diseased or damaged mammalian tissues and
organs. Particularly useful areas for human and veterinary therapeutics include reconstructive
surgery, the treatment of tissue degenerative diseases including, for example, arthritis,
emphysema, osteoporosis, cardiomyopathy, cirrhosis, degenerative nerve diseases, inflammatory
25 diseases, and cancer, and in the regeneration of tissues, organs and limbs. In this and related
applications, the terms "morphogenetic" and "morphogenic" are used interchangeably.

- 2 -

A number of different factors have been isolated in recent years which appear to play a role in cell differentiation. Recently, a distinct subfamily of the "superfamily" of structurally related proteins referred to in the art as the "transforming growth factor- β (TGF- β) superfamily of proteins" have been identified as true tissue morphogens.

5 The members of this distinct "subfamily" of true tissue morphogenic proteins share substantial amino acid sequence homology within their morphogenetically active C-terminal domains (at least 50% identity in the C-terminal 102 amino acid sequence), including a conserved six or seven cysteine skeleton, and share the *in vivo* activity of inducing tissue-specific morphogenesis in a variety of organs and tissues. The proteins apparently contact and interact
10 with progenitor cells e.g., by binding suitable cell surface molecules, predisposing or otherwise stimulating the cells to proliferate and differentiate in a morphogenetically permissive environment. These morphogenic proteins are capable of inducing the developmental cascade of cellular and molecular events that culminate in the formation of new organ-specific tissue, including any vascularization, connective tissue formation, and nerve innervation as required by
15 the naturally occurring tissue. The proteins have been shown to induce morphogenesis of both bone cartilage and bone, as well as periodontal tissues, dentin, liver, and neural tissue, including retinal tissue.

True tissue morphogenic proteins identified to date include proteins originally identified as bone inductive proteins. These include OP-1, (osteogenic protein-1, also referred to in related
20 applications as "OP1"), its *Drosophila* homolog, 60A, with which it shares 69% identity in the C-terminal "seven cysteine" domain, and the related proteins OP-2 (also referred to in related applications as "OP2") and OP-3, both of which share approximately 65-75% identity with OP-1 in the C-terminal seven cysteine domain, as well as BMP5, BMP6 and its murine homolog, Vgr-1, all of which share greater than 85% identity with OP-1 in the C-terminal seven cysteine
25 domain, and the BMP6 *Xenopus* homolog, Vgl, which shares approximately 57% identity with OP-1 in the C-terminal seven cysteine domain. Other bone inductive proteins include the CBMP2 proteins (also referred to in the art as BMP2 and BMP4) and their *Drosophila* homolog, DPP. Another tissue morphogenic protein is GDF-1 (from mouse). See, for example, PCT documents US92/01968 and US92/07358, the disclosures of which are incorporated herein by
30 reference. Members of the BMP/OP subfamily and the amino acid sequence identities

(expressed as percentages) between selected members of the TGF- β superfamily are shown in Fig. 6.

As stated above, these true tissue morphogenic proteins are recognized in the art as a distinct subfamily of proteins different from other members of the TGF- β superfamily in that they share a high degree of sequence identity in the C-terminal domain and in that the true tissue morphogenic proteins are able to induce, on their own, the full cascade of events that result in formation of functional tissue rather than merely inducing formation of fibrotic (scar) tissue. Specifically, members of the family of morphogenic proteins are capable of all of the following in a morphogenetically permissive environment: stimulating cell proliferation and cell differentiation, and supporting the growth and maintenance of differentiated cells. The morphogenic proteins apparently also may act as endocrine, paracrine or autocrine factors.

The morphogenic proteins are capable of significant species "crosstalk." That is, xenogenic (foreign species) homologs of these proteins can substitute for one another in functional activity. For example, dpp and 60A, two *Drosophila* proteins, can substitute for their mammalian homologs, BMP2/4 and OP-1, respectively, and induce endochondral bone formation at a non-bony site in a standard rat bone formation assay. Similarly, BMP2 has been shown to rescue a dpp⁻ mutation in *Drosophila*. In their native form, however, the proteins appear to be tissue-specific, each protein typically being expressed in or provided to one or only a few tissues or, alternatively, expressed only at particular times during development. For example, GDF-1 appears to be expressed primarily in neural tissue, while OP-2 appears to be expressed at relatively high levels in early (e.g., 8-day) mouse embryos. The endogenous morphogens may be synthesized by the cells on which they act, by neighboring cells, or by cells of a distant tissue, the secreted protein being transported to the cells to be acted on.

A particularly potent tissue morphogenic protein is OP-1. This protein, and its xenogenic homologs, are expressed in a number of tissues, primarily in tissues of urogenital origin, as well as in bone, mammary and salivary gland tissue, reproductive tissues, and gastrointestinal tract tissue. It is expressed also in different tissues during embryogenesis, its presence coincident with the onset of morphogenesis of that tissue.

- 4 -

The morphogenic protein signal transduction across a cell membrane appears to occur as a result of specific binding interaction with one or more cell surface receptors. Recent studies on cell surface receptor binding of various members of the TGF- β protein superfamily suggests that the ligands mediate their activity by interaction with two different receptors, referred to as Type I and Type II receptors to form a hetero-complex. A cell surface bound beta-glycan also may enhance the binding interaction. The Type I and Type II receptors are both serine /threonine kinases, and share similar structures: an intracellular domain that consists essentially of the kinase, a short, extended hydrophobic sequence sufficient to span the membrane one time, and an extracellular domain characterized by a high concentration of conserved cysteines.

Morphogenic proteins are disulfide-linked dimers which are expressed as large precursor polypeptide chains containing a hydrophobic signal sequence, a long and relatively poorly conserved N-terminal pro region of several hundred amino acids, a cleavage site and a mature domain comprising an N-terminal region which varies among the family members and a more highly conserved C-terminal region. The C-terminal region, which is present in the processed mature proteins of all known morphogen family members, contains approximately 100 amino acids with a characteristic motif having a conserved six or seven cysteine skeleton. Each of the morphogenic proteins isolated to date are dimeric structures wherein the monomer subunits are held together by non-covalent interactions or by one or more disulfide bonds. The morphogenic proteins are active as dimeric proteins but are inactive as individual monomer subunits.

As a result of their biological activities, significant effort has been directed toward the development of morphogen-based therapeutics for treating injured or diseased mammalian tissue, including, for example, therapeutic compositions for inducing regenerative healing of bone defects such as fractures, as well as therapeutic compositions for preserving or restoring healthy metabolic properties in diseased bone tissue, e.g., osteopenic bone tissue. Complete descriptions of efforts to develop and characterize morphogen-based therapeutics for non-chondrogenic tissue applications in mammals, particularly humans, are set forth, for example, in: EP 0575,555; WO93/04692; WO93/05751; WO94/06399; WO94/03200; WO94/06449; WO94/10203; and WO94/06420, the disclosures of each of which are incorporated herein by reference.

Certain difficulties may be experienced upon administration of naturally isolated or recombinantly produced morphogenic proteins to a mammal. These difficulties may include, for

example, loss of morphogenic activity due to disassociation of the biologically active morphogen dimer into its inactive monomer subunits, and/or handling problems due to low solubility under physiological conditions.

Accordingly, a need remains for the identification of morphogen analogs, which mimic or
5 enhance the physiological effects of a morphogenic protein, for example OP-1. The analogs may be modified, morphogenically active hOP-1 protein dimers, or fragments or truncated analogs thereof, peptides or small organic molecules. Preferably the analogs have enhanced therapeutic value, for example, by being more stable and/or more soluble under physiological conditions than naturally occurring hOP-1, or, for example, by having enhanced tissue targeting specificity,
10 enhanced biodistribution or a reduced clearance rate in the body.

It is an object of the present invention to provide a database defining the atomic coordinates of the three-dimensional structure of mature hOP-1, all or a portion of which can be used as part of a computer system for designing and/or identifying a functional analog of hOP-1. Another object is to provide means for designing and/or identifying a molecule having enhanced
15 solubility and/or stability under physiological conditions as compared with hOP-1 and which is capable of mimicking or enhancing the biological activity of hOP-1 in a mammal. Another object of the invention is to provide a therapeutic composition comprising an analog designed and/or identified, and produced by the methods of the invention, and suitable for administration to a mammal in need thereof, such as a mammal afflicted with a metabolic bone disease, e.g., a
20 disease characterized by osteopenia. Another object of the invention is to provide methods and compositions useful for designing and/or identifying, and producing an hOP-1 antagonist capable of, for example, competing with hOP-1 for receptor binding, but incapable of inducing a receptor-mediated downstream biological effect.

These and other objects and features of the invention will be apparent from the
25 description, drawings, and claims which follow.

Summary of the Invention

The present invention is based, in part, upon the X-ray crystallographic determination of the three-dimensional structure of mature, dimeric human osteogenic protein-1 (hOP-1). The three-dimensional structure of hOP-1 has been resolved to 2.3Å. Provided herein are two sets of

atomic X-ray crystallographic co-ordinates for hOP-1, one set defining a hOP-1 structure resolved to a resolution of 2.8Å, and the other set defining a hOP-1 structure resolved to a resolution of 2.3Å. With this disclosure, the skilled artisan is provided with sets of atomic co-ordinates for use in conventional computer aided design (CAD) methodologies to identify or design protein or peptide analogs of OP-1, or alternatively, to identify or design small organic molecules that functionally mimic OP-1.

In one aspect, the invention provides a computer system comprising a memory and a processor in electrical communication with the memory. The memory has disposed therein, atomic X-ray crystallographic co-ordinates which together define at least a portion of the three-dimensional structure of hOP-1. In a preferred embodiment, the atomic co-ordinates are defined by either a portion or all of the atomic co-ordinates set forth in Figure 15 or Figure 16.

The processor, in electrical communication with the memory, comprises a process which generates a molecular model having a three-dimensional shape representative of at least a portion of human OP-1. In a preferred embodiment, the processor is capable of producing a molecular model having, in addition to the three-dimensional shape, a solvent accessible surface representative of at least a portion of human OP-1.

As used herein, the term "computer system" is understood to mean any general or special purpose system which includes a processor in electrical communication with both a memory and at least one input/output device, such as a terminal. Such a system may include, but is not limited to, personal computers, workstations or mainframes. The processor may be a general purpose processor or microprocessor or a specialized processor executing programs located in RAM memory. The programs may be placed in RAM from a storage device, such as a disk or preprogrammed ROM memory. The RAM memory in one embodiment is used both for data storage and program execution. The term computer system also embraces systems where the processor and memory reside in different physical entities but which are in electrical communication by means of a network.

In the present invention, the processor executes a modeling program which accesses data representative of the X-ray crystallographic co-ordinates of hOP-1 thereby to construct a three-dimensional model of the molecule. In addition, the processor also can execute another program,

a solvent accessible surface program, which uses the three-dimensional model of hOP-1 to construct a solvent accessible surface of at least a portion of the hOP-1 molecule and optionally calculate the solvent accessible areas of atoms. In one embodiment the solvent accessible surface program and the modeling program are the same program. In another embodiment, the modeling program and the solvent accessible surface program are different programs. In such an embodiment the modeling program may either store the three-dimensional model of hOP-1 in a region of memory accessible both to it and to the solvent accessible surface program, or the three-dimensional model may be written to external storage, such as a disk, CD ROM, or magnetic tape for later access by the solvent accessible surface program.

The memory may have stored therein the entire set of X-ray crystallographic co-ordinates which define mature biologically active human OP-1, or may comprise a subset of such co-ordinates including, for example, one or more of: a finger 1 region; a finger 2 region; and a heel region. The protein structures which correspond to the finger and heel regions are described in detail below.

In another preferred embodiment, the processor also is capable of identifying a morphogen analog, or a morphogen antagonist for example, a protein, peptide or small organic molecule, having a three-dimensional shape and preferably, in addition, a solvent accessible surface corresponding to at least a portion of human OP-1 and competent to mimic an OP-1 specific activity.

As used herein, with respect to OP-1 (or related morphogens), or with respect to a region of OP-1, the phrase "at least a portion of the three-dimensional structure of" or "at least a portion of" is understood to mean a portion of the three-dimensional surface structure of the morphogen, or region of the morphogen, including charge distribution and hydrophilicity/hydrophobicity characteristics, formed by at least three, more preferably at least three to ten, and most preferably at least ten contiguous amino acid residues of the OP-1 monomer or dimer. The contiguous residues forming such a portion may be residues which form a contiguous portion of the primary structure of the OP-1 molecule, residues which form a contiguous portion of the three-dimensional surface of the OP-1 monomer, residues which form a contiguous portion of the three-dimensional surface of the OP-1 dimer, or a combination thereof. Thus, the residues forming a portion of the three-dimensional structure of OP-1 need not be contiguous in the

- 8 -

primary sequence of the morphogen but, rather, must form a contiguous portion of the surface of the morphogen monomer or dimer. In particular, such residues may be non-contiguous in the primary structure of a single morphogen monomer or may comprise residues from different monomers in the dimeric form of the morphogen. As used herein, the residues forming "a
5 portion of the three-dimensional structure of " a morphogen, or "a portion of" a morphogen, form a contiguous three-dimensional surface in which each atom or functional group forming the portion of the surface is separated from the nearest atom or functional group forming the portion of the surface by no more than 40 Å, preferably by no more than 20 Å, more preferably by no more than 5-10 Å, and most preferably by no more than 1-5 Å.

10 As used herein the term "X-ray crystallographic co-ordinates" refers to a series of mathematical co-ordinates (represented as "X", "Y" and "Z" values) that relate to the spatial distribution of reflections produced by the diffraction of a monochromatic beam of X-rays by atoms of an hOP-1 molecule in crystal form. The diffraction data are used to generate electron density maps of the repeating units of a crystal, and the resulting electron density maps are used
15 to define the positions of individual atoms within the unit cell of the crystal.

As will be apparent to those of ordinary skill in the art, the hOP-1 structure presented herein is independent of its orientation, and that the atomic co-ordinates listed in Figs. 15 and 16 merely represent one possible orientation of the hOP-1 structure. It is apparent, therefore, that the atomic co-ordinates listed in Figs. 15 and 16, may be mathematically rotated, translated,
20 scaled, or a combination thereof, without changing the relative positions of atoms or features of the hOP-1 structure. Such mathematical manipulations are intended to be embraced herein. Furthermore, it will be apparent to the skilled artisan that the X-ray atomic co-ordinates defined herein have some degree of uncertainty in location (see, for example, column "δ" in Fig. 16 which shows the thermal uncertainty in location of each atom, as expressed in Å). Accordingly,
25 for purposes of this invention, a preselected protein or peptide having the same amino acid sequence as at least a portion of hOP-1 is considered to have the same structure as the corresponding portion of hOP-1, when a set of atomic co-ordinates defining backbone Cα atoms of the preselected protein or peptide can be superimposed onto the corresponding Cα atoms for hOP-1 (as listed in Figure 16) to a root mean square deviation of preferably less than about 1.5
30 Å, and most preferably less than about 0.75 Å.

As used herein, the term "morphogen analog", is understood to mean any molecule capable of mimicking OP-1's receptor binding activity and/or inducing a receptor mediated downstream biological effect characteristic of a morphogenic protein. Inducing alkaline phosphatase activity is a characteristic biological effect. The analog may be a protein, peptide, or non-peptidyl based organic molecule. Accordingly, the term morphogen analog embraces any substance having such OP-1 like activity, regardless of the chemical or biochemical nature thereof. The present morphogen analog can be a simple or complex substance produced by a living system or through chemical or biochemical synthetic techniques. It can be a large molecule, e.g., a modified hOP-1 dimer produced by recombinant DNA methodologies, or a small molecule, e.g., an organic molecule prepared *de novo* according to the principles of rational drug design. It can be a substance which is a mutein (or mutant protein) of hOP-1, a substance that structurally resembles a solvent-exposed surface epitope of hOP-1 and binds an OP-1 specific receptor, or a substance that otherwise stimulates an OP-1 specific receptor displayed on the surface of an OP-1 responsive cell.

As used herein, the terms "OP-1 or OP-1-like biological activity" are understood to mean any biological activities known to be induced or enhanced by OP-1. OP-1 and OP-1-like biological activities include, but are not limited to, stimulating proliferation of progenitor cells; stimulating differentiation of progenitor cells; stimulating proliferation of differentiated cells; and supporting growth and maintenance of differentiated cells. The term "progenitor cells" includes uncommitted cells, preferably of mammalian origin that are competent to differentiate into one or more specific types of differentiated cells, depending on their genomic repertoire and the tissue specificity of the permissive environment where morphogenesis is induced. Specifically, with regard to bone, cartilage, nerve, and liver tissue, the OP-1 stimulated morphogenic cascade culminates in the formation of new or regenerative differentiated tissue appropriate to the selected local environment. OP-1 mediated morphogenesis, therefore, differs significantly from simple reparative healing processes in which scar tissue (e.g., fibrous connective tissue) is formed and fills a lesion or other defect in differentiated functional tissue.

As used herein a "morphogen antagonist" is a molecule competent to mimic OP-1 receptor binding activity but which cannot induce a receptor-mediated downstream effect.

- 10 -

In yet another preferred embodiment, the processor is capable of identifying amino acids defined by the co-ordinates, which upon site-directed modification, either by chemical modification or amino acid substitution, enhance the solubility and/or stability of human OP-1.

In a related aspect, the invention provides a method of producing a morphogen analog that mimics or enhances an OP-1 or OP-1-like biological activity. The method comprises the steps of: (a) providing a molecular model defining a three-dimensional shape representative of at least a portion of human OP-1, (b) identifying a compound having a three-dimensional shape corresponding to the three-dimensional shape representative of at least the portion of human OP-1; and (c) producing the compound identified in step (b). The method can comprise the additional step of testing the compound in a biological system to determine whether the resultant candidate compound mimics or agonizes the biological activity of OP-1. It is contemplated that, in the aforementioned method, step (a) and/or (b) may be performed by means of an electronic processor using commercially available software packages.

It is contemplated that, upon determination of whether the candidate compound modulates OP-1 activity, the candidate compound can be iteratively improved using conventional CAD and/or rational drug design methodologies, well known and thoroughly documented in the art. Furthermore, it is contemplated that the resultant compound identified thus far, may be produced in a commercially useful quantity for administration into a mammal.

In another embodiment, the morphogen analog is created using atomic co-ordinates set forth in either Figs. 15 or 16. By reviewing the atomic co-ordinates set forth in Figs. 15 and 16, the skilled artisan can observe the three-dimensional structure of particular amino acid sequences located *in situ* within the three-dimensional structure of hOP-1. Preferred amino acid sequences are defined by one or more of the peptides selected from the group consisting of: H1, H-n2, H-c2, F1-2, F2-2 and F2-3, as discussed hereinbelow. The peptides provide templates which can be used in the production of more effective morphogen analogs. In a preferred embodiment, the C α atoms of amino acid residues in the morphogen analog are located within 6Å, preferably within 3Å, and most preferably within 2Å of the corresponding C α atom as defined by the respective atomic co-ordinates in Figs. 15 or 16. In another preferred embodiment, the C α atoms of amino acid residues in the morphogen analog are located within 6Å, preferably within 3Å, and most preferably within 2Å of the corresponding C α atoms of at least three amino acids in the peptide

sequences H1, H-n2, H-c2, F1-2, F2-2 and F2-3, wherein each of the C α atoms in the peptides are defined by the respective atomic co-ordinates set forth in Figs. 15 or 16.

In another embodiment, the invention provides morphogen analogs having greater solubility and/or stability in aqueous buffers than native dimeric hOP-1. In yet another
5 embodiment, the invention provides a morphogen analog which is a modified form of dimeric hOP-1, in which the modification eliminates an epitope or region on OP-1 normally recognized by an antibody or by a cellular scavenging protein for clearing OP-1 from the body.

In another embodiment, the invention provides means for creating an analog with altered receptor binding characteristics. For example, provided with the structure, charge distribution,
10 and solvent accessible surface information pertaining to the putative receptor binding site, one can alter or modify receptor binding specificity and avidity. In one embodiment, amino acid replacements in this region are made with reference to the corresponding amino acids of other known morphogens, disclosed for example, in WO 94/06449 or WO 93/05751.

After having determined the three-dimensional structure of human OP-1, a skilled artisan,
15 in possession of the atomic co-ordinates defining the OP-1 structure is hereby enabled to use conventional CAD and/or rational drug design methodologies to identify or design protein or peptide analogs, or other small organic molecules which, after having been produced using conventional chemistries and methodologies, can be tested either *in vitro* or *in vivo* to assess whether they mimic or enhance the biological activity of human OP-1.

20 The foregoing and other objects, features and advantages of the present invention will be made more apparent from the following detailed description of preferred embodiments of the invention.

Brief Description of the Drawings.

The file of this patent contains at least one drawing executed in color. Copies of this patent with color drawing(s) will be provided by the Patent and Trademark Office upon request and payment of the necessary fee.

5 The objects and features of the invention may be better understood by reference to the drawings described below, wherein like referenced features identify common features in corresponding figures.

Figure 1A is a simplified line drawing useful in describing the structure of a monomeric subunit of hOP-1. See the Summary of the Invention, *infra*, for explanation. Figures 1B, 1C,
10 and 1D are monovision ribbon tracings of the respective peptide backbones of hOP-1 finger-1, heel, and finger-2 regions. Figures 1E and 1F are schematic representations of monomeric and dimeric forms of hOP-1, respectively, as represented by a left hand motif.

Fig. 2 is a schematic drawing of a monomeric subunit of hOP-1. The hOP-1 cysteine knot comprising three disulfide bonds constitutes the core of the hOP-1 monomer subunit. Two
15 disulfide bonds which connect residues Cys 67 - Cys 136 and Cys 71-Cys 138 produce an eight residue ring through which the third disulfide bond which connects residues Cys 38 - Cys 104 passes. Four strands of antiparallel β -sheet, which emanate from the knot, form the two finger like projections. An α -helix located on the opposite end of the knot, lies perpendicular to the axis of the two fingers thereby forming the heel. The N-terminus of the monomer subunit
20 remains unresolved. The β -sheets are displayed as arrows and labeled from $\beta 1$ through $\beta 8$. The α -helix is displayed as a tube and labeled $\alpha 1$. The intra-subunit disulfide bonds that constitute the cysteine knot are shown in solid lines. Starting from Gln 36 (" N_{36} "), the first residue shown in this figure, the amino acid residues which produce secondary structure in the Finger 1 region include: Lys 39 - His 41 ($\beta 1$), Tyr 44-Ser 46 ($\beta 2$), Glu 60 - Ala 63 ($\beta 3$), Tyr 65 - Glu 70 ($\beta 4$); the
25 amino acid residues which produce secondary structure in the Finger 2 region include: Cys 103-Asn 110 ($\beta 5$); Ile 112 - Asp 118 ($\beta 6$); Asn 122 - Tyr 128 ($\beta 7$); Val 132 - His 139 ($\beta 8$); and the amino acid residues which produce secondary structure in the heel region include: Thr 82 - Ile 94($\alpha 1$).

- 13 -

Figure 3 is a structure-based sequence alignment of the hOP-1 and TGF- β 2 finger-1, heel, and finger-2 regions. Amino acid residues in the heel regions which constitute inter-chain contacts in the dimers of hOP-1 and TGF- β 2 are highlighted as white on black. Amino acid residues in the finger-1 and finger-2 regions which contact the other chain are highlighted as black on gray. In hOP-1 and TGF- β 2, the amino acids located at the same residue positions constitute the inter-chain contacts.

Figures 4A and 4B are stereo peptide backbone ribbon trace drawings illustrating the three-dimensional shape of hOP-1: A) from the "top" (down the two-fold axis of symmetry between the subunits) with the axes of the helical heel regions generally normal to the paper and the axes of each of the finger 1 and finger 2 regions generally vertical, and B) from the "side" with the two-fold axis between the subunits in the plane of the paper, with the axes of the heels generally horizontal, and the axes of the fingers generally vertical. The hOP-1 monomer has an accessible non-polar surface area of approximately 4394\AA^2 , while that for the dimer is approximately 6831\AA^2 resulting in a hidden area upon dimerization of approximately 979\AA^2 per monomer. The reader is encouraged to view the stereo alpha carbon trace drawings in wall-eyed stereo, for example, using a standard stereo viewer device, to more readily visualize the spatial relationships of amino acids sequences in the morphogen analog design.

Figure. 5A is a backbone ribbon trace drawing illustrating the hOP-1 dimer comprising the two hOP-1 monomer subunits resolved to 2.8\AA . One monomer subunit is shown in green and the other monomer subunit is shown in gold. Amino acid residues disposed within the purported receptor binding domain having solvent accessible side chains are shown as atomic spheres. The tips of the finger 1 and finger 2 regions of one OP-1 monomeric subunit and a loop at the C-terminal end of the heel of the other OP-1 monomeric subunit are believed to constitute the receptor binding domain. Amino acids located at positions of variable amino acid sequence shown in white while amino acids located at more conserved positions are shown in red. Figures 5B and 5C are pictures showing the respective solvent accessible surfaces of OP-1 and TGF- β 2 dimers colored based on their electrostatic potential. Surface regions having an electrostatic potential of -3 kT or less are shown in red while surface regions of $+3\text{ kT}$ or greater are shown in blue. Neutral regions are shown in green or gold to correspond to the backbone ribbons shown in 5A.

- 14 -

Figure 6 is a table showing an identity matrix for the TGF- β superfamily. The matrix comprises members of the TGF- β superfamily having an amino acid sequence identity relative to OP-1 of greater than 36%. In the matrix, the TGF- β superfamily members are placed in order of decreasing amino acid identity relative to OP-1. TGF- β 2 has an amino acid sequence of identity of 36% relative to OP-1 and is positioned the bottom of the matrix. Boxes enclose families of sequences having 50% or higher identity with a majority of the other members of the family; with sequences having identities of 75% or higher are shown in gray. Recombinantly expressed OP/BMP family members which have been shown to make bone are denoted by a "+" in the left margin. In the left margin, TGF- β superfamily members with three-dimensional structures determined are highlighted white on black. The sequences are referenced in Kingsley (Kingsley. (1994) *Genes and Development* 8:133-146), except for the following: UNIVIN (Stenzel *et al.* (1994) *Develop. Biol.* 166:149-158.), SCREW (Arora, *et al.* (1994) *Genes and Dev.* 8:2588-2601.), BMP-9 (Wozney, *et al.* (1993) PCT/WO 93/00432, SEQ. ID. NO.9), BMP-10 (Celeste *et al.* (1994) PCT/WO 94/26893, SEQ. ID. NO. 1), GDF-5 (Storm *et al.* (1994) *Nature* 368:639-643) (also called CDMP-1 (Chang *et al.* (1994) *J. Biol. Chem.* 269: 28227-28234.), GDF-6 (Storm, *et al.* (1994) *Nature* 368:639-643), GDF-7 (Storm *et al.* (1994) *Nature* 368:639-643), CDMP-2 (Chang *et al.* (1994) *J. Biol. Chem.* 269: 28227-28234.), OP-3 (Özkaynak *et al.* (1994) PCT/WO 94/10203, SEQ. ID. NO. 1), Inhibin β c (Hötten, *et al.* (1995) *Bioch. Biophys. Res. Comm.* 206:608-613), and GDF-10 (Cunningham, *et al.* (1995) *Growth Factors* 12:99-109.).

The disclosures of the aforementioned citations are incorporated herein by reference. Several sequences in the matrix have alternate names: OP-1 (BMP-7), BMP-2 (BMP-2a), BMP-4 (BMP-2b), BMP-6 (Vgr1), OP-2 (BMP-8), 60A (Vgr-D), BMP-3 (osteogenin), GDF-5 (CDMP-1, MP-52), GDF-6 (CDMP-2, BMP-13) and GDF-7 (CDMP-3, BMP-12).

Figure 7 is a summary of amino acid residues which, according to the 2.8Å resolution structure, together define the solvent accessible surfaces of dimeric hOP-1. Figures 7A, 7B, and 7C show the amino acid sequences defining the human OP-1 finger 1, heel, and finger 2 regions, respectively. The amino acid residues having 40% or greater of their sidechain exposed to solvent are boxed, wherein the solvent accessible amino acid residues that are highly variable among the BMP/OP family of the TGF- β superfamily are identified by shaded boxes.

Figure 8 is a table, based on the 2.8Å structure, which summarizes the percentage surface accessibility of the amino acid side chains in a hOP-1 monomer subunit and in a hOP-1 dimer. Amino acid residues believed to constitute putative epitopes are designated "EPITOPE" and amino acid residues which are potential candidates as surface modifiable amino acids are marked with an asterisk. In addition, surface modifiable amino acids which are preferred candidates for enhancing solubility are marked with an asterisk.

Figure 9 is a table, based on the 2.8Å structure, which summarizes amino acid residues believed to define the ridge. Amino acid residues believed to constitute the receptor binding domain in the ridge are marked with an asterisk.

Figure 10 is a schematic representation of a computer system useful in the practice of the invention.

Figures 11A and 11B are tables, produced by reference to the 2.8Å structure, which summarize amino acid pairs believed to be useful as sites for introducing additional inter-chain (11A) or intra-chain (11B) disulfide bonds in the hOP-1 dimer.

Figure 12 is an amino acid sequence alignment showing the amino acid sequence of mature human OP-1, and peptides defining the finger-1, finger-2 and heel regions of human OP-1.

Figures 13A-13D are bar graphs illustrating the effect of finger-2 and heel peptides on the alkaline phosphatase activity of ROS cells incubated in either the presence or absence of soluble OP-1. Figures 13A, 13B, 13C, and 13D show the effect of peptides F2-2, F2-3, Hn-2 and Hn-3, respectively, on the alkaline phosphatase activity of ROS cells incubated in the presence (shaded bars) or absence of soluble OP-1 (unshaded bars).

Figures 14A and 14B are graphs showing the displacement of radiolabelled soluble OP-1 from ROS cell membranes by finger 1, finger 2, and heel peptides. Figure 14A shows the displacement of radiolabelled OP-1 from ROS cell membranes by unlabeled soluble OP-1 (open circles and triangles), finger 2 peptide F2-2 (closed circles) and finger 2 peptide F2-3 (closed triangles). Figure 14B shows the displacement of radiolabelled OP-1 from ROS cell membranes by unlabeled soluble OP-1 (open triangles), finger 1 peptide F1-2 (closed boxes), heel peptide Hn2 (open diamonds) and heel peptide H-c2 (open circles).

- 16 -

Figure 15 is a table summarizing the atomic co-ordinates of hOP-1 resolved to 2.8Å.

Figure 16 is a table summarizing the atomic co-ordinates of hOP-1 resolved to 2.3Å.

Further particulars concerning the drawings are disclosed in the following description which discloses details of the three-dimensional structure of hOP-1, methods for identifying morphogen
5 analogs, and methods for making, testing and using such morphogen analogs.

Detailed Description of Preferred Embodiments

I. Introduction

As described hereinbelow, the three-dimensional crystal structure of mature hOP-1 now has been solved to 2.3Å. The disclosure provides two sets of atomic co-ordinates for hOP-1, wherein one set of co-ordinates (see Fig. 15) represents the structure of hOP-1 resolved to 2.8Å, and the other set of co-ordinates (see Fig. 16) represents the structure of hOP-1 resolved to 2.3Å. This disclosure thus provides, the atomic co-ordinates defining the relative positions, in three-dimensional space, of at least the C-terminal 104 amino acids of human OP-1 which are sufficient for imparting biological activity. The disclosure provides also an analysis of the structural features of hOP-1. The skilled artisan now can use some or all of these co-ordinates in a database for making morphogenic protein analogs, particularly OP-1 analogs. Specifically, the artisan can select part or all of the database to create templates of part, or all of the hOP-1 structure in three-dimensions, and using this template, create a desired analog or variant which may be amino acid-based, or alternatively composed, in whole or in part, by non-amino acid-based organic components.

Provided below is a detailed description of the three-dimensional crystal structure of hOP-1, along with a detailed description on how to use co-ordinates in a database to design a morphogen analog or structural variant of interest. Amino acid sequences as exemplary templates are provided as examples for designing, identifying, and producing an OP-1 analog using one of the OP-1 atomic co-ordinate databases. Specifically contemplated herein as useful analogs include: small amino molecules which mimic the receptor binding region of the protein; analogs having enhanced stability or solubility; analogs having reduced clearance rates from the body; or enhanced target tissue specificity. The reader will appreciate that these examples are merely exemplary. Given the disclosure of the co-ordinates, the three-dimensional structure, the use of the co-ordinates in a database, and the level of skill in the art today, still other analogs, not specifically recited herein, are contemplated and enabled by this disclosure. In particular, it will be appreciated that, given the disclosure herein, and the known amino acid sequences for other, closely related morphogens, the methods can be used to create other morphogen analogs of, for example, BMP2, BMP4, OP2, BMP5 and BMP6.

II. Structural Determination of hOP-1

A. Determination of the 2.8Å Structure

Crystals of mature hOP-1 were grown by mixing equal volumes of purified protein (Özkaynak *et al.* (1990) *EMBO J.* 9:2085-20893; and Sampath *et al.* (1992) *J. Biol. Chem.* 267:20352-20362) at 10 mg/ml, with 8% saturated ammonium sulfate in 50 mM sodium acetate buffer (pH 5.0) (Griffith *et al.* (1994) *J. Mol. Biol.* 244:657-658). The crystals have the symmetry of space group $P3_221$ with unit cell dimensions $a=b=99.46\text{\AA}$, and $c=42.09\text{\AA}$. One crystal was used to collect a complete native data set to 2.8Å resolution at 4°C. Two heavy atom derivative data sets were collected at 4°C, one from a crystal soaked for seven days in 0.3mM uranyl nitrate and the other from a crystal soaked for eight hours in 0.5 mM sodium gold (III) tetra chloride (Griffith *et al.* (1994) *supra*).

The native and derivative data sets were integrated and reduced with the R-AXIS-IIC software suite (Higashi (1990) A Program for Indexing and Processing R-AXIS IIC Imaging Plate Data, Rigaku Corp.) and scaled together using the CCP4 program ANSC (Collaborative Computation Project (1994) *Acta Cryst. D50*:760-763). Inspection of the Harker sections of the difference Patterson map reveals a single uranyl site. The position of the single gold site was determined by using cross-Fourier techniques using the uranyl position as the phasing site. The heavy atom x,y,z parameters and occupancy were refined with the program TENEYCK (Ten Eyck *et al.* (1976) *J. Mol. Biol.* 100:3-11). Using these two derivatives and their anomalous signals, an initial phase set was calculated to 4.0Å resolution with a mean figure of merit of 0.72. The phases were improved and extended to 3.5Å resolution by cycles of solvent flattening (Wang (1985) *Meth. Enzymol.* 115:90-112) and phase combination (Reed (1986) *Acta Cryst. A42*:140-149) using the CCP4 (Collaborative Computation Project (1994) *supra*) crystallographic package. A completely interpretable 3.5Å resolution electron density map permitted the unambiguous tracing of the polypeptide chain and identification of the amino acids from Gln 36 to His 139 using the graphic program "O" (Jones *et al.* (1991) *Acta Crystallogr. A47*:110-119). The model was refined with the program XPLOR (Brunger *et al.* (1987) *Science* 235:458-460) by using all reflections between 10Å and 2.8Å resolution for which $F_{\text{obs}} > 2.0\sigma$ (F_{obs}). There were no water molecules included in the refinement. The root mean square (rms)

deviation from ideality is 0.02 Å for bond lengths, 3.2° for bond angles. Good stereochemistry was observed for backbone torsion angles. The current R factor is 22.8%.

The atomic co-ordinates defining the 2.8Å resolution structure are listed in Fig. 15. In Fig. 15, the columns entitled "Atom" denote atoms whose co-ordinates have been measured.

- 5 The first letter in the column defines the element. The columns entitled "Residue" denote the amino acid residues in the hOP-1 monomer which contain an atom whose co-ordinates have been measured. The column entitled "Chain" denotes whether the atom of interest is located within the first (A) or second (B) monomer subunit of the hOP-1 dimer. The columns "X, Y, Z" are the Cartesian co-ordinates that define the atomic position of the atom measured.

10 B. Determination of the 2.3Å Structure

- Crystals of mature hOP-1 were produced as described in the previous section. One crystal, frozen in liquid nitrogen, was used to collect a data set to 2.3Å resolution that was 91% complete. The data were collected on imaging plates at beam line X12C (National Synchrotron Light Source) with an oscillation range of 0.5 degrees (overlap of 0.1 degrees) and exposure times of 60-90 seconds. The digitalized data were processed, merged and scaled with DENZO and SCALEPACK (available from Molecular Structure Corporation, Texas). An initial $2Fo-FC$ map, calculated after X-PLOR rigid-body refinement using the 2.8Å model, was readily interpretable. Portions of the model were manually refitted to the electron-density map with the interactive graphics programs "O" and "Chain". Subsequent cycles of refinement (XPLOR/PROFFT) and manual rebuilding (QUANTA) rapidly converged to the present model.
- 15
20

- The current model yielded a conventional crystallographic R factor of 23.5% for data from 10 to 2.3Å (1.5σ cutoff) and a R_{free} of 27%. The refined structure was analyzed using the PROCHECK (available from Protein Data Bank, Brookhaven, NY) algorithm and corrected where appropriate. The root mean square (rms) deviation from ideality is 0.015Å for bond distances, 0.034Å for angle distances, and 0.142Å for planar 1-4 distances. The rms deviation from ideality is 1.7° for bond angles. The upper estimate of the error in the atomic positions from the Luzzati plots (EXPLOR) using the free R factor is 0.25-0.33Å. The final model, comprising one monomer subunit, consists of 828 protein atoms (i.e., all non-hydrogen atoms)
- 25

- 20 -

and 33 water molecules. The average temperature (B) factor is 33\AA^2 for protein atoms and 37\AA^2 for solvent atoms.

The atomic co-ordinates defining the 2.3\AA resolution structure are listed in Figure 16. In Fig. 16, the columns entitled "Atom" denote atoms whose co-ordinates have been measured.

- 5 The first letter in the column defines the element. The columns entitled "Residue" denote the amino acid residues in the hOP-1 monomer which contain an atom whose co-ordinates have been measured. The column entitled "Chain" denotes whether the atom of interest is located within the first (A) or second (B) monomer subunit of the hOP-1 dimer. The columns "X, Y, Z" are the Cartesian co-ordinates that define the atomic position of the atom measured. The column
- 10 denoted " δ " represents the uncertainty in the position of the co-ordinate as derived from the temperature factor (B) of each corresponding atom. The uncertainty of each co-ordinate was derived from the formula $\delta = \sqrt{\frac{B}{8\pi^2}}$ (see "*Protein Crystallography*" (1976) T.L. Blundell and L.N. Johnson, Academic Press, p. 121) and is expressed in units of \AA .

III. Structural Features of hOP-1 Monomer Subunits

- 15 Human OP-1, like TGF- β 2, is a dimeric protein having a unique folding pattern involving six of the seven C-terminal cysteine residues, as illustrated in Figure 1A. Each of the subunits in OP-1, like TGF β 2 (See Daopin *et al.* (1992) *Science* 257:369-373; and Schulnegger *et al.* (1992) *Nature* 358:430-434) have a characteristic folding pattern, illustrated schematically in Fig. 1A, that involves six of the seven C-terminal cysteine residues.

- 20 Referring to Fig. 1A, four of the cysteine residues in each subunit form two disulfide bonds which together create an eight residue ring, while two additional cysteine residues form a disulfide bond that passes through the ring to form a knot-like structure (cysteine knot). With a numbering scheme beginning with the most N-terminal cysteine of the 7 conserved cysteine residues assigned number 1, the 2nd and 6th cysteine residues are disulfide bonded to close one
- 25 side of the eight residue ring while the 3rd and 7th cysteine residues are disulfide bonded to close the other side of the ring. The 1st and 5th conserved cysteine residues are disulfide bonded through the center of the ring to form the core of the knot. Amino acid sequence alignment patterns suggest this structural motif is conserved between members of the TGF- β superfamily.

The 4th cysteine is semi-conserved and when present typically forms an inter-chain disulfide bond (ICDB) with the corresponding cysteine residue in the other subunit.

Each hOP-1 monomer subunit comprises three major tertiary structural elements and an N-terminal region. The structural elements are made up of regions of contiguous polypeptide chain that possess over 50% secondary structure of the following types: (1) loop, (2) helix and (3) β -sheet. Furthermore, in these regions the N-terminal and C-terminal strands are not more than 7 Å apart.

The amino acid sequence between the 1st and 2nd conserved cysteines (Fig. 1A) form a structural region characterized by an anti-parallel β -sheet finger, referred to herein as the finger 1 region (F1). A ribbon trace of the human OP-1 finger 1 peptide backbone is shown in Fig. 1B. Similarly the residues between the 5th and 6th conserved cysteines in Fig. 1A also form an anti-parallel β -sheet finger, referred to herein as the finger 2 region (F2). A ribbon trace of the human OP-1 finger 2 peptide backbone is shown in Fig. 1D. A β -sheet finger is a single amino acid chain, comprising a β -strand that folds back on itself by means of a β -turn or some larger loop so that the entering and exiting strands form one or more anti-parallel β -sheet structures. The third major structural region, involving the residues between the 3rd and 4th conserved cysteines in Fig. 1A, is characterized by a three turn α -helix referred to herein as the heel region (H). A ribbon trace of the human OP-1 heel peptide backbone is shown in Fig. 1C.

The organization of the monomer structure is similar to that of a left hand (see Fig. 1E) where the knot region is located at the position equivalent to the palm (16), the finger 1 region is equivalent to the index and middle fingers (12 and 13, respectively), the α -helix, or heel region, is equivalent to the heel of the hand (17), and the finger 2 region is equivalent to the ring and small fingers (14 and 15, respectively). The N-terminal region (undefined in the 2.8 Å resolution map disclosed herein) is predicted to be located at a position roughly equivalent to the thumb (11).

Monovision ribbon tracings illustrating the alpha carbon backbones of each of the three major independent structural elements of the monomer are illustrated in Figures 1B-1D. Specifically, the finger 1 region comprising the first anti-parallel β -sheet segment is shown in

- 22 -

Fig. 1B, the heel region comprising the three turn α -helical segment is shown in Fig. 1C, and the finger 2 region comprising second and third anti-parallel β -sheet segments is shown in Fig. 1D.

For the sake of comparison, Fig. 3 shows an alignment of the amino acid sequences defining the finger 1, finger 2 and heel regions of hOP-1 and TGF- β 2. In Fig. 3, the OP-1 and TGF- β 2 amino acid sequences were aligned according to the corresponding regions of local structural identity in the OP-1 and TGF- β 2 structures. Alignment gaps were positioned in loop regions, which is where the local conformational homology of the α -carbon traces tends to be the lowest.

The structure-based alignment of OP-1 and TGF- β 2 then was used as a template for the alignment of the 7-cysteine domain sequences of other TGF- β superfamily members (other members of the TGF- β superfamily are set forth in Fig. 6). Alignment gaps were positioned in regions which are loops in both the OP-1 and TGF- β 2 structures. Percent identity between pairs of sequences was calculated as the number of identical aligned sequence positions, excluding gaps, normalized to the geometric mean of the lengths of the sequences and multiplied by 100.

Fig. 6 is a matrix of the resulting pair wise percent identities between super family sequences so aligned. Using such principles, it is contemplated that the hOP-1 and TGF- β 2 structures, either alone or in combination, may be used for homology modeling of other proteins belonging to the TGF- β superfamily whose three-dimensional structures have not yet been determined (see, for example, the other members of the TGF- β superfamily listed in Figure 6). It is contemplated that such models may be useful in designing morphogen analogs for the particular candidate morphogens of interest, however, for simplicity, the disclosure hereinbelow refers specifically the design, identification, and production of morphogen analogs of hOP-1.

Fig. 3 also shows, based on an analysis of the 2.8Å resolution structure, a comparison of interchain contact residues in OP-1 and TGF- β 2. Residues were designated as contact residues if the distance between the centers of at least one non-hydrogen atom from each side chain was less than the sum of their Van der Waals radii plus 1.1Å. Despite the low level of sequence identity between OP-1 and TGF- β 2, the inter chain contacts between residues in the heel of one chain and residues in finger 1 and finger 2 of the other chain are well conserved.

Upon detailed inspection of the 2.8Å resolution structure of hOP-1, the finger 1 region of hOP-1 is an antiparallel β -sheet containing a thirteen residue omega loop (Phe 47-Glu 60) (Fig. 2). The structural alignment of the OP-1 and TGF- β 2 sequences in Fig. 3 places two gaps in the omega loop. The first gap represents a deletion in hOP-1 that aligns with Arg 26 in the α 2 helix of TGF- β 2. This deletion results in a tighter, non- α -helical turn in OP-1 as compared with TGF- β 2. The second gap corresponds to the insertion of Gln 53 in OP-1, which has the result of directing both Gln 53 and Asp 54 side chains into the solvent. By comparison, in the corresponding region of TGF- β 2, only Lys 31 is in contact with the solvent. These differences in the conformation of the omega loop also result in the conserved proline (Pro 59) adopting a trans conformation in hOP-1 rather than cis, as in TGF- β 2. The conformation of the omega loop orients six non-polar residues so they can contribute to a solvent inaccessible interface with Finger 2. Of these six, four are aromatic (Phe 47, Trp 55, Tyr 62 and Tyr 65), and two are aliphatic (Ile 56 and Ile 57). In all, the conformation of the omega loop backbone places five polar residues (Arg 48, Asp 49, Gln 53, Asp 54, and Glu 60) in contact with solvent. The net surface charge in this region is -2 whereas it is +2 for TGF- β 2 (Fig. 5).

According to the 2.8Å structure, the only α helix in the monomer is located between the third and fifth cysteines (Cys 71 and Cys 104). This helix extends for three and one-half turns from residues Thr 82 to Ile 94, is amphipathic, and contains a number of hydrophobic residues which in the dimer make contact with residues from Finger 1 and Finger 2 of the other monomer (Fig. 3). Several hydrophilic residues (Thr 82, His 84, and Gln 88) form one wall of an internal solvent pocket near the 2-fold axis of the dimer, while others (Asn 83, His 92, and Asn 95) are in contact with the external solvent. The conformation of the loop leading from the C-terminal end of the helix back to the cysteine knot is similar in OP-1 and TGF- β 2. By comparison, the loop located at the N-terminal end of the helix is 3 residues longer in OP-1, resulting in a different fold than in TGF- β 2. In this loop of OP-1, it is believed that an N-linked sugar moiety is attached to Asn 80, however, no such corresponding glycosylation site exists in TGF- β 2. Further, this loop is uncharged in OP-1 whereas it is negatively charged in TGF- β 2.

According to the 2.8Å structure, Finger 2 is the second antiparallel β -sheet in OP-1 (Fig. 2). The polypeptide chain reverses direction between segments β 6 and β 7 through a 3:5 turn (Sibanda, *et al.* (1991) *Methods in Enzymol.* 202:59-82) beginning at residue Asp 118 and ending

at residue Asn 122. In contrast, TGF- β 2 has one less residue in this loop and adopts a 2:2 turn (Sibanda *et al.* (1991) *supra*). Residues Arg 129 to Val 132, located between segments β 7 and β 8, form a peptide bridge that crosses over the C-terminal end of strand β 5 and produces a 180° twist in the Finger 2 antiparallel β -structure. A similar structure is observed in other cysteine knot growth factors, however the peptide bridge length varies (McDonald *et al.* (1991) *Nature* 354:411-414). Within the monomer, Finger 2 makes intra-chain contacts with Finger 1 by contributing aromatic residues Tyr 116, Phe 117 and Tyr 128, and aliphatic residues Val 114, Leu 115, Val 123, Met 131 and Val 133 to a solvent inaccessible interface. OP-1 and TGF- β 2 differ by three charges in the region of the Finger 2 turn; OP-1 has two negative charges while TGF- β 2 has one positive charge. In the region between the turn and the peptide bridge, OP-1 has a net charge of +3 while TGF- β 2 is neutral (Fig. 5).

The N-terminus of each monomeric subunit is believed to be highly mobile and has not been resolved in the 2.8Å resolution structure of hOP-1. The N-terminal region can be deleted without affecting biological activity and, therefore, it is contemplated that this portion of mature hOP-1 may be removed and replaced with other protein or peptide sequences, such as antibodies, and/or radiolabel binding sites for enhancing targeting to a particular locus *in vivo* or for use in *in vivo* imaging experiments. In addition, the N-terminal region may be replaced with an ion chelating motif (e.g., His₆) for use in affinity purification schemes, or replaced with proteins or peptides for enhancing solubility in aqueous solvents.

IV. Structural Features of the hOP-1 Dimer

Fig. 4 shows stereo ribbon trace drawings representative of the peptide backbone of the hOP-1 dimer complex, based on the 2.8Å structure. The two monomer subunits in the dimer complex are oriented symmetrically such that the heel region of one subunit contacts the finger regions of the other subunit with the knot regions of the connected subunits forming the core of the molecule. The 4th cysteine forms an inter-chain disulfide bond with its counterpart on the second chain thereby equivalently linking the chains at the center of the palms. The dimer thus formed is an ellipsoidal (cigar shaped) molecule when viewed from the top looking down the two-fold axis of symmetry between the subunits (Fig. 4A). Viewed from the side, the molecule resembles a bent "cigar" since the two subunits are oriented at a slight angle relative to each other (Fig. 4B).

As shown in Fig. 4, each of the structural elements which together define the native monomer subunits of the dimer are labeled 43, 43', 44, 44', 45, 45', 46, and 46', wherein, elements 43, 44, 45, and 46 are defined by one subunit and elements 43', 44', 45', and 46' belong to the other subunit. Specifically, 43 and 43' denote the finger 1 regions; 44 and 44' denote heel regions; 45 and 45' denote the finger 2 regions; and 46 and 46' denote disulfide bonds which connect the 1st and 5th conserved cysteines of each subunit to form the knot-like structure. From Fig. 4, it can be seen that the heel region from one subunit, e.g., 44, and the finger 1 and finger 2 regions, e.g., 43' and 45', respectively from the other subunit, interact with one another. These three elements are believed to co-operate with one another to define a structure interactive with the ligand binding interactive surface of the cognate receptor.

The helical axis is defined as the line equi-distant from the alpha carbons in the helical region. A sequence of four points is needed to define the dihedral angle between the axes of the helices in the dimer. The two inner points were chosen to lie on the helical axes adjacent to the α -carbon of residue His 84 in OP-1 or His 58 in TGF- β 2, respectively. The two outer points were chosen to lie on their respective helical axes, but their location is arbitrary. To measure the angle between the helices, the first two points used to define the dihedral angle were translated so as to superimpose the inner points. The resulting three points define the angle.

A major difference between the OP-1 and TGF- β 2 dimers is the relative orientation of the helices in the heel region. The angle between the axes of the helices in the heel region of OP-1 is 43° which is 10° larger than that measured for TGF- β 2. The measured dihedral angle between the helices is -20° for OP-1 which is 14° more negative than for TGF- β 2. Despite these differences in helical orientation, the same helix and finger residue positions are involved in making inter-chain contacts, as evidenced by the shaded residues in Fig. 3.

A. Differences in the hOP-1 Dimer Relative To Individual Monomer Subunits

During dimerization of the monomer subunits, several amino acids on the surface of each monomer subunit become buried in the hOP-1 dimer. Figure 8 highlights differences in the surface accessibility of particular amino acid residues located in the hOP-1 monomer subunit relative to those in the hOP-1 dimer, as determined from the 2.8 Å structure.

Loss of non-polar surface area during dimerization was calculated using ACCESS (version 2.1) with a 1.4Å probe (Lee *et al.* (1971) *J. Mol. Biol.* 55:379-400). Non-polar surface area is defined as the contribution to the total accessible surface from carbon and sulfur atoms. The surface area measurement algorithm in ACCESS slices the structure into 0.25Å slabs
5 perpendicular to the Z-axis. As a consequence, the results are sensitive to the orientation of a structure relative to the Z-axis (Lee *et al.* (1971) *supra*). In order to minimize this effect, we evaluated three perpendicular and one intermediate orientations of each structure. The results of these calculations were combined by accepting, for each non-polar atom, the largest accessible area measured among the four orientations. The values for TGF-β2 reported here were
10 calculated using coordinates from entry 2TG1 (Daopin *et al.* (1992) *supra*) and entry 1TFG (Schlunegger *et al.* (1992) *supra*) obtained from the January 1994 release of the Protein Data Bank (Bernstein *et al.* (1977) *J. Mol. Biol.* 112:535-542) at Brookhaven National Laboratory.

In Figure 8, the column entitled "Residue" denotes an amino acid of interest. The column entitled "Monomer % Area" denotes the percentage of the amino acid that is exposed on the
15 surface of the hOP-1 monomer, the column entitled "Dimer % Area" denotes the percentage of the amino acid that is exposed on the surface of the hOP-1 dimer, and the column denoted "Hidden % Area" denotes amount of surface area for each amino acid that is lost upon dimerization of each monomer subunit to produce the hOP-1 dimer. This analysis reveals amino acids which become buried during dimerization and, thus, likely are located at the interface of
20 the two monomer subunits. For example, 70.75% of the surface area of His 84 becomes hidden upon dimerization. A review of the structure of dimeric hOP-1 reveals that His 84 is located at the interface between the two monomers.

B. Solution Electrostatic Potentials on the Surface of OP-1 and TGF-β2

The solution electrostatic potentials surrounding the OP-1 and TGF-β2 (1TFG)
25 (Schlunegger *et al.* (1992) *supra*) dimers were calculated using DELPHI (Gilson *et al.* (1987) *Nature* 330:84-86; and Nicholls *et al.* (1991) *J. Comput. Chem.* 12:435-445) (Biosym Technologies, Inc., San Diego, CA). The calculations were performed using a solvent dielectric constant of 80, a solvent radius of 1.4Å, an ionic strength of 0.145M and an ionic radius of 2.0 Å. The interior of the protein was modeled using a dielectric constant of 2.0. Formal charges were
30 used and distributed as follows: atoms OD1 and OD2 of Asp were each charged -0.5, atoms

OE1 and OE2 of Glu were each charged -0.5, atoms ND1 and NE2 of His were each charged 0.25, atom NZ of Lys was charged +1.0, atoms NH1 and NH2 of Arg were each charged +0.5, and atom OXT of the C-terminal carboxyl group was charged -1.0.

The differences in charge distribution on the surfaces of OP-1 and TGF- β 2 can be
5 observed by comparing the color distributions of Figs. 5B and 5C, respectively. Surface regions having an electrostatic potential of -3kT or less are shown in red while surface regions of +3kT or greater are shown in blue. Neutral regions are shown in green or gold to correspond to the backbone ribbons shown in Fig. 5A. As mentioned in the following section, the differences in electrostatic potential on the surfaces of OP-1 and TGF- β 2 may play an important role in the
10 specific interactions of the TGF- β superfamily members with their cognate receptors.

C. Receptor Binding Domain

Without wishing to be bound by theory, it is contemplated that the receptor binding regions of hOP-1 includes amino acids that are both solvent accessible and lie at positions of heterogeneous composition, as determined from the amino acid sequence of hOP-1 when aligned
15 with other members of the TGF- β superfamily (See Fig. 3). Divergent structural features in hOP-1, like TGF- β 2, occur primarily in the external loops of finger 1 and finger 2, the loops bordering the helix in the heel region, and the residues in the N-terminal domain preceding the first cysteine of the cysteine knot. These regions are solvent accessible. In both the OP-1 and TGF- β 2 dimer structures, the tip of finger 2 and the omega loop of finger 1 from one chain, and
20 the C-terminal end of the α -helix in the heel of the other chain form a contiguous ridge approximately 40 Å long and 15 Å wide (Fig. 5A). It is contemplated that this ridge contains the primary structural features that interact with the cognate receptor, and that the binding specificity between different TGF- β superfamily members derives from conformational and electrostatic variations on the surface of this ridge.

25 Differences in the conformation of the finger 1 omega loop, which constitutes the mid section of the ridge, and in the turn at the end of finger 2, which forms one end of the ridge are noted. However, there are striking differences in the surface charge of the ridge in hOP-1 relative to TGF- β 2 (see Figs. 5B and 5C). In hOP-1, the ends of the finger regions are negatively charged whereas in TGF- β 2, the ends of the finger regions are positively charged. This results in

- 28 -

a net charge of -4 for the receptor binding ridge of hOP-1 versus +3 for TGF- β 2. Conversely, the β -strand located C-terminal to the turn of finger 2 (β 7, Fig. 2) is positively charged in OP-1 whereas it is negatively charged in TGF- β 2 (Figs. 5B and 5C). These features suggests that electrostatic charge distribution plays an important role in the specific interactions of the TGF- β superfamily members with their cognate receptors.

Figure 9 summarizes the amino acid residues which, according to the 2.8 Å structure, are believed to constitute the ridge, and also indicates whether each amino acid residue is disposed within the heel, finger 1, or finger 2 domains. Figure 9 also provides a list of amino acid residues which are believed to constitute at least part, if not all of the receptor binding domain of hOP-1.

V. Design of Morphogen Analogs

Although it is contemplated that the design of morphogen analogs can be facilitated by conventional ball and stick type modeling procedures, it is contemplated that the ability to design morphogen analogs is enhanced significantly using modern computer-driven modeling and design procedures.

It is contemplated that the design of morphogen analogs, as discussed in detail hereinbelow, is facilitated using conventional molecular modeling computers or workstations, commercially available from, for example, Silicon Graphics, Inc. or Evans and Sutherland Computer Corp., which implement equally conventional computer modeling programs, for example, INSIGHTII, DISCOVER, and DELPHI, commercially available from Biosym, Technologies Inc., and QUANTA, and CHARMM commercially available from Molecular Simulations, Inc.

Furthermore, it is understood that any computer system having the overall characteristics set forth in Fig. 10 may be useful in the practice of the instant invention. More specifically, Fig. 10, is a schematic representation of a typical computer work station having in electrical communication (100) with one another via, for example, an internal bus or external network, a processor (101), a RAM (102), a ROM (103), a terminal (104), and optionally an external storage device, for example, a diskette, CD ROM, or magnetic tape (105).

It is contemplated, that the co-ordinates can be used not only to provide a basis for re-engineering hOP-1 dimers by using, for example, site-directed mutagenesis methodologies, to enhance, for example, the solubility and or/stability of the active hOP-1 dimer in physiological buffers, but also to provide a starting point for the *de novo* design and production of peptides or other small molecules which mimic the bioactivity of hOP-1. Set forth below are illustrative examples demonstrating the usefulness of hOP-1 atomic co-ordinates in the design of morphogen analogs, however, it is understood the examples below are illustrative and not meant to be limiting in any way.

A. Engineering hOP-1 Dimers

In one aspect, the availability of the atomic co-ordinates for hOP-1, enables the artisan to perform theoretical amino acid replacements and to determine by calculation, in advance of actually making and testing the candidate molecule in a laboratory setting, whether a particular amino acid substitution disrupts the packing of the OP-1 dimer and whether a morphogen analog is likely to be more stable and/or soluble than the template OP-1 molecule. Such procedures assist the artisan to eliminate non viable replacements and to focus efforts on more promising candidate analogs.

(i) Enhancing the Stability of hOP-1 Dimers

It is contemplated that the skilled artisan in possession of the atomic co-ordinates defining hOP-1 can introduce additional inter- or intra-chain covalent and/or non-covalent interactions into the hOP-1 dimer to stabilize the dimer by preventing disassociation or unfolding of each monomer subunit. Preferred engineered covalent interactions include, for example, engineered disulfide bonds, and preferred engineered non-covalent interactions include, for example, hydrogen bonds, salt bridges, and hydrophobic interactions.

For example, in order to introduce additional disulfide bonds, the skilled artisan can identify sites suitable for the introduction of a pair of cysteine amino acid residues by using standard molecular modeling programs, for example, INSIGHT, DISCOVER, CHARMM and QUANTA. Another program useful in identifying pairs of amino acids as potential sites for introducing stabilizing disulfide bonds is described in U.S. Patent No. 4,908,773, the disclosure of which is incorporated herein by reference.

- 30 -

For example, the skilled artisan using the INSIGHT program can screen for pairs of amino acids, where the distance between the C β atoms of each amino acid is in the range of about 3.0 to about 5.0, or more preferably about 3.5 to about 4.5 Å apart. For this purpose, glycines, which contain no C β -C β bond, are first converted to alanines on the computer. The possible range of C β -C β distances in a disulfide bond are 3.1 Å to 4.6 Å, but separations outside this range can be accommodated by small shifts in the neighboring atoms. Searching C β , rather than C α distances, ensures both reasonable spacing as well as proper orientation of the C α -C β bond. The effects of adding such an additional linkage on protein structure are determined by mutating the two candidates residues to Cys; rotating each new Cys about the C α -C β bond to bring the two γ sulfurs as close to within 2Å as possible; creating a disulfide between the γ sulfurs; and energy minimizing structural regions within 5 Å of the disulfide bond. Any deformation of the structure caused by introduction of the additional disulfide bond is revealed by inspection when the minimized, mutated model structure is superimposed on the native structure.

It is contemplated that the introduction of additional linkages will improve solubility by preventing transient exposure of non-polar interface or buried residues. Figure 11A lists amino acid residues, based on the 2.8Å structure, which may be mutated to cysteine residues for introducing additional inter-chain disulfide bonds, based upon the selection criteria presented above. For reference purposes, Table 11A includes also the length of the naturally occurring inter-chain disulfide linkage in wild type hOP-1, that is, the disulfide linkage connecting Cys-103 of one monomer subunit with the counterpart Cys-103 of the other monomer subunit.

A preferred pair of residues suitable for modification include the residue at position 83 of one chain and the residue at position 130 of the other chain. It is contemplated that the additional inter-chain linkage stabilizes the dimeric structure by connecting the N-terminal end of the Heel helix of the first subunit to the middle of the Finger 2 region in the second subunit. A disulfide bond between position 82 on one chain and position 130 of the other chain also is geometrically feasible, but because Thr 82 is part of the NAT glycosylation site in OP-1, its modification may inhibit proper glycosylation.

Figure 11B summarizes amino acid residues which can be mutated to cysteine residues for introducing additional intra-chain disulfide bonds, based upon the selection criteria presented

above. As noted previously, the putative receptor binding region comprises at least two physically proximal, but sequentially separate regions, namely the tips of Finger 1 and Finger 2. It is contemplated that the structural integrity of the putative receptor binding ridge can be stabilized by engineering an intra-chain disulfide link between residues of Finger 1 and Finger 2.

5 In a preferred embodiment, the residue at position 58 in Finger 1 can be disulfide bonded with the residue at position 114 in Finger 2. It is contemplated that a link between the residues at positions 58 and 115 also would be viable, however, this would move the disulfide bond nearer to the putative receptor binding region on Finger 2. Also a link between positions 65 and 133 is possible, however, this would be located near to the knot region of each chain and, thus may

10 have little effect on stabilizing the putative receptor binding regions at the tips of Finger 1 and Finger 2. Additionally, the proximity of such a linkage to the disulfide bonds in the knot region might interfere with the proper formation of those structures.

With regard to non-covalent interactions, it is contemplated that the structural stability of the hOP-1 dimer can be enhanced by increasing inter-chain hydrogen bonding.

15 The electrostatic potential (due to other charges in a protein) in the region of a charged residue affects the pK of that residue. Because the pK's of both histidine and the N-terminal primary amino group are near neutrality, it may be possible to modify their pKs through the placement of charges on the surface of the molecule. It is contemplated that the buried His at position His 84 in hOP-1 helps stabilize the structure of the dimer by participating in hydrogen

20 bonds with backbone carboxyl groups of residues Ala 64 and Tyr 65 of the other chain. Accordingly, it is contemplated that the introduction of surface charges may enhance this effect and thereby further stabilize the structure of the molecule. For example, mutating Tyr 65 or Val 132 to Asp may further polarize the carbonyl bonds of the amino acid residues at positions 64 and 65, as well as raise the pK of His 84. The pK of His 84 may further be affected by replacing

25 residues Tyr 44, Ala 63, or Asn 110 by an Asp. It is contemplated that the preferred modification for this purpose is Tyr 65-> Asp 65.

Using the same basic principles, the skilled artisan likewise can identify pairs of amino acids whose replacement can facilitate the introduction of an inter-chain salt bridge, internal hydrogen bond, or hydrophobic interaction. Such determinations are within the scope of an

30 artisan having an ordinary level of skill in the field of molecular modeling.

Once a pair of target amino acids has been identified, the site-directed replacement of the target amino acids with the desirable replacements can be facilitated by the use of conventional site-directed mutagenesis procedures, for example, by cassette mutagenesis or oligonucleotide-directed mutagenesis. Such techniques are thoroughly documented in the art and so are not
5 discussed herein. The effect of the site-specific replacements on the stability of resulting modified hOP-1 dimers or muteins can be measured, after production and purification, using standard methodologies well known in the art, for example, circular dichroism, analytical centrifugation, differential scanning calorimetry, fluorescence or other spectroscopic techniques.

(ii) Enhancing Water Solubility of hOP-1 Dimer

10 OP-1 has limited solubility in aqueous solvents. It is contemplated, however, that by using the hOP-1 atomic co-ordinates that the artisan can replace amino acids at the solvent accessible surface of the dimer thereby to increase the dielectric properties of dimeric hOP-1. For example, solvent accessible hydrophobic amino acid residues, such as, glycine, alanine, valine, leucine and isoleucine may be replaced by more polar residues, such as, lysine, arginine,
15 histidine, aspartate, asparagine, glutamate and glutamine.

The solvent accessible amino acids can be identified using a computer program, such as ACCESS (version 2.1) using a 1.4 Å probe (Lee *et al.* (1972) *supra*). In Fig. 7 amino acid residues having at least 20% of their side chain areas exposed to solvent are boxed. When modifying surface residues it is important not to produce new epitopes that can be recognized as
20 non-host especially, if the hOP-1 analogs are to be used as injectable molecules. It is believed that, amino acid side chains seen by a 10 Å spherical probe likely are part of surface epitopes. One skilled in the art can use ACCESS with a 10 Å spherical probe to identify potential epitopes, however this process can be carried out manually using a graphics package, such as, INSIGHT II. In Figure 8, residue side chains so identified as potential epitopes are highlighted. Residue
25 positions that are candidates for modification so as to improve the solubility of the dimer are highlighted. Preferred candidate amino acids for replacement include, for example, Ala 63, Ala 72, Ala 81, Ala 111 and Ala 135, Ile 86, Ile 112, Tyr 52, Tyr 65, and Tyr 128.

Once solvent accessible hydrophobic or non polar amino acids have been identified (see Fig. 9), these amino acids theoretically may be replaced, via a computer, with more polar amino

acids. The effect of the amino acid replacements on the solution electrostatic potentials surrounding the modified hOP-1 dimer as well as the free energy of the dimer can be calculated using the program DELPHI (Gilson *et al* (1987) *supra*; Nicholls *et al.* (1991) *supra*). Preferred amino acid substitutions lower the free energy of the hOP-1 dimer without introducing potential antigenic sites. As mentioned above, such antigenic sites may be detected by implementing a computer program like ACCESS (version 2.1) using a 10 Å probe. In addition, it is contemplated that preferred surface residues suitable for replacement do not constitute part of the receptor binding domain.

The resulting candidate morphogen analogs can be produced using conventional site-directed mutagenesis methodologies and the effect of the site-directed modification on the solubility of the hOP-1 dimer can be measured, for example, by comparing the partition coefficient or "salting out properties" of the modified hOP-1 dimer versus the native hOP-1 dimer. See for example, Scopes (1987) in *Protein Purification: Principles and Practice, 2nd Edition* (Springer-Verlag); and England *et al.* (1990) *Methods in Enzymology* 182: 285-300.

(iii) Engineering Glycosylation Sites

In addition to replacing single, solvent accessible amino acid residues with more polar or hydrophobic amino acid residue, one or more solvent accessible amino acid residues may be replaced so as to create a new eukaryotic glycosylation site or alternatively to eliminate or alter an existing glycosylation site. Glycosylation sites are well known and are thoroughly described in the art. Addition of a new glycosylation site or alteration of an existing site may result in the addition of one or more glycosyl groups, e.g., N-acetyl-sialic acid, which may enhance the solubility of the morphogen analog. As described herein, such sites can be introduced by site-directed mutagenesis methodologies which are well known in the art. Preferably, such sites do not create new antigenic determinants (although these may be tolerable for short duration therapeutic uses). Reference to Table 8 identifies surface accessible amino acid residues, based on the 2.8Å structure, which likely are not part of an antigenic epitope and which may be used as candidates for introducing an additional glycosylation site.

B. Engineering Small Molecules Based Upon The hOP-1 Structure

The availability of atomic co-ordinates for hOP-1 enables the skilled artisan to design small molecules, for example, peptides or non-peptidyl based organic molecules having certain chemical features, which mimic the biological activity of hOP-1. Chemical features of interest may include, for example, the three-dimensional structure of a particular protein domain, solvent accessible surface of a particular protein domain, spatial distribution of charged and/or hydrophobic chemical moieties, electrostatic charge distribution, or a combination thereof. Such chemical features may readily be determined from the three-dimensional representation of hOP-1.

(i) Peptides

After having determined which amino acid residues contribute to the receptor binding domain (*supra*), it is possible for the skilled artisan to design synthetic peptides having amino acid sequences that define a pre-selected receptor binding motif. A computer program useful in designing potentially bioactive peptido-mimetics is described in U.S. Patent No. 5,331,573, the disclosure of which is incorporated by reference herein.

In addition to choosing a desirable amino acid sequence, the skilled artisan using standard molecular modeling software packages, *infra*, can design specific peptides having, for example, additional cysteine amino acids located at pre-selected positions to facilitate cyclization of the peptide of interest. Oxidation of the additional cysteine residues results in cyclization of the peptide thereby constraining the peptide in a conformation which mimics the conformation of the corresponding amino acid sequence in native hOP-1. It is contemplated, that any standard covalent linkage, for example, disulfide bonds, typically used to cyclize synthetic peptides maybe useful in the practice of the instant invention. Alternative cyclization chemistries are discussed in International Application PCT/WO 95/01800, the disclosure of which is incorporated herein by reference.

In addition, it is contemplated that a single peptide containing amino acid sequences derived from separate hOP-1 subunit domains, for example, a single peptide having an amino acid sequence defining the tip of the finger 1 region linked by means of a polypeptide linker to an amino acid sequence defining the tip of the finger 2 region. The amino sequence defining

each of the finger regions may further comprise a means, for example, disulfide bonds for cyclizing each finger region motif. The resulting peptide therefore comprises a single polypeptide chain having a first amino acid sequence defining a three-dimensional domain mimicking the tip of the finger 1 region and a second said sequence defining a three-dimensional domain mimicking the tip of the finger 2 region.

Such peptides may be synthesized and screened for OP-1 like activity using any of the standard protocols described below.

(ii) Organic molecules

As discussed above, upon determination of the receptor binding domain of hOP-1, it is contemplated that the skilled artisan, can design non-peptidyl based small molecules, for example, small organic molecules, whose structural and chemical features mimic the same features displayed on at least part of the surface of the receptor binding domain of hOP-1.

Because a major contribution to the receptor binding surface is the spatial arrangement of chemically interactive moieties present within the sidechains of amino acids which together define the receptor binding surface, a preferred embodiment of the present invention relates to designing and producing a synthetic organic molecule having a framework that carries chemically interactive moieties in a spatial relationship that mimics the spatial relationship of the chemical moieties disposed on the amino acid sidechains which constitute the receptor binding site of hOP-1. Preferred chemical moieties, include but are not limited to, the chemical moieties defined by the amino acid side chains of amino acids believed to constitute the receptor binding domain of hOP-1 (See Fig. 9). It is understood, therefore, that the receptor binding surface of the morphogen analog need not comprise amino acid residues but the chemical moieties disposed thereon.

For example, upon identification of relevant chemical groups, the skilled artisan using a conventional computer program can design a small molecule having the receptor interactive chemical moieties disposed upon a suitable carrier framework. Useful computer programs are described in, for example, Dixon (1992) *Tibtech* 10: 357-363; Tschinke *et al.* (1993) *J. Med. Chem* 36: 3863-3870; and Eisen *et al.* (1994) *Proteins: Structure, Function, and Genetics* 19: 199-221, the disclosures of which are incorporated herein by reference.

- 36 -

One particular computer program entitled "CAVEAT" searches a database, for example, the Cambridge Structural Database, for structures which have desired spatial orientations of chemical moieties (Bartlett *et al.* (1989) in "*Molecular Recognition: Chemical and Biological Problems*" (Roberts, S.M., ed) pp 182-196). The CAVEAT program has been used to design
5 analogs of tendamistat, a 74 residue inhibitor of α -amylase, based on the orientation of selected amino acid side chains in the three-dimensional structure of tendamistat (Bartlett *et al.* (1989) *supra*).

Alternatively, upon identification of a series of analogs which mimic the biological activity of OP-1, as determined by *in vivo* or *in vitro* assays, the skilled artisan may use a variety
10 of computer programs which assist the skilled artisan to develop quantitative structure activity relationships (QSAR) and further to assist in the *de novo* design of additional morphogen analogs. Other useful computer programs are described in, for example, Connolly-Martin (1991) *Methods in Enzymology* 203:587-613; Dixon (1992) *supra*; and Waszkowycz *et al.* (1994) *J. Med. Chem.* 37: 3994-4002.

Thus, for example, one can begin with a portion of the three dimensional structure of OP-1 (or a related morphogen) corresponding to a region of known or suspected biological importance. One such region is the solvent accessible loop or "tip" of the finger 2 region between the $\beta 6$ and $\beta 7$ sheets (i.e., from approximately residues 118-122). Synthetic, cyclic peptides (i.e., F2-2 and F2-3) were produced including this region (and several flanking residues)
20 and were shown to possess OP-1-like biological activity (see Examples below). Based upon the three-dimensional structure of this region, disclosed herein, one is now enabled to produce more effective OP-1-like (or, generally, morphogen-like) analogs. For example, as shown in great detail in Figures 7-9 and 15, the charged γ -carboxy groups of Asp 118 and Asp 119, and the relatively hydrophilic hydroxyl groups of Ser 120 and Ser 121, are solvent accessible and
25 believed to be involved in OP-1 receptor binding. The relative positions of these groups in three dimensions in OP-1 are given in Figures 15 and 16. These functional groups define a contiguous portion of the three dimensional structure of the OP-1 surface. The peptide backbone of these residues, however, is not solvent accessible and, therefore, is not believed to form a portion of the three-dimensional surface of the OP-1 molecule. Thus, one of ordinary skill in the art, when
30 choosing or designing an OP-1 or morphogen analog, can choose or design a molecule having

the same or substantially equivalent (e.g., thiol v. hydroxyl) functional groups in substantially the same (e.g., $\pm 1-3$ Å) three-dimensional conformation. The same is true for other regions of interest in the OP-1 monomers or dimers (e.g., the receptor binding domain, the finger 1, finger 2, or heel regions, or solvent accessible portions thereof). By using the three-dimensional structures disclosed herein, including the disclosure of the positions of solvent accessible and probable receptor contact residues, one of ordinary skill in the art can choose a portion of the three-dimensional structure of the OP-1 (or a related morphogen) molecule and, using this "portion" as a template select or design an analog which functionally mimics the template structure.

10 The molecular framework or backbone of the morphogen analog can be freely chosen by one of ordinary skill in the art so that it (1) joins the functional groups which mimic the portion of the morphogen's contiguous three-dimensional surface, including charge distribution and hydrophobicity/hydrophilicity characteristics, and (2) maintains or, at least, allows the functional groups to maintain the appropriate three-dimensional surface interaction and spatial relationships, including any hydrogen bonding and electrostatic interactions. As described above, peptides are obvious choices for the production of such morphogen analogs because they can provide all of the necessary functional groups and can assume appropriate three-dimensional structures. Several examples of peptide analogs of the finger regions are described herein, below. The peptides are cyclized to maintain hydrogen bonds and create a structure which mimics that of the template. These peptides are synthesized from a linear primary sequence of amino acids in finger 2. An alternative peptide can be created, for example, which combines portions of finger 1 and finger 2, constructed to mimic the structure of the tips of fingers 1 and 2 together as they occur in the folded OP1 monomer. Biologically active peptides such as F2, F3 or others, then can be used as is or, more preferably, become lead compounds for iterative modification to create a compound that is more stable or more active *in vivo*. For example, the peptide backbone can be reduced or replaced to reduce hydrolysis *in vivo*. Alternatively, structural modifications can be introduced to the backbone or by amino acid substitutions which more accurately mimic the protein's structure when bound to the receptor. These second generation structures then can be tested for enhanced binding. In addition, iterative amino acid

replacements with alanines, ("alanine scan") can be used to determine the minimum residue contacts required for binding.

Once these minimum functional groups are known, a fully synthetic molecule can be created which mimics the charge or electrostatic distribution of the minimum required functional groups, and provides the appropriate bulk and structure to functionally mimic a second generation molecule having the desired binding affinity.

VI. Production of Morphogen Analogs.

As mentioned above, the morphogen analogs of the invention may comprise modified hOP-1 dimeric proteins or small molecules, for example, peptides or small organic molecules. It is contemplated that any appropriate methods can be used for producing a pre-selected morphogen analog. For example, such methods may include, but are not limited to, methods of biological production from suitable host cells or synthetic production using synthetic organic chemistries.

For example, modified hOP-1 dimeric proteins or hOP-based peptides may be produced using conventional recombinant DNA technologies, well known and thoroughly documented in the art. Under these circumstances, the proteins or peptides may be produced by the preparation of nucleic acid sequences encoding the respective protein or peptide sequences, after which, the resulting nucleic acid can be expressed in an appropriate host cell. By way of example, the proteins and peptides may be manufactured by the assembly of synthetic nucleotide sequences and/or joining DNA restriction fragments to produce a synthetic DNA molecule. The DNA molecules then are ligated into an expression vehicle, for example an expression plasmid, and transfected into an appropriate host cell, for example *E. coli*. The protein encoded by the DNA molecule then is expressed, purified, folded if necessary, tested *in vitro* for binding activity with an OP-1 receptor, and subsequently tested to assess whether the morphogen analog induces or stimulates hOP-1-like biological activity.

The processes for manipulating, amplifying, and recombining DNA which encode amino acid sequences of interest generally are well known in the art, and therefore, are not described in detail herein. Methods of identifying and isolating genes encoding hOP-1 and its cognate receptors also are well understood, and are described in the patent and other literature.

Briefly, the construction of DNAs encoding the biosynthetic constructs disclosed herein is performed using known techniques involving the use of various restriction enzymes which make sequence specific cuts in DNA to produce blunt ends or cohesive ends, DNA ligases, techniques enabling enzymatic addition of sticky ends to blunt-ended DNA, construction of synthetic DNAs by assembly of short or medium length oligonucleotides, cDNA synthesis techniques, polymerase chain reaction (PCR) techniques for amplifying appropriate nucleic acid sequences from libraries, and synthetic probes for isolating OP-1 genes or genes encoding other members of the TGF- β superfamily as well as their cognate receptors. Various promoter sequences from bacteria, mammals, or insects to name a few, and other regulatory DNA sequences used in achieving expression, and various types of host cells are also known and available. Conventional transfection techniques, and equally conventional techniques for cloning and subcloning DNA are useful in the practice of this invention and known to those skilled in the art. Various types of vectors may be used such as plasmids and viruses including animal viruses and bacteriophages. The vectors may exploit various marker genes which impart to a successfully transfected cell a detectable phenotypic property that can be used to identify which of a family of clones has successfully incorporated the recombinant DNA of the vector.

One method for obtaining DNA encoding the biosynthetic constructs disclosed herein is by assembly of synthetic oligonucleotides produced in a conventional, automated, oligonucleotide synthesizer followed by ligation with appropriate ligases. For example, overlapping, complementary DNA fragments may be synthesized using phosphoramidite chemistry, with end segments left unphosphorylated to prevent polymerization during ligation. One end of the synthetic DNA is left with a "sticky end" corresponding to the site of action of a particular restriction endonuclease, and the other end is left with an end corresponding to the site of action of another restriction endonuclease. The complementary DNA fragments are ligated together to produce a synthetic DNA construct.

After the appropriate DNA molecule has been synthesized, it may be integrated into an expression vector and transfected into an appropriate host cell for protein expression. Useful prokaryotic host cells include, but are not limited to, *E. coli*, and *B. subtilis*. Useful eukaryotic host cells include, but are not limited to, yeast cells, insect cells, myeloma cells, fibroblast 3T3 cells, monkey kidney or COS cells, chinese hamster ovary (CHO) cells, mink-lung epithelial

cells, human foreskin fibroblast cells, human glioblastoma cells, and teratocarcinoma cells. Alternatively, the genes may be expressed in a cell-free system such as the rabbit reticulocyte lysate system.

The vector additionally may include various sequences to promote correct expression of the recombinant protein, including transcriptional promoter and termination sequences, enhancer sequences, preferred ribosome binding site sequences, preferred mRNA leader sequences, preferred protein processing sequences, preferred signal sequences for protein secretion, and the like. The DNA sequence encoding the gene of interest also may be manipulated to remove potentially inhibiting sequences or to minimize unwanted secondary structure formation. The morphogenic protein analogs proteins also may be expressed as fusion proteins. After being translated, the protein may be purified from the cells themselves or recovered from the culture medium and then cleaved at a specific protease site if so desired.

For example, if the gene is to be expressed in *E. coli*, it is cloned into an appropriate expression vector. This can be accomplished by positioning the engineered gene downstream of a promoter sequence such as Trp or Tac, and/or a gene coding for a leader peptide such as fragment B of protein A (FB). During expression, the resulting fusion proteins accumulate in refractile bodies in the cytoplasm of the cells, and may be harvested after disruption of the cells by French press or sonication. The isolated refractile bodies then are solubilized, and the expressed proteins folded and the leader sequence cleaved, if necessary, by methods already established with many other recombinant proteins.

Expression of the engineered genes in eukaryotic cells requires cells and cell lines that are easy to transfect, are capable of stably maintaining foreign DNA with an unrearranged sequence, and which have the necessary cellular components for efficient transcription, translation, post-translation modification, and secretion of the protein. In addition, a suitable vector carrying the gene of interest also is necessary. DNA vector design for transfection into mammalian cells should include appropriate sequences to promote expression of the gene of interest as described herein, including appropriate transcription initiation, termination, and enhancer sequences, as well as sequences that enhance translation efficiency, such as the Kozak consensus sequence. Preferred DNA vectors also include a marker gene and means for amplifying the copy number of the gene of interest. A detailed review of the state of the art of the production of foreign proteins

in mammalian cells, including useful cells, protein expression-promoting sequences, marker genes, and gene amplification methods, is disclosed in Bendig (1988) *Genetic Engineering* 7:91-127.

5 The best characterized transcription promoters useful for expressing a foreign gene in a particular mammalian cell are the SV40 early promoter, the adenovirus promoter (AdMLP), the mouse metallothionein-I promoter (mMT-I), the Rous sarcoma virus (RSV) long terminal repeat (LTR), the mouse mammary tumor virus long terminal repeat (MMTV-LTR), and the human cytomegalovirus major intermediate-early promoter (hCMV). The DNA sequences for all of these promoters are known in the art and are available commercially.

10 The use of a selectable DHFR gene in a dhfr⁻ cell line is a well characterized method useful in the amplification of genes in mammalian cell systems. Briefly, the DHFR gene is provided on the vector carrying the gene of interest, and addition of increasing concentrations of the cytotoxic drug methotrexate, which is metabolized by DHFR, leads to amplification of the DHFR gene copy number, as well as that of the associated gene of interest. DHFR as a
15 selectable, amplifiable marker gene in transfected chinese hamster ovary cell lines (CHO cells) is particularly well characterized in the art. Other useful amplifiable marker genes include the adenosine deaminase (ADA) and glutamine synthetase (GS) genes.

The choice of cells/cell lines is also important and depends on the needs of the experimenter. COS cells provide high levels of transient gene expression, providing a useful
20 means for rapidly screening the biosynthetic constructs of the invention. COS cells typically are transfected with a simian virus 40 (SV40) vector carrying the gene of interest. The transfected COS cells eventually die, thus preventing the long term production of the desired protein product but provide a useful technique for testing preliminary analogs for binding activity.

The various cells, cell lines and DNA sequences that can be used for mammalian cell
25 expression of the single-chain constructs of the invention are well characterized in the art and are readily available. Other promoters, selectable markers, gene amplification methods and cells also may be used to express the proteins of this invention. Particular details of the transfection, expression, and purification of recombinant proteins are well documented in the art and are understood by those having ordinary skill in the art. Further details on the various technical

aspects of each of the steps used in recombinant production of foreign genes in mammalian cell expression systems can be found in a number of texts and laboratory manuals in the art, such as, for example, Ausubel *et al.*, ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, (1989).

5 Alternatively, morphogen analogs which are small peptides, usually up to 50 amino acids in length, may be synthesized using standard solid-phase peptide synthesis procedures, for example, procedures similar to those described in Merrifield (1963) *J. Am. Chem. Soc.*, 85:2149. For example, during synthesis, N- α -protected amino acids having protected side chains are added stepwise to a growing polypeptide chain linked by its C-terminal end to an insoluble
10 polymeric support, e.g., polystyrene beads. The peptides are synthesized by linking an amino group of an N- α -deprotected amino acid to an α -carboxy group of an N- α -protected amino acid that has been activated by reacting it with a reagent such as dicyclohexylcarbodiimide. The attachment of a free amino group to the activated carboxyl leads to peptide bond formation. The most commonly used N- α -protecting groups include Boc which is acid labile and Fmoc which is
15 base labile.

Briefly, the C-terminal N- α -protected amino acid is first attached to the polystyrene beads. Then, the N- α -protecting group is removed. The deprotected α -amino group is coupled to the activated α -carboxylate group of the next N- α -protected amino acid. The process is repeated until the desired peptide is synthesized. The resulting peptides are cleaved from the insoluble
20 polymer support and the amino acid side chains deprotected. Longer peptides, for example greater than about 50 amino acids in length, typically are derived by condensation of protected peptide fragments. Details of appropriate chemistries, resins, protecting groups, protected amino acids and reagents are well known in the art and so are not discussed in detail herein. See for example, Atherton *et al.* (1963) *Solid Phase Peptide Synthesis: A Practical Approach* (IRL
25 Press,), and Bodanszky (1993) *Peptide Chemistry, A Practical Textbook, 2nd Ed*, Springer-Verlag, and Fields *et al.* (1990) *Int. J. Peptide Protein Res.* 35:161-214, the disclosures of which are incorporated herein by reference.

Purification of the resulting peptide is accomplished using conventional procedures, such as preparative HPLC, e.g., gel permeation, partition and/or ion exchange chromatography. The

choice of appropriate matrices and buffers are well known in the art and so are not described in detail herein.

With regard to the production of non-peptide small organic molecules which induce OP-1 like biological activities, these molecules can be synthesized using standard organic chemistries well known and thoroughly documented in the patent and other literatures.

VII. Screening For Binding and Biological Activity.

As a first step in determining whether a morphogen analog induces an OP-1 like biological activity, the skilled artisan can use a standard ligand-receptor assay to determine whether the morphogen analog binds preferentially to OP-1 receptor. For standard receptor-ligand assays, the artisan is referred to, for example, Legerski *et al.* (1992) *Biochem. Biophys. Res. Comm.* 183: 672-679; Frakar *et al.* (1978) *Biochem. Biophys. Res. Comm.* 80:849-857; Chio *et al.* (1990) *Nature* 343: 266-269; Dahlman *et al.* (1988) *Biochem* 27: 1813-1817; Strader *et al.* (1989) *J. Biol. Chem.* 264: 13572-13578; and D'Dowd *et al.* (1988) *J. Biol. Chem.* 263: 15985-15992.

In a typical ligand/receptor binding assay useful in the practice of this invention, purified OP-1 having a known, quantifiable affinity for a pre-selected OP-1 receptor (see, for example, Ten Dijke *et al.* (1994) *Science* 264:101-103, the disclosure of which is incorporated herein by reference) is labeled with a detectable moiety, for example, a radiolabel, a chromogenic label, or a fluorogenic label. Aliquots of purified receptor, receptor binding domain fragments, or cells expressing the receptor of interest on their surface are incubated with labeled OP-1 in the presence of various concentrations of the unlabeled morphogen analog. The relative binding affinity of the morphogen analog may be measured by quantitating the ability of the candidate (unlabeled morphogen analog) to inhibit the binding of labeled OP-1 with the receptor. In performing the assay, fixed concentrations of the receptor and the OP-1 are incubated in the presence and absence of unlabeled morphogen analog. Sensitivity may be increased by pre-incubating the receptor with the candidate morphogen analog before adding labeled OP-1. After the labeled competitor has been added, sufficient time is allowed for adequate competitor binding, and then free and bound labeled OP-1 are separated from one another, and one or the other measured.

Labels useful in the practice of the screening procedures include radioactive labels (e.g., ^{125}I , ^{131}I , ^{111}In or ^{77}Br), chromogenic labels, spectroscopic labels (such as those disclosed in Haughland (1994) "*Handbook of Fluorescent and Research Chemicals 5 ed.*" by Molecular Probes, Inc., Eugene, OR), or conjugated enzymes having high turnover rates, for example, 5 horseradish peroxidase, alkaline phosphatase, or β -galactosidase, used in combination with chemiluminescent or fluorogenic substrates.

The biological activity, namely the agonist or antagonist properties of the resulting morphogen analogs subsequently may be characterized using any conventional *in vivo* and *in vitro* assays that have been developed to measure the biological activity of OP-1. A variety of 10 specific assays believed to be useful in the practice of the invention are set forth in detail in Example 1, hereinbelow.

Furthermore, it is appreciated that many of the standard OP-1 assays may be automated thereby facilitating the screening of a large number of morphogen analogs at the same time. Such automation procedures are within the level of skill in the art of drug screening and, 15 therefore, are not discussed herein.

Following the identification of useful morphogen analogs, the morphogenic analogs may be produced in commercially useful quantities (e.g., without limitation, gram and kilogram quantities), for example, by producing cell lines that express the morphogen analogs of interest or by producing synthetic peptides defining the appropriate amino acid sequence. It is 20 appreciated, however, that conventional methodologies for producing the appropriate cell lines and for producing synthetic peptides are well known and thoroughly documented in the art, and so are not discussed in detail herein.

VIII. Formulation and Bioactivity.

Morphogen analogs, including OP-1 analogs, can be formulated for administration to a 25 mammal, preferably a human in need thereof as part of a pharmaceutical composition. The composition can be administered by any suitable means, e.g., parenterally, orally or locally. Where the morphogen analog is to be administered locally, as by injection, to a desired tissue site, or systemically, such as by intravenous, subcutaneous, intramuscular, intraorbital, ophthalmic, intraventricular, intracranial, intracapsular, intraspinal, intracisternal, intraperitoneal,

buccal, rectal, vaginal, intranasal or aerosol administration, the composition preferably comprises an aqueous solution. The solution preferably is physiologically acceptable, such that administration thereof to a mammal does not adversely affect the mammal's normal electrolyte and fluid volume balance. The aqueous solution thus can comprise, e.g., normal physiologic saline (0.9% NaCl, 0.15M), pH 7-7.4.

Useful solutions for oral or parenteral systemic administration can be prepared by any of the methods well known in the pharmaceutical arts, described, for example, in "*Remington's Pharmaceutical Sciences*" (Gennaro, A., ed., Mack Pub., 1990, the disclosure of which is incorporated herein by reference). Formulations can include, for example, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, hydrogenated naphthalenes, and the like. Formulations for direct administration, in particular, can include glycerol and other compositions of high viscosity. Biocompatible, preferably bioresorbable polymers, including, for example, hyaluronic acid, collagen, tricalcium phosphate, polybutyrate, polylactide, polyglycolide and lactide/glycolide copolymers, may be useful excipients to control the release of the morphogen analog *in vivo*.

Other potentially useful parenteral delivery systems for the present analogs can include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation administration can contain as excipients, for example, lactose, or can be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate or deoxycholate, or oily solutions for administration in the form of nasal drops or as a gel to be applied intranasally.

Alternatively, the morphogen analogs, including OP-1 analogs, identified as described herein may be administered orally. For example, liquid formulations of morphogen analogs can be prepared according to standard practices such as those described in "*Remington's Pharmaceutical Sciences*" (*supra*). Such liquid formulations can then be added to a beverage or another food supplement for administration. Oral administration can also be achieved using aerosols of these liquid formulations. Alternatively, solid formulations prepared using art-recognized emulsifiers can be fabricated into tablets, capsules or lozenges suitable for oral administration.

Optionally, the analogs can be formulated in compositions comprising means for enhancing uptake of the analog by a desired tissue. For example, tetracycline and diphosphonates (bisphosphonates) are known to bind to bone mineral, particularly at zones of bone remodeling, when they are provided systemically in a mammal. Accordingly, such components can be used to enhance delivery of the present analogs to bone tissue. Alternatively, an antibody or portion thereof that binds specifically to an accessible substance specifically associated with the desired target tissue, such as a cell surface antigen, also can be used. If desired, such specific targeting molecules can be covalently bound to the present analog, e.g., by chemical crosslinking or by using standard genetic engineering techniques to create, for example, an acid labile bond such as an Asp-Pro linkage. Useful targeting molecules can be designed, for example, according to the teachings of U.S. 5,091,513.

It is contemplated also that some of the morphogen analogs may exhibit the highest levels of activity *in vivo* when combined with carrier matrices i.e., insoluble polymer matrices. See for example, U.S. Patent No. 5,266,683 the disclosure of which is incorporated by reference herein. Currently preferred carrier matrices are xenogenic, allogenic or autogenic in nature. It is contemplated, however, that synthetic materials comprising polylactic acid, polyglycolic acid, polybutyric acid, derivatives and copolymers thereof may also be used to generate suitable carrier matrices. Preferred synthetic and naturally derived matrix materials, their preparation, methods for formulating them with the morphogen analogs of the invention, and methods of administration are well known in the art and so are not discussed in detailed herein. See for example, U.S. Patent No. 5,266,683.

Still further, the present analogs can be administered to the mammal in need thereof either alone or in combination with another substance known to have a beneficial effect on tissue morphogenesis. Examples of such substances (herein, cofactors) include substances that promote tissue repair and regeneration and/or inhibit inflammation. Examples of useful cofactors for stimulating bone tissue growth in osteoporotic individuals, for example, include but are not limited to, vitamin D₃, calcitonin, prostaglandins, parathyroid hormone, dexamethasone, estrogen and IGF-I or IGF-II. Useful cofactors for nerve tissue repair and regeneration can include nerve growth factors. Other useful cofactors include symptom-alleviating cofactors, including antiseptics, antibiotics, antiviral and antifungal agents, analgesics and anesthetics.

Analog preferably are formulated into pharmaceutical compositions by admixture with pharmaceutically acceptable, nontoxic excipients and carriers. As noted above, such compositions can be prepared for systemic, e.g., parenteral, administration, particularly in the form of liquid solutions or suspensions; for oral administration, particularly in the form of tablets or capsules; or intranasally, particularly in the form of powders, nasal drops or aerosols. Where adhesion to a tissue surface is desired, the composition can comprise a fibrinogen-thrombin dispersant or other bioadhesive such as is disclosed, for example, in PCT US91/09275, the disclosure of which is incorporated herein by reference. The composition then can be painted, sprayed or otherwise applied to the desired tissue surface.

10 The compositions can be formulated for parenteral or oral administration to humans or other mammals in therapeutically effective amounts, e.g., amounts which provide appropriate concentrations of the morphogen analog to target tissue for a time sufficient to induce the desired effect. Preferably, the present compositions alleviate or mitigate the mammal's need for a morphogen-associated biological response, such as maintenance of tissue-specific function or
15 restoration of tissue-specific phenotype to senescent tissues (e.g., osteopenic bone tissue).

As will be appreciated by those skilled in the art, the concentration of the compounds described in a therapeutic composition will vary depending upon a number of factors, including the dosage of the drug to be administered, the chemical characteristics (e.g., hydrophobicity) of the compounds employed, and the route of administration. The preferred dosage of drug to be
20 administered also is likely to depend on such variables as the type and extent of a disease, tissue loss or defect, the overall health status of the particular patient, the relative biological efficacy of the compound selected, the formulation of the compound, the presence and types of excipients in the formulation, and the route of administration. In general terms, the therapeutic molecules of this invention may be provided to an individual where typical doses range from about 10 ng/kg
25 to about 1 g/kg of body weight per day; with a preferred dose range being from about 0.1 mg/kg to 100 mg/kg of body weight.

IX. Examples

Practice of the invention will be more fully understood from the following examples, which are presented herein for illustrative purposes only, and should not be construed as limiting the invention in any way.

5 Example 1. Introduction of Inter-chain Disulfide Bonds to Stabilize the hOP-1 Dimer.

As discussed in section V.A.(i) it is contemplated that introduction of one or more additional inter-chain disulfide may stabilize further the hOP-1 dimer. The introduction of additional inter-chain disulfide bonds is described here.

10 A Sma I to Bam HI fragment of the human OP-1 cDNA as described in Ozkaynak *et al.* (1990) *supra* is cloned into Bluescript KS+ (available from Stratagene Cloning Systems, La Jolla, CA), previously cleaved with Eco RV and Bam HI. Upon transformation into *E. coli*, the resulting colonies are screened by a blue-white selection process wherein the desired colonies containing the OP-1 cDNA insert are blue. The correct clone may be identified by restriction screening to give the following expected restriction fragments.

<u>Restriction Enzyme</u>	<u>Fragment size (bp)</u>
EcoR I	84, 789, 3425
Xho I	161, 1223, 2914
Sac II	97, 650, 3551

15

In order to introduce two additional inter-chain disulfide bridges, a double cysteine mutant containing Asn 83 to Cys and Asn 130 to Cys replacements is produced. The cysteine mutant can be prepared by site-directed mutagenesis using synthetic oligonucleotides and either PCR or the site-directed mutagenesis methods, see for example, Kunkel *et al.* (1985) *Proc. Natl. Acad. Sci. USA* 822: 488; Kunkel *et al.* (1985) *Meth. Enzymol.* 154: 367 and U.S. Patent No. 4,873,192. Neither mutation causes a frameshift and, therefore, *E. coli* transformed with mutagenesis products that give white colonies indicate an error in the sequence. The presence of the appropriate mutation is verified by conventional dideoxy sequencing.

20

Then, linkers are introduced into the N- and C-termini of the mutant gene by oligonucleotide-directed mutagenesis using appropriate oligonucleotides. A preferred N terminal linker introduces a unique Not I site and a preferred C terminal linker introduces a non-suppressible stop codon TAA at the end of the mutein gene followed by a unique Bgl II site (AGATCT). Each of the resulting mutant genes are excised from the cloning vector by the restriction enzymes Nde I and Bgl II, isolated, and ligated independently into pET vector (New England Biolabs, Beverly, MA) previously cleaved with Nde I and Bam HI. The ligation products then are transformed into *E. coli* and transformants containing, and expressing each individual mutant protein are identified.

Expression of the double cysteine containing mutant analog is induced after the expression of T7 RNA polymerase (initiated through infected with λ CE6 phage). During expression, the mutant analog is produced as inclusion granules which are harvested from the cell paste. Then, the mutant protein is dissolved in 6M guanidine-HCl, 0.2M Tris-HCl, pH 8.2 and 0.1 M 2-mercaptoethanol, and the mixture dialyzed exhaustively against 6M urea, 2.5 mM Tris-HCl, pH 7.5 and 1 mM EDTA. 2-mercaptoethanol is added to a final concentration of 0.1M and the solution incubated at room temperature. The mixture is dialyzed exhaustively against buffer containing 2.5 mM Tris-HCl, pH 7.5 and 1mM EDTA. Folded mutant protein is purified by affinity chromatography on a column packed with surface immobilized OP-1 receptor. Unbound material is removed by washing as described above and the specific OP-1 receptor binding material eluted.

Following purification the stabilizing effect of the additional bond is determined by fluorescence polarization. For example, the rotational rates of morphogen analog (mutein) and natural hOP-1 are determined as a function of temperature using a fluorescence spectrophotometer modified for fluorescence anisotropy (Photon Technology International). It is anticipated that the mutein dimer will exhibit a lower rotational rate upto a higher temperature than natural hOP-1 dimer, thereby indicating that the mutein dimer remains as a dimer and is more stable upto a higher temperature than is the wild type protein.

The biological activity of the resulting mutant protein or mutein can be tested using any of the bioassays developed to date for determining the biological activity of native hOP-1. A variety of such exemplary assays are described below. The assays which follow are recited for

- 50 -

ease of testing. Specific *in vivo* assays for testing the efficacy of a morphogenic protein or analog in an application to repair or regenerate damaged bone, liver, kidney, or nerve tissue, periodontal tissue, including cementum and/or periodontal ligament, gastrointestinal and renal tissues, and immune-cell mediated damages tissues are disclosed in publicly available documents, which include, for example, EP 0575,555; WO93/04692; WO93/05751; WO/06399; WO94/03200; WO94/06449; and WO94/06420. The skilled artisan can test an analog in any of these assays without undue experimentation.

A. Mitogenic Effect on Rat and Human Osteoblasts

The following example is a typical assay useful in determining whether an OP-1 morphogen analog induces proliferation of osteoblasts *in vitro*. It is contemplated that in this, and all other examples using osteoblast cultures, preferably uses rat osteoblast-enriched primary cultures. Although these cultures are heterogeneous in that the individual cells are at different stages of differentiation, the culture is believed to more accurately reflect the metabolism and function of osteoblasts *in vivo* than osteoblast cultures obtained from established cell lines. Unless otherwise indicated, all chemicals referenced are standard, commercially available reagents, readily available from a number of sources, including Sigma Chemical, Co., St. Louis; Calbiochem, Corp., San Diego and Aldrich Chemical Co., Milwaukee.

Briefly, rat osteoblast-enriched primary cultures are prepared by sequential collagenase digestion of newborn rat calvaria (e.g., from 1-2 day-old animals, Long-Evans strain, Charles River Laboratories, Wilmington, MA), following standard procedures, such as are described, for example, in Wong *et al.* (1975) *Proc. Natl. Acad. Sci. USA* 72: 3167-3171. Rat osteoblast single cell suspensions then are plated onto a multi-well plate (e.g., a 24 well plate at a concentration of 50,000 osteoblasts per well) in alpha MEM (modified Eagle's medium, Gibco, Inc., NY) containing 10% FBS (fetal bovine serum), L-glutamine and penicillin/streptomycin. The cells are incubated for 24 hours at 37°C, at which time the growth medium is replaced with alpha MEM containing 1% FBS and the cells incubated for an additional 24 hours so that cells are in serum-deprived growth medium at the time of the experiment.

The cultured cells are divided into four groups: (1) wells which receive, for example, 0.1, 1.0, 10.0, 40.0 and 80.0 ng of the OP-1 morphogen analog (mutein), (2) wells which receive 0.1,

1.0, 10.0 and 40.0 ng of wild type OP-1; (3) wells which receives 0.1, 1.0, 10.0, and 40.0 ng of TGF- β , and (4) the control group, which receive no growth factors. The cells then are incubated for an additional 18 hours after which the wells are pulsed with 2mCi/well of ^3H -thymidine and incubated for six more hours. The excess label then is washed off with a cold solution of 0.15 M NaCl and then 250 ml of 10% trichloroacetic acid is added to each well and the wells incubated at room temperature for 30 minutes. The cells then are washed three times with cold distilled water, and lysed by the addition of 250 ml of 1% sodium dodecyl sulfate (SDS) for a period of 30 minutes at 37°C. The resulting cell lysates are harvested using standard means and the incorporation of ^3H -thymidine into cellular DNA determined by liquid scintillation as an indication of mitogenic activity of the cells. In the experiment, it is contemplated that the OP-1 morphogen analog construct (mutein), like natural OP-1, will stimulate ^3H -thymidine incorporation into DNA, and therefore promote osteoblast cell proliferation. In contrast, the effect of the TGF- β is expected to be transient and biphasic. Furthermore, it is contemplated that at higher concentrations, TGF- β will have no significant effect on osteoblast cell proliferation.

The *in vitro* effect of the OP-1 morphogen analog on osteoblast proliferation also may be evaluated using human primary osteoblasts (obtained from bone tissue of a normal adult patient and prepared as described above) and on human osteosarcoma-derived cell lines.

B. Progenitor Cell Stimulation.

The following example is designed to demonstrate the ability of OP-1 morphogen analogs to stimulate the proliferation of mesenchymal progenitor cells. Useful naive stem cells include pluripotent stem cells, which may be isolated from bone marrow or umbilical cord blood using conventional methodologies, (see, for example, Faradji *et al.* (1988) *Vox Sang.* 55 (3): 133-138 or Broxmeyer *et al.* (1989) *Proc. Natl. Acad. Sci. USA.* 86: 3828-3832), as well as naive stem cells obtained from blood. Alternatively, embryonic cells (e.g., from a cultured mesodermal cell line) may be used.

Another method for obtaining progenitor cells and for determining the ability of OP-1 morphogen analogs to stimulate cell proliferation is to capture progenitor cells from an *in vivo* source. For example, a biocompatible matrix material able to allow the influx of migratory progenitor cells may be implanted at an *in vivo* site long enough to allow the influx of migratory

- 52 -

progenitor cells. For example, a bone-derived, guanidine-extracted matrix, formulated as disclosed for example in Sampath *et al.* (1983) *Proc. Natl. Acad. Sci. USA* 80: 6591-6595, or U.S. Patent No. 4,975,526, may be implanted into a rat at a subcutaneous site, essentially following the method of Sampath *et al.* After three days the implant is removed, and the progenitor cells associated with the matrix dispersed and cultured.

Progenitor cells, however obtained, then are incubated *in vitro* with the candidate OP-1 morphogen analog under standard cell culture conditions, such as those described hereinbelow. In the absence of external stimuli, the progenitor cells do not, or only minimally, proliferate on their own in culture.

However, progenitor cells cultured in the presence of a biologically active OP-1 morphogen analog, like OP-1, will proliferate. Cell growth can be determined visually or spectrophotometrically using standard methods well known in the art.

C. Morphogen-Induced Cell Differentiation.

A variety of assays also can be used to determine OP-1 based morphogen analog-induced cellular differentiation.

(1) Embryonic Mesenchyme Differentiation

As with natural OP-1, it is contemplated that the OP-1 morphogen analog (mucin) can induce cell differentiation. The ability of OP-1 morphogen analogs to induce cell differentiation can be demonstrated by culturing early mesenchymal cells in the presence of OP-1 morphogen analog and then studying the histology of the cultured cells by staining with toluidine blue using standard cell culturing and cell staining methodologies well described in the art. For example, it is known that rat mesenchymal cells destined to become mandibular bone, when separated from the overlying epithelial cells at stage 11 and cultured *in vitro* under standard tissue culture conditions, e.g., in a chemically defined, serum-free medium, containing for example, 67% DMEM (Dulbecco's modified Eagle's medium), 22% F-12 medium, 10mM Hepes pH 7, 2mM glutamine, 50 mg/ml transferrin, 25 mg/ml insulin, trace elements, 2mg/ml bovine serum albumin coupled to oleic acid, with HAT (0.1 mM hypoxanthine, 10mM aminopterin, 12 mM thymidine, will not continue to differentiate. However, if these same cells are left in contact with the overlying endoderm for an additional day, at which time they become stage 12 cells, they

will continue to differentiate on their own *in vitro* to form chondrocytes. Further differentiation into osteoblasts and, ultimately, mandibular bone, requires an appropriate local environment, e.g., a vascularized environment.

It is anticipated that, as with natural OP-1, stage 11 mesenchymal cells, cultured *in vitro*
5 in the presence of OP-1 morphogen analog (mutein), e.g., 10-100 ng/ml, will continue to differentiate *in vitro* to form chondrocytes just as they continue to differentiate *in vitro* if they are cultured with the cell products harvested from the overlying endodermal cells. This experiment can be performed with different mesenchymal cells to demonstrate the cell differentiation capability of OP-1 morphogen analog in different tissues.

10 As another example of morphogen-induced cell differentiation, the ability of OP-1 morphogen analogs to induce osteoblast differentiation can be demonstrated *in vitro* using primary osteoblast cultures, or osteoblast-like cells lines, and assaying for a variety of bone cell markers that are specific markers for the differentiated osteoblast phenotype, e.g., alkaline phosphatase activity, parathyroid hormone-mediated cyclic AMP (cAMP) production,
15 osteocalcin synthesis, and enhanced mineralization rates.

(2) Induction of Alkaline Phosphatase Activity in Osteoblasts.

Cultured osteoblasts in serum-free medium are incubated with a range of OP-1 morphogen analog concentrations, for example, 0.1, 1.0, 10.0, 40.0 or 80.0 ng OP-1 morphogen analog/ml medium; or with a similar concentration range of natural OP-1 or TGF- β . After a 72
20 hour incubation the cell layer is extracted with 0.5 ml of 1% Triton X-100. The resultant cell extract is centrifuged, and 100 μ l of the extract is added to 90 μ l of para-nitroso-phenylphosphate (PNPP)/glycerine mixture and incubated for 30 minutes in a 37°C water bath and the reaction stopped with 100 μ l NaOH. The samples then are run through a plate reader (e.g., Dynatech MR700 plate reader, and absorbance measured at 400 nm, using p-nitrophenol as
25 a standard) to determine the presence and amount of alkaline phosphate activity. Protein concentrations are determined by the BioRad method. Alkaline phosphatase activity is calculated in units/mg protein, where 1 unit=1 nmol p-nitrophenol liberated/30 minutes at 37°C.

It is contemplated that the OP-1 morphogen analog, like natural OP-1, will stimulate the production of alkaline phosphatase in osteoblasts thereby promoting the growth and expression

of the osteoblast differentiated phenotype. The long term effect of OP-1 morphogen analog on the production of alkaline phosphatase by rat osteoblasts also can be demonstrated as follows.

Rat osteoblasts are prepared and cultured in multi-well plates as described above. In this example six sets of 24 well plates are plated with 50,000 rat osteoblasts per well. The wells in each plate, prepared as described above, then are divided into three groups: (1) those which receive, for example, 1 ng of OP-1 morphogen analog per ml of medium; (2) those which receive 40 ng of OP-1 morphogen analog per ml of medium; and (3) those which receive 80 ng of OP-1 morphogen analog per ml of medium. Each plate then is incubated for different lengths of time: 0 hours (control time), 24 hours, 48 hours, 96 hours, 120 hours and 144 hours. After each incubation period, the cell layer is extracted with 0.5 ml of 1% Triton X-100. The resultant cell extract is centrifuged, and alkaline phosphatase activity determined using para-nitrophenylphosphate (PNPP), as above. It is contemplated that the OP-1 morphogen analog, like natural OP-1, will stimulate the production of alkaline phosphatase in osteoblasts in a dose-dependent manner so that increasing doses of OP-1 morphogen analog will further increase the level of alkaline phosphatase production. Moreover, it is contemplated that the OP-1 morphogen analog-stimulated elevated levels of alkaline phosphatase in the treated osteoblasts will last for an extended period of time.

(3) Induction of PTH-Mediated cAMP

This experiment is designed to test the effect of OP-1 morphogen analogs on parathyroid hormone-mediated cAMP production in rat osteoblasts *in vitro*. Briefly, rat osteoblasts are prepared and cultured in a multiwell plate as described above. The cultured cells then are divided into four groups: (1) wells which receive, for example, 1.0, 10.0 and 40.0 ng OP-1 morphogen analog/ml medium); (2) wells which receive for example, natural OP-1, at similar concentration ranges; (3) wells which receive for example, TGF- β , at similar concentration ranges; and (4) a control group which receives no growth factors. The plate then is incubated for another 72 hours. At the end of the 72 hours the cells are treated with medium containing 0.5% bovine serum albumin (BSA) and 1mM 3-isobutyl-1-methylxanthine for 20 minutes followed by the addition into half of the wells of human recombinant parathyroid hormone (hPTH, Sigma, St. Louis) at a concentration of 200 ng/ml for 10 minutes. The cell layer then is extracted from each well with 0.5 ml of 1% Triton X-100. The cAMP levels then are determined using a

radioimmunoassay kit (e.g., Amersham, Arlington Heights, Illinois). It is contemplated that OP-1 morphogen analog alone, like OP-1, will stimulate an increase in the PTH-mediated cAMP response, thereby promoting the growth and expression of the osteoblast differentiated phenotype.

5 (4) Induction of Osteocalcin Production.

Osteocalcin is a bone-specific protein synthesized by osteoblasts which plays an integral role in the rate of bone mineralization *in vivo*. Circulating levels of osteocalcin in serum are used as a marker for osteoblast activity and bone formation *in vivo*. Induction of osteocalcin synthesis in osteoblast-enriched cultures can be used to demonstrate OP-1 morphogen analog efficacy *in*
10 *vitro*.

Rat osteoblasts are prepared and cultured in a multi-well plate as above. In this experiment the medium is supplemented with 10%FBS, and on day 2, cells are fed with fresh medium supplemented with fresh 10 mM β -glycerophosphate (Sigma, Inc.). Beginning on day 5 and twice weekly thereafter, cells are fed with a complete mineralization medium containing all
15 of the above components plus fresh L(+)-ascorbate, at a final concentration of 50mg/ml medium. OP-1 morphogen analog then is added to the wells directly, e.g., in 50% acetonitrile (or 50% ethanol) containing 0.1% trifluoroacetic acid (TFA), at no more than 5ml morphogen analog/ml medium. Control wells receive solvent vehicle only. The cells then are re-fed and the conditioned medium sample diluted 1:1 in standard radioimmunoassay buffer containing
20 standard protease inhibitors and stored at -20°C until assayed for osteocalcin. Osteocalcin synthesis is measured by standard radioimmunoassay using a commercially available osteocalcin-specific antibody.

Mineralization is determined on long term cultures (13 day) using a modified von Kossa staining technique on fixed cell layers: cells are fixed in fresh 4% paraformaldehyde at 23°C for
25 10 min, following rinsing cold 0.9% NaCl. Fixed cells then are stained for endogenous alkaline phosphatase at pH 9.5 for 10 min, using a commercially available kit (Sigma, Inc.). Purple stained cells then are dehydrated with methanol and air dried. After 30 min incubation in 3% AgNO₃ in the dark, H₂O-rinsed samples are exposed for 30 sec to 254 nm UV light to develop

- 56 -

the black silver-stained phosphate nodules. Individual mineralized foci (at least 20 mm in size) are counted under a dissecting microscope and expressed as nodules/culture.

It is contemplated that the OP-1 morphogen analog, like natural OP-1, will stimulate osteocalcin synthesis in osteoblast cultures. Furthermore, it is contemplated that the increased
5 osteocalcin synthesis in response to OP-1 morphogen analog will be in a dose dependent manner thereby showing a significant increase over the basal level after 13 days of incubation. Enhanced osteocalcin synthesis also can be confirmed by detecting the elevated osteocalcin mRNA message (20-fold increase) using a rat osteocalcin-specific probe. In addition, the increase in osteocalcin synthesis correlates with increased mineralization in long term osteoblast cultures as
10 determined by the appearance of mineral nodules. It is contemplated also that OP-1 morphogen analog, like natural OP-1, will increase significantly the initial mineralization rate as compared to untreated cultures.

(5) Morphogen-Induced CAM Expression

Members of the BMP/OP family (see Figure 6) induce CAM expression, particularly N-
15 CAM expression, as part of their induction of morphogenesis (see copending U.S.S.N. 922,813). CAMs are morphoregulatory molecules identified in all tissues as an essential step in tissue development. N-CAMs, which comprise at least 3 isoforms (N-CAM-180, N-CAM-140 and N-CAM-120, where "180", "140" and "120" indicate the apparent molecular weights of the isoforms as measured by SDS polyacrylamide gel electrophoresis) are expressed at least
20 transiently in developing tissues, and permanently in nerve tissue. Both the N-CAM-180 and N-CAM-140 isoforms are expressed in both developing and adult tissue. The N-CAM-120 isoform is found only in adult tissue. Another neural CAM is L1.

The ability of OP-1 based morphogen analogs to stimulate CAM expression may be demonstrated using the following protocol, using NG108-15 cells. NG108-15 is a transformed
25 hybrid cell line (neuroblastoma x glioma, America Type Culture Collection (ATCC), Rockville, MD), exhibiting a morphology characteristic of transformed embryonic neurons. As described in Example D, below, untreated NG108-15 cells exhibit a fibroblastic, or minimally differentiated, morphology and express only the 180 and 140 isoforms of N-CAM normally associated with a developing cell. Following treatment with members of the vg/dpp subgroup these cells exhibit a

morphology characteristic of adult neurons and express enhanced levels of all three N-CAM isoforms.

In this example, NG108-15 cells are cultured for 4 days in the presence of increasing concentrations of either the OP-1 morphogen analog or natural OP-1 using standard culturing procedures, and standard Western blots are performed on whole cell extracts. N-CAM isoforms are detected with an antibody which crossreacts with all three isoforms, mAb H28.123, obtained from Sigma Chemical Co., St. Louis, the different isoforms being distinguishable by their different mobilities on an electrophoresis gel. Control NG108-15 cells (untreated) express both the 140 kDa and the 180 kDa isoforms, but not the 120 kDa, as determined by Western blot analyses using up to 100 mg of protein. It is contemplated that treatment of NG108-15 cells with OP-1 morphogen analog, like natural OP-1 may result in a dose-dependent increase in the expression of the 180 kDa and 140 kDa isoforms, as well as the induction of the 120 kDa isoform. In addition, it is contemplated that the OP-1 morphogen analog, like natural OP-1-induced CAM expression may correlate with cell aggregation, as determined by histology.

15 (D) OP-1 Morphogen Analog-Induced Redifferentiation of Transformed Phenotype

It is contemplated that OP-1 morphogen analog, like natural OP-1, also induces redifferentiation of transformed cells to a morphology characteristic of untransformed cells. The examples provided below detail morphogen-induced redifferentiation of a transformed human cell line of neuronal origin (NG108-15); as well as mouse neuroblastoma cells (N1E-115), and human embryo carcinoma cells, to a morphology characteristic of untransformed cells.

As described above, NG108-15 is a transformed hybrid cell line produced by fusing neuroblastoma x glioma cells (obtained from ATCC, Rockville, MD), and exhibiting a morphology characteristic of transformed embryonic neurons, e.g., having a fibroblastic morphology. Specifically, the cells have polygonal cell bodies, short, spike-like processes and make few contacts with neighboring cells. Incubation of NG108-15 cells, cultured in a chemically defined, serum-free medium, with 0.1 to 300 ng/ml of morphogen analog or natural OP-1 for four hours induces an orderly, dose-dependent change in cell morphology.

For example, NG108-15 cells are subcultured on poly-L-lysine coated 6 well plates. Each well contains 40-50,000 cells in 2.5 ml of chemically defined medium. On the third day,

2.5 ml of OP-1 morphogen analog or natural OP-1 in 60% ethanol containing 0.025% trifluoroacetic is added to each well. The media is changed daily with new aliquots of morphogen. It is contemplated that OP-1 morphogen analog, like OP-1, may induce a dose-dependent redifferentiation of the transformed cells, including a rounding of the soma, an
5 increase in phase brightness, extension of the short neurite processes, and other significant changes in the cellular ultrastructure. After several days it is contemplated also that treated cells may begin to form epithelioid sheets that then become highly packed, multi-layered aggregates, as determined visually by microscopic examination.

Moreover, it is contemplated that the redifferentiation may occur without any associated
10 changes in DNA synthesis, cell division, or cell viability, making it unlikely that the morphologic changes are secondary to cell differentiation or a toxic effect of the morphogen. In addition, it is contemplated that the morphogen analog-induced redifferentiation may not inhibit cell division, as determined by ³H-thymidine uptake, unlike other molecules such as butyrate, DMSO, retinoic acid or Forskolin, which have been shown to stimulate differentiation of
15 transformed cells in analogous experiments. Thus, it is contemplated that the OP-1 morphogen analog, like natural OP-1, may maintain cell stability and viability after inducing redifferentiation.

The morphogen described herein would, therefore, provide useful therapeutic agents for the treatment of neoplasias and neoplastic lesions of the nervous system, particularly in the
20 treatment of neuroblastomas, including retinoblastomas, and gliomas.

(E) Maintenance of Phenotype.

OP-1 morphogen analogs, like natural OP-1, also may be used to maintain a cell's differentiated phenotype. This application is particularly useful for inducing the continued expression of phenotype in senescent or quiescent cells.

(1) In Vitro Model for Phenotypic Maintenance

25 The phenotypic maintenance capability of morphogens is determined readily. A number of differentiated cells become senescent or quiescent after multiple passages *in vitro* under standard tissue culture conditions well described in the art (e.g., *Culture of Animal Cells: A Manual of Basic Techniques*, C.R. Freshney, ed., Wiley, 1987). However, if these cells are

cultivated *in vitro* in association with a morphogen such as OP-1, cells are stimulated to maintain expression of their phenotype through multiple passages. For example, the alkaline phosphatase activity of cultured osteoblasts, such as cultured osteosarcoma cells and calvaria cells, is significantly reduced after multiple passages *in vitro*. However, if the cells are cultivated in the presence of OP-1, alkaline phosphatase activity is maintained over extended periods of time. Similarly, phenotypic expression of myocytes also is maintained in the presence of a morphogen. In the experiment, osteoblasts are cultured as described in Example A. The cells are divided into groups, incubated with varying concentrations of either OP-1 morphogen analog or natural OP-1 (e.g., 0-300 ng/ml) and passaged multiple times (e.g., 3-5 times) using standard methodology. Passaged cells then are tested for alkaline phosphatase activity, as described in Example C as an indication of differentiated cell metabolic function. It is contemplated that osteoblasts cultured in the absence of OP-1 morphogen analog may have reduced alkaline phosphatase activity, as compared to OP-1 morphogen analog, or natural OP-1-treated cells.

(2) *In Vivo Model for Phenotypic Maintenance.*

Phenotypic maintenance capability also may be demonstrated *in vivo*, using a standard rat model for osteoporosis. Long Evans female rats (Charles River Laboratories, Wilmington, MA) are sham-operated (control animals) or ovariectomized using standard surgical techniques to produce an osteoporotic condition resulting from decreased estrogen production. Following surgery, e.g., 200 days after ovariectomy, rats are systemically provided with phosphate buffered saline (PBS) or morphogen, (e.g., OP-1 morphogen analog, or natural OP-1, 1-100 mg) for 21 days (e.g., by daily tail vein injection.) The rats then are sacrificed and serum alkaline phosphatase levels, serum calcium levels, and serum osteocalcin levels are determined, using standard methodologies as described therein and above. It is contemplated that the OP-1 morphogen analog treated rats, like the OP-1 treated rats may exhibit elevated levels of osteocalcin and alkaline phosphatase activity. It is contemplated also that histomorphometric analysis on the tibial diaphyseal bone may show improved bone mass in OP-1 morphogen analog-treated animals as compared with untreated, ovariectomized rats.

F. Proliferation of Progenitor Cell Populations

Progenitor cells may be stimulated to proliferate *in vivo* or *ex vivo*. It is contemplated that cells may be stimulated *in vivo* by injecting or otherwise providing a sterile preparation containing the OP-1 morphogen analog into the individual. For example, the hematopoietic pluripotent stem cell population of an individual may be stimulated to proliferate by injecting or otherwise providing an appropriate concentration of OP-1 morphogen analog to the individual's bone marrow.

Progenitor cells may be stimulated *ex vivo* by contacting progenitor cells of the population to be enhanced with a morphogenically active OP-1 morphogen analog under sterile conditions at a concentration and for a time sufficient to stimulate proliferation of the cells. Suitable concentrations and stimulation times may be determined empirically, essentially following the procedure described in Example A, above. It is contemplated that a OP-1 morphogen analog concentration of between about 0.1-100 ng/ml and a stimulation period of from about 10 minutes to about 72 hours, or, more generally, about 24 hours, typically should be sufficient to stimulate a cell population of about 10^4 to 10^6 cells. The stimulated cells then may be provided to the individual as, for example, by injecting the cells to an appropriate *in vivo* locus. Suitable biocompatible progenitor cells may be obtained by any of the methods known in the art or described hereinabove.

G. Regeneration of Damaged or Diseased Tissue

It is contemplated that OP-1 morphogen analogs may be used to repair diseased or damaged mammalian tissue. The tissue to be repaired preferably is assessed first, and excess necrotic or interfering scar tissue removed as needed, e.g., by ablation or by surgical, chemical, or other methods known in the medical arts.

OP-1 morphogen analog then may be provided directly to the tissue locus as part of a sterile, biocompatible composition, either by surgical implantation or injection. The morphogen analog also may be provided systemically, as by oral or parenteral administration. Alternatively, a sterile, biocompatible composition containing progenitor cells stimulated by a morphogenically active OP-1 morphogen analog may be provided to the tissue locus. The existing tissue at the

locus, whether diseased or damaged, provides the appropriate matrix to allow the proliferation and tissue-specific differentiation of progenitor cells. In addition, a damaged or diseased tissue locus, particularly one that has been further assaulted by surgical means, provides a morphogenically permissive environment. Systemic provision of OP-1 morphogen analog may be sufficient for certain applications (e.g., in the treatment of osteoporosis and other disorders of the bone remodeling cycle).

In some circumstances, particularly where tissue damage is extensive, the tissue may not be capable of providing a sufficient matrix for cell influx and proliferation. In these instances, it may be necessary to provide progenitor cells stimulated by the OP-1 morphogen analog to the tissue locus in association with a suitable, biocompatible, formulated matrix, prepared by any of the means described below. The matrix preferably is *in vivo* biodegradable. The matrix also may be tissue-specific and/or may comprise porous particles having dimensions within the range of 70-850 μm , most preferably 150-420 μm .

OP-1 morphogen analog also may be used to prevent or substantially inhibit immune/inflammatory response-mediated tissue damage and scar tissue formation following an injury. OP-1 morphogen analog may be provided to a newly injured tissue locus, to induce tissue morphogenesis at the locus, preventing the aggregation of migrating fibroblasts into non-differentiated connective tissue. Preferably the OP-1 morphogen analog may be provided as a sterile pharmaceutical preparation injected into the tissue locus within five hours of the injury. Where an immune/inflammatory response is unavoidably or deliberately induced, as part of, for example, a surgical or other aggressive clinical therapy, OP-1 morphogen analog preferably may be provided prophylactically to the patient prior to, or concomitant with, the therapy.

Described below is a protocol for demonstrating whether a OP-1 morphogen analog induces tissue morphogenesis in bone.

(1) OP-1 Morphogen Analog-Induced Bone Morphogenesis.

A particularly useful mammalian tissue model system for demonstrating and evaluating the morphogenic activity of a morphogen analog is the endochondral bone tissue morphogenesis model known in the art and described, for example, in U.S. Pat. No. 4,968,590, incorporated herein by reference. The ability to induce endochondral bone formation includes the ability to

induce proliferation and differentiation of progenitor cells into chondroblasts and osteoblasts, the ability to induce cartilage matrix formation, cartilage calcification, and bone remodeling, and the ability to induce formation of an appropriate vascular supply and hematopoietic bone marrow differentiation.

5 The local environment in which the morphogenic material is placed is important for tissue morphogenesis. As used herein, "local environment" is understood to include the tissue structural matrix and the environment surrounding the tissue. For example, in addition to needing an appropriate anchoring substratum for their proliferation, the cells stimulated by morphogens need signals to direct the tissue-specificity of their differentiation. These signals
10 vary for the different tissues and may include cell surface markers. In addition, vascularization of new tissue requires a local environment which supports vascularization.

 The following sets forth various procedures for evaluating the *in vivo* morphogenic utility of OP-1 morphogen analogs and OP-1 morphogen analog containing compositions. The compositions may be injected or surgically implanted in a mammal, following any of a number
15 of procedures well known in the art. For example, surgical implant bioassays may be performed essentially following the procedure of Sampath *et al.* (1983) *Proc. Natl. Acad. Sci. USA* 80: 6591-6595 and U.S. Pat No. 4,968,590.

 Histological sectioning and staining is preferred to determine the extent of morphogenesis *in vivo*, particularly in tissue repair procedures. Excised implants are fixed in Bouins Solution,
20 embedded in paraffin, and cut into 6-8 μ m sections. Staining with toluidine blue or hemotoxylin/eosin demonstrates clearly the ultimate development of the new tissue. Twelve day implants are usually sufficient to determine whether the implants contain newly induced tissue.

 Successful implants exhibit a controlled progression through the stages of induced tissue development allowing one to identify and follow the tissue-specific events that occur. For
25 example, in endochondral bone formation the stages include: (1) leukocytes on day one; (2) mesenchymal cell migration and proliferation on days two and three; (3) chondrocyte appearance on days five and six; (4) cartilage matrix formation on day seven; (5) cartilage calcification on day eight; (6) vascular invasion, appearance of osteoblasts, and formation of new bone on days nine and ten; (7) appearance of osteoclastic cells, and the commencement of bone

remodeling and dissolution of the implanted matrix on days twelve to eighteen; and
(8) hematopoietic bone marrow differentiation in the resulting ossicles on day twenty-one.

In addition to histological evaluation, biological markers may be used as markers for tissue morphogenesis. Useful markers include tissue-specific enzymes whose activities may be
5 assayed (e.g., spectrophotometrically) after homogenization of the implant. These assays may be useful for quantitation and for rapidly obtaining an estimate of tissue formation after the implants are removed from the animal. For example, alkaline phosphatase activity may be used as a marker for osteogenesis.

Incorporation of systemically provided OP-1 morphogen analog may be followed using
10 labeled protein (e.g., radioactively labeled) and determining its localization in the new tissue, and/or by monitoring their disappearance from the circulatory system using a standard labeling protocol and pulse-chase procedure. OP-1 morphogen analog also may be provided with a tissue-specific molecular tag, whose uptake may be monitored and correlated with the concentration of OP-1 morphogen analog provided. As an example, ovary removal in female
15 rats results in reduced bone alkaline phosphatase activity, and renders the rats predisposed to osteoporosis (as described in Example E). If the female rats now are provided with OP-1 morphogen analog, a reduction in the systemic concentration of calcium may be seen, which correlates with the presence of the provided OP-1 morphogen analog and which is anticipated to correspond with increased alkaline phosphatase activity.

20 Example 2. Enhancing the Solubility of a hOP-1 Dimer.

As described in section V.A.(ii), *supra*, it is contemplated that the solubility of the hOP-1 dimer can be enhanced by replacing hydrophobic amino acid residues located at the solvent accessible surface of hOP-1 dimer with more polar or hydrophilic amino acid residues. This example provides a description of such an approach.

25 A Sma I to Bam HI fragment of the human OP-1 cDNA as described in Ozkaynak *et al.* (1990) *supra* is cloned into a vector to produce a plasmid similar to the plasmid called pW24 in International Application PCT/US94/12063, the disclosure of which is incorporated herein by reference. The pW24 plasmid contains OP-1 cDNA under the transcriptional control of the CMV (cytomegalovirus) immediate early promoter. The selective marker on pW24 is the

- 64 -

neomycin gene which provides resistance to the cytostatic drug G418. The pW24 plasmid also employs an SV40 origin of replication (ori). The early SV40 promoter is used to drive transcription of the neomycin marker gene.

5 Then, the alanine at position 63 is mutated to a serine by site-directed mutagenesis using, for example, synthetic oligonucleotides and either PCR or the site-directed mutagenesis methods. See, for example, Kunkel *et al.* (1985) *Proc. Natl. Acad. Sci. USA* 822: 488; Kunkel *et al.* (1985) *Meth. Enzymol.* 154: 367 and U.S. Patent No. 4,873,192. The resulting mutation is confirmed by dideoxy sequencing.

10 Two additional vectors have been developed for use in a triple transfection procedure along with pW24 to enhance OP-1 expression. One of the vectors employs the adenovirus E1A gene under the VA1 gene as translation stimulation for the gene DHFR gene. The other vector employs the adenovirus E1A gene under the control of the thymidine kinase promoter as a transactivating transcription activator. Both additional vectors, known as pH1130 and pH1176, as well as preferred transfection and screening procedures are described in International
15 Application PCT/US94/12063.

Briefly, triple transfections are performed using the calcium phosphate coprecipitation procedure. CHO cells are cultured in α MEM, containing 5% or 10% fetal bovine serum (FBS), non-essential amino acids, glutamine and antibiotics: penicillin and streptomycin. Stable cell line transfections are carried out by seeding $1-2 \times 10^6$ cells in a 9 cm. petri dish. Following an
20 incubation period of up to 24-hour, each petri dish is transfected with 10-30 μ g total vector DNA in equimolar amounts, by calcium phosphate coprecipitation followed by glycerol shock using standard methodology. Cells are incubated at 37°C in growth medium for 24 hours, then transferred to selection medium. All cultures are fed once or twice weekly with fresh selective medium. After 10 - 21 days, resistant colonies are picked and assayed for protein production.

25 Approximately 30 individual clones are selected, transferred to a 24-well petri dish, and allowed to grow to confluence in serum-containing media. The conditioned media from all surviving clones is screened for protein production using a standard ELISA (enzyme-linked immunosorbent assay) or Western blot. The methodologies for these assay protocols as well as

for generating antibodies for use in these assays are well described in the art (see e.g., Ausubel, *supra*).

Under such conditions, the VA1 and E1A genes typically act synergistically to enhance OP1 expression in unamplified transfected CHO cells. Candidate cell lines identified by the
5 screening protocol, then are seeded on ten 100 mm petri dishes at a cell density of either 50 or 100 cells per plate, and with a higher drug concentration (e.g., 1.0- μ m).

After 10-21 days of growth, the clones are isolated using cloning cylinders and standard procedures, and cultured in 24-well plates. Then, clones are screened for OP-1 expression by Western immunoblots using standard procedures, and OP-1 expression levels compared to
10 parental lines. Candidate cells showing higher protein production than cells of parental lines then are replated and grown in the presence of a still higher drug concentration (e.g., 5-20 μ m). Generally, no more than 2-3 rounds of these "amplification" cloning steps are necessary to achieve cell lines with high protein productivity. Useful high producing cell lines may be further subcloned to improve cell line homogeneity and product stability.

15 A currently preferred method of large scale protein production e.g., at least 2 liters, is by suspension culture of the host Chinese hamster ovary (CHO) cells. CHO cells prefer attachment but can be adapted to grow in suspension mode of cultivation. The cells are trypsinized from a culture dish, introduced to growth media containing 10% FBS and completely suspended to produce a single cell suspension. The single cell suspension is introduced to a spinner flask and
20 placed in a 37°C 95% air/5% CO₂ humidified incubator. Over a period of time the cells are subcultured in medium with descending concentrations of serum.

Specifically, the adapted cells are introduced into a 3L spinner flask at an initial viable cell density of approximately 2×10^5 cells/ml. Preferred culture medium is DMEM/F-12 (1:1) (GIBCO, New York) supplemented with 2% FBS, and preferred agitation is approximately 50-
25 60 rpm with a paddle impeller. After 7 days, the culture media is harvested, centrifuged at 1500 rpm and the clarified conditioned media stored at 4°C.

A representative purification scheme for purifying recombinant morphogenic protein involves three chromatographic steps (S-Sepharose, phenyl-Sepharose and C-18 HPLC) and is described in International Application PCT/US94/12063. Morphogen analog containing culture

- 66 -

media is diluted to 6M urea, 0.05M NaCl, 13mM HEPES, pH 7.0 and loaded onto an S-Sephacrose column, which acts as a strong cation exchanger. The column subsequently is developed with two salt elutions. The first elution employs a solution containing 0.1M NaCl, and the second elution employs a buffer containing 6M urea, 0.3M NaCl, 20mM HEPES, pH 7.0.

Ammonium sulfate is added to the 0.3M NaCl fraction to give a solution containing 6M urea, 1M $(\text{NH}_4)_2\text{SO}_4$, 0.3M NaCl, 20mM HEPES, pH 7.0. Then, the sample is loaded onto a phenyl-Sepharose column in the presence of 1M $(\text{NH}_4)_2\text{SO}_4$. Then, the column is developed with two step elutions using decreasing concentrations of ammonium sulfate. The first elution employs 0.6M $(\text{NH}_4)_2\text{SO}_4$ and the second elution employs 6M urea, 0.3M NaCl, 20mM HEPES, pH 7.0 buffer. The material harvested from the second elution is dialyzed against water, followed by 30% acetonitrile (0.1% TFA), and then applied to a C-18 reverse phase HPLC column. Purified morphogen analog is harvested from the HPLC column.

The enhanced solubility of the resulting morphogen analog is measured by comparing the partition coefficient of the Ala 63-> Ser 63 mutein versus wild type hOP-1 dimer. It is contemplated that the Ala 63-> Ser 63 mutein has a higher solubility than native hOP-1. It is contemplated that, additional muteins having multiple hydrophobic to hydrophilic substitutions can be produced and characterized using the protocols described in this Example. The biological activity of the resulting morphogen analogs can be determined using one or more of the OP-1 activity assays described Example 1.

Example 3. Biological Activity of Finger 1, Finger 2, and Heel Peptides

The hOP-1-based peptides described in this example were produced and characterized prior to determination of the three-dimensional structure of hOP-1. These peptides either agonize or antagonize the biological activity of hOP-1. It is contemplated that, further refinements based upon the hOP-1 crystal structure, for example, the choice of more suitable sites for cyclizing peptides which constrain the peptide into a conformation that more closely mimics the shape of the corresponding region in hOP-1, may be used to further enhance the agonistic or antagonistic properties of such hOP-1-based peptides.

All of the peptides used in the following experiments, as well as their relationships with the mature hOP-1 amino acid sequence, are shown in Fig. 12. The finger 1-based peptides are designated F1 - 2; the heel-based peptides are designated H-1, H-n2 and H-c2; and the finger 2-based peptides are designated F2-2, and F2-3. Potential intra-peptide disulfide linkages are shown for each peptide. All the peptides were synthesized on a standard peptide synthesizer in accordance with the manufacturer's instructions. The peptides were deprotected, cyclized by oxidation, and then cleaved from resin prior to use.

In a first series of experiments, increasing concentrations of peptides F2-2 (Fig. 13A), F2-3 (Fig. 13B), Hn-2 (Fig. 13C) and Hc-2 (Fig. 13D) were added to ROS cells either alone (open bars) or in combination with 40ng/ml soluble OP-1 (filled bars) and their effect on alkaline phosphatase activity measured. Soluble OP-1 is the form of OP-1 in which the pro-domain is still attached to the mature portion of OP-1 (see WO94/03600). A basal alkaline phosphatase activity is shown by the line and represents the alkaline phosphatase activity of cells incubated in the absence of both soluble OP-1 and peptide.

In Fig. 13A, peptide F2-2 at a concentration of about 60 μ M appears to double the basal alkaline phosphatase level and, in the presence of soluble OP-1, increases alkaline phosphatase activity by about 20% relative to soluble OP-1 alone. In Fig. 13B, peptide F2-3 at a concentration of about 0.01 μ M appears to increase the basal alkaline phosphatase level and, in the presence of soluble OP-1, increases alkaline phosphatase activity by about 20% relative to

soluble OP-1 alone. Accordingly, both peptides F2-2 and F2-3, in the alkaline phosphatase assay, appear to act as weak OP-1 agonists.

In Fig. 13C, peptide H-n2 displays little or no effect on alkaline phosphatase activity either alone or in combination with soluble OP-1. Fig. 13D, peptide H-c2, at concentrations greater than about 5 μ M, appears to antagonize the activity of soluble OP-1.

In a second series of experiments, the ability of unlabeled soluble OP-1 and unlabeled peptides F1-2, F2-2, F2-3, H-n2 and H-c2 to displace 125 I labeled soluble OP-1 from ROS cell membranes was measured. The activities of peptides F2-2 and F2-3 relative to soluble OP-1 are shown in Fig. 14A, and the activities of peptides F1-2, H-n2 and H-c2 relative to soluble OP-1 are shown in Fig. 14B. OP-1 receptor-enriched plasma membranes of ROS cells were incubated for 20 hrs at 4°C with 125 I-labeled soluble OP-1 and unlabeled peptide. Receptor bound material was separated from unbound material by centrifugation at 39,500 x g. The resulting pellet was harvested and washed with 50mM HEPES buffer, pH7.4 containing 5mM $MgCl_2$ and 1mM $CaCl_2$. Radioactivity remaining in the pellet was determined by means of a gamma counter.

In Fig. 14A, peptide F2-2 (filled circles) soluble competes with soluble OP-1 with an Effective Dose₅₀ (ED₅₀) of about 1 μ M, but cannot completely displace soluble OP-1. ED₅₀ is the concentration of peptide to produce half maximal displacement of labeled soluble OP-1. Peptide F2-3 (filled triangles) competes and is able to completely displace soluble OP-1 with an ED₅₀ of about 5 μ M. In Fig. 14B, peptide F1-2 (filled boxes), peptide H-n2 (open diamonds) and peptide H-c2 (open circles) all appear to exhibit little or no ability to displace iodinated soluble OP-1 from ROS cell membranes.

Although the peptide experiments appear promising, it is contemplated that resolution of the hOP-1 structure will enable the skilled practitioner to design constrained peptides that more closely mimic the receptor binding domains of human OP-1 and which are more effective at agonizing or antagonizing an hOP-1 mediated biological effect.

Example 4. Elimination of a Binding Site on the Surface of OP-1

α -2 macroglobulin, a protease scavenging protein known to bind proteins in serum and target them to the kidney for clearance from the body, binds OP-1. As described herein, α -2's interaction sites on the OP-1 protein have been mapped. Accordingly, using the database and

structural information provided herein, one can design an analog of OP-1 which eliminates one or more α -2 macroglobulin interaction sites and provide an analog having enhanced bioavailability in the body. This same strategy can be applied for identifying and/or eliminating interaction sites for other binding proteins on the OP-1 surface.

5 A. Identifying α -2 macroglobulin Binding Sites

OP-1 was determined to interact specifically with α -2 macroglobulin in a standard competition binding assay, using immobilized, commercially available α -2 macroglobulin, and labeled and unlabeled OP-1 protein. Truncated mature OP-1, wherein the first 22 amino acids have been cleaved from the mature form of OP-1 in a standard trypsin digest, bound α -2 with 10-
10 fold less affinity, indicating that the N terminal portion of the mature protein is involved in binding. This N-terminal portion of the protein, which is not part of the crystal structure, is positively charged and likely is highly flexible in solution. Elimination of this sequence does not interfere with OP-1 activity. Two cyclized peptides to all or a portion of the heel region, H-n2 and H1 (Cys₇₁ - Pro₁₀₂, where Pro₁₀₂ has been changed to a cysteine to allow a disulfide bond
15 between the two cysteines) also compete for binding; while peptides to the finger regions (F2-2, F2-3) do not compete.

α -2 macroglobulin was determined not to interfere with OP-1's ability to stimulate alkaline phosphatase activity in a ROS cell assay. Accordingly, α -2 macroglobulin binding does not appear to sterically inhibit OP-1 receptor binding.

20 B. Design of Modified OP-1 Analog

The precise α -2 macroglobulin interaction sites on OP-1 now can be mapped and an analog designed using the structure information provided herein. For example, the exact contact residues can be identified by creating model peptides like H-N2 and/or H1 in conjunction with an "alanine scan" mutagenesis program, wherein each residue is individually changed to an
25 alanine in turn, and the constructs then tested for their ability to compete for binding. Once the contact residues are mapped, an analog can be designed which eliminates the contact residues without altering the overall structure of the heel region. Specifically, a template of the region can be called up on the computer from the database, and candidate replacement residues tested. The information in Table 8 identifies particularly useful candidate residues in the heel region which

- 70 -

are solvent accessible, which likely are not available as epitopes and make good candidates for modification.

Equivalents.

5 The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting on the invention described herein. Scope of the invention is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to
10 be embraced therein.

- 71 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: KECK, PETER
GRIFFITH, DIANA L
CARLSON, WILLIAM D
RUEGER, DAVID C
SAMPATH, KUBER T
- (ii) TITLE OF INVENTION: METHODS AND COMPOSITIONS FOR
PRODUCING MORPHOGEN ANALOGS
- (iii) NUMBER OF SEQUENCES: 8
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: PATENT ADMINISTRATOR, CREATIVE
BIOMOLECULES, INC.
 - (B) STREET: 45 SOUTH STREET
 - (C) CITY: HOPKINTON
 - (D) STATE: MA
 - (E) COUNTRY: USA
 - (F) ZIP: 01748
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: PITCHER, EDMUND R
 - (B) REGISTRATION NUMBER: 27,829
 - (C) REFERENCE/DOCKET NUMBER: CRP-102
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (508) 435-9001
 - (B) TELEFAX: (508) 435-0992

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 102 amino acids
 - (B) TYPE: amino acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (ix) FEATURE:
 - (A) NAME/KEY: Protein
 - (B) LOCATION: 1..102
 - (D) OTHER INFORMATION: /product= "hOP-1"
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

- 72 -

Cys Lys Lys His Glu Leu Tyr Val Ser Phe Arg Asp Leu Gly Trp Gln
 1 5 10 15
 Asp Trp Ile Ile Ala Pro Glu Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly
 20 25 30
 Glu Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn Ala Thr Asn His Ala
 35 40 45
 Ile Val Gln Thr Leu Val His Phe Ile Asn Pro Glu Thr Val Pro Lys
 50 55 60
 Pro Cys Cys Ala Pro Thr Gln Leu Asn Ala Ile Ser Val Leu Tyr Phe
 65 70 75 80
 Asp Asp Ser Ser Asn Val Ile Leu Lys Lys Tyr Arg Asn Met Val Val
 85 90 95
 Arg Ala Cys Gly Cys His
 100

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 15 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..15
- (D) OTHER INFORMATION: /product= "PEPTIDE F1-2"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Cys Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Cys
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..32
- (D) OTHER INFORMATION: /product= "PEPTIDE H-1"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Cys Ala Phe Pro Leu Asn Ser Tyr Met Asn Ala Thr Asn His Ala Ile
 1 5 10 15

- 73 -

Val Gln Thr Leu Val His Phe Ile Asn Pro Glu Thr Val Pro Lys Cys
20 25 30

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 13 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..13
- (D) OTHER INFORMATION: /product= "PEPTIDE H-N2"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Cys Leu Asn Ser Tyr Met Asn Ala Thr Asn His Ala Cys
1 5 10

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /product= "PEPTIDE H-C2"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Cys Cys Phe Ile Asn Pro Glu Thr Val Cys Cys
1 5 10

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..11
- (D) OTHER INFORMATION: /product= "PEPTIDE F2-2"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

- 74 -

Cys Tyr Phe Asp Asp Ser Ser Asn Val Ile Cys
 1 5 10

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 16 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Peptide
- (B) LOCATION: 1..16
- (D) OTHER INFORMATION: /product= "PEPTIDE F2-3"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Cys Tyr Phe Asp Asp Ser Ser Asn Val Ile Cys Lys Lys Tyr Arg Ser
 1 5 10 15

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 98 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(ix) FEATURE:

- (A) NAME/KEY: Protein
- (B) LOCATION: 1..98
- (D) OTHER INFORMATION: /product= "TGFB2"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Cys Cys Leu Arg Pro Leu Tyr Ile Asp Phe Lys Arg Asp Leu Gly Trp
 1 5 10 15

Lys Trp Ile His Glu Pro Lys Gly Tyr Asn Ala Asn Phe Cys Ala Gly
 20 25 30

Ala Cys Pro Tyr Leu Trp Ser Ser Asp Thr Gln His Ser Arg Val Leu
 35 40 45

Ser Leu Tyr Asn Thr Ile Asn Pro Glu Ala Ser Ala Ser Pro Cys Cys
 50 55 60

Val Ser Gln Asp Leu Glu Pro Leu Thr Ile Leu Tyr Tyr Ile Gly Lys
 65 70 75 80

Thr Pro Lys Ile Glu Gln Leu Ser Asn Met Ile Val Lys Ser Cys Lys
 85 90 95

Cys Ser

CLAIMS

What is claimed is:

- 1 1. A computer system comprising:
2 (a) a memory having disposed therein atomic X-ray crystallographic co-ordinates
3 defining at least a portion of human OP-1; and
4 (b) a processor in electrical communication with the memory; the processor comprising a
5 process which generates a molecular model having a three-dimensional shape representative of at
6 least a portion of human OP-1.
- 1 2. The system of claim 1, wherein the processor further comprises a process which
2 generates the molecular model having a solvent accessible surface representative of at least a
3 portion of human OP-1.
- 1 3. The system of claim 1, wherein said co-ordinates are stored on a computer readable
2 diskette.
- 1 4. The system of claim 1, wherein the molecular model is representative of at least a portion
2 of human OP-1 finger 1 region.
- 1 5. The system of claim 1 or 4, wherein the molecular model is representative of at least a
2 portion of the human OP-1 heel region.
- 1 6. The system of claim 1 or 4, wherein the molecular model is representative of at least a
2 portion of the human OP-1 finger 2 region.
- 1 7. The system of claim 6, wherein the molecular model is representative of at least a portion
2 of the human OP-1 heel region.
- 1 8. The system of claim 1, wherein the processor further identifies a morphogenic analog
2 having a three-dimensional shape and a solvent accessible surface corresponding to at least a
3 portion of the three-dimensional shape and the solvent accessible surface of human OP-1.

1 9. The system of claim 1, wherein the processor further identifies at least one candidate
2 amino acid defined by the co-ordinates, which upon modification enhances water solubility or
3 stability of human OP-1.

1 10. A method of producing a morphogenic analog having osteogenic protein-1 (OP-1) like
2 biological activity, the method comprising the steps of:

3 (a) providing a molecular model defining a three dimensional shape representative of at
4 least a portion of human OP-1;

5 (b) identifying a candidate analog having a three dimensional shape corresponding to the
6 three dimensional shape representative of at least a portion of human OP-1; and

7 (c) producing the candidate analog identified in step (b).

1 11. The method of claim 10, further comprising the step of determining whether the
2 compound produced in step (c) has an OP-1-like biological activity.

1 12. The method of claim 10, wherein the molecular model provided in step (a) is
2 representative of at least a portion of a finger 1 region of human OP-1.

1 13. The method of claim 10 or 12, wherein the molecular model provided in step (a) is
2 representative of at least a portion of a heel region of human OP-1.

1 14. The method of claim 10 or 12, wherein the model provided in step (a) is representative of
2 at least a portion of a finger 2 region of human OP-1.

1 15. The method of claim 14, wherein the molecular model provided in step (a) is
2 representative of at least a portion of a heel region of human OP-1.

1 16. The method of claim 10, wherein the analog comprises a plurality of charged moieties
2 spaced about the solvent accessible surface thereof and disposed in a spaced-apart relation
3 corresponding to charged moieties spaced about a portion of the solvent accessible surface of
4 human OP-1.

1 17. The method of claim 10, wherein steps (a) and (b) are performed by means of an
2 electronic processor.

1 18. The method of claim 17, wherein step (a) comprises storing a representation of at least a
2 portion of the atomic co-ordinates of human OP-1 in a computer memory.

1 19. A method of producing a morphogen analog that modulates an osteogenic protein-1 (OP-
2 1) mediated biological effect, the method comprising the steps of:

3 (a) providing in a computer memory atomic X-ray crystallographic co-ordinates defining
4 at least a portion of human OP-1;

5 (b) generating with a processor a molecular model having a three-dimensional shape and
6 a solvent accessible surface representative of at least a portion of human OP-1,

7 (c) identifying a candidate morphogen analog having a three-dimensional structure shape
8 and a solvent accessible surface corresponding to the three-dimensional shape and the solvent
9 accessible surface of at least a portion of human OP-1;

10 (d) producing the candidate morphogen analog identified in step (c); and

11 (e) determining whether the candidate morphogen analog produced in step (d) modulates
12 the OP-1 mediated biological effect.


1 20. The method of claim 11 or 19, further comprising the additional step of producing the
2 compound in a commercially useful quantity.

1 21. The method of claim 11 or 19, wherein said compound is a peptide.

1 22. A compound that modulates an OP-1 mediated biological effect produced by the method
2 of claim 11 or 19.

1 23. The compound of claim 22, wherein said compound agonizes the biological activity of
2 human OP-1.

FIG. 1A



2/98

FIG. 1B




FIG. 1C




FIG. 1D








FIG. 1E*FIG. 1F*

FIG. 2



5/98

Fig. 3



6/98

FIG. 4A




FIG. 4B





Fig. 5A^{7/98}



8/98

Fig. 5B





^{9/98}
Fig. 5C




Fig. 6

	OP-1	BMP-5	BMP-6	OP-2	OP-3	60A	UNIVIN	BMP-2	BMP-4	dpp	Vg-1	GDF-6	GDF-7	CDMP-2	GDF-5	BMP-9	DORSALIN	BMP-10	GDF-3	GDF-1	SCREW	GDF-10	BMP-3	NODAL	Inhibin β A	Inhibin β B	Inhibin β C	TGF- β 4	TGF- β 5	TGF- β 1	TGF- β 3
+ OP-1	100																														
+ BMP-5	88	100																													
+ BMP-6	87	91	100																												
+ OP-2	74	74	75	100																											
+ OP-3	67	67	68	77	100																										
+ 60A	69	74	71	63	55	100																									
+ UNIVIN	63	62	63	61	58	54	100																								
+ BMP-2	60	61	61	55	57	57	67	100																							
+ BMP-4	58	59	60	55	56	54	64	92	100																						
+ dpp	58	57	59	53	51	54	56	74	76	100																					
+ Vg-1	57	56	58	55	56	51	64	58	56	48	100																				
+ GDF-6	53	54	53	52	49	48	60	57	56	53	50	100																			
+ GDF-7	53	52	52	53	51	48	60	57	57	53	49	86	100																		
+ CDMP-2	53	54	53	51	48	47	57	54	55	52	51	96	82	100																	
+ GDF-5	51	52	51	50	48	48	58	57	57	52	52	86	80	82	100																
+ BMP-9	51	53	53	48	44	47	50	50	50	51	48	52	50	53	50	100															
+ DORSALIN	49	51	53	47	44	48	47	53	54	53	46	55	56	54	53	79	100														
+ BMP-10	47	48	48	46	45	48	49	55	52	49	50	57	55	56	49	63	66	100													
+ GDF-3	49	49	52	54	52	46	56	53	51	46	56	47	43	45	47	45	43	44	100												
+ GDF-1	45	44	45	44	44	39	49	40	41	37	56	44	47	43	47	40	41	39	51	100											
+ SCREW	47	48	48	47	43	52	42	43	44	39	42	43	42	42	39	41	45	43	42	41	100										
+ GDF-10	42	43	44	40	40	40	41	46	45	43	48	42	42	40	43	40	37	37	38	40	31	100									
+ BMP-3	42	43	44	41	41	41	48	48	47	43	49	46	46	44	47	38	39	42	42	42	33	83	100								
+ NODAL	41	40	42	45	38	40	49	42	41	41	42	44	45	42	42	42	41	41	40	37	35	39	41	100							
+ Inhibin β A	43	43	44	41	39	38	41	43	42	38	42	40	38	40	38	39	36	38	42	31	29	34	35	37	100						
+ Inhibin β B	38	38	38	37	36	36	42	41	41	37	38	37	36	36	35	34	37	41	34	27	35	38	36	63	100						
+ Inhibin β C	39	38	38	35	37	35	37	38	37	34	34	38	39	36	35	34	32	35	39	30	31	33	37	24	53	100					
+ TGF- β 4	38	36	39	34	36	37	34	35	34	36	35	35	37	34	32	28	30	34	31	26	29	29	29	33	32	37	100				
+ TGF- β 5	37	37	38	34	34	37	33	35	34	36	33	38	35	37	35	31	32	35	33	28	26	29	29	30	36	30	33	79	100		
+ TGF- β 1	35	35	36	31	33	38	31	35	34	35	33	33	34	32	31	29	30	35	32	29	25	28	29	30	37	32	34	86	82	100	
+ TGF- β 3	37	36	38	36	37	40	33	36	35	36	37	35	35	34	34	30	31	36	32	33	31	30	32	31	36	33	36	74	73	78	100
+ TGF- β 2	36	34	37	34	33	40	32	35	34	36	37	35	35	34	34	30	33	37	30	29	29	31	33	34	38	31	33	68	70	74	82

86/10



11/98



12/98

Fig. 8.1

Residue	Monomer % Area	Dimer % Area	Hidden Epitope % Area Residues	Surface Modifiable	Modifiable to Improve Solubility
36 GLN	71.89	71.89	0.00		
37 ALA	52.51	52.51	0.00		
39 LYS	62.19	62.19	0.00 EPITOPE		
40 LYS	39.26	39.26	0.00 EPITOPE		
41 HIS	27.13	27.13	0.00	*	
42 GLU	79.09	79.09	0.00 EPITOPE		
43 LEU	51.26	11.83	-39.43		
44 TYR	50.51	50.51	0.00 EPITOPE		
45 VAL	15.22	0.51	-14.71		
46 SER	23.02	23.02	0.00	*	
47 PHE	3.26	3.26	0.00		
48 ARG	76.89	76.89	0.00 EPITOPE		
49 ASP	68.71	52.15	-16.56 EPITOPE		
50 LEU	37.77	0.00	-37.77		
51 GLY	0.00	0.00	0.00		
52 TRP	40.99	34.53	-6.46	*	*
53 GLN	54.47	54.47	0.00 EPITOPE		
54 ASP	54.22	54.22	0.00 EPITOPE		
55 TRP	62.99	62.99	0.00 EPITOPE		
56 ILE	9.68	9.68	0.00		y
57 ILE	33.58	33.58	0.00 EPITOPE		
58 ALA	0.00	0.00	0.00		
59 PRO	34.01	34.01	0.00 EPITOPE		
60 GLU	60.90	60.90	0.00 EPITOPE		
61 GLY	0.00	0.00	0.00		
62 TYR	8.93	2.09	-6.84		
63 ALA	39.31	39.31	0.00	*	*

13/98
Fig. 8.2

64	ALA	14.78	0.00	-14.78		
65	TYR	26.22	26.22	0.00	*	*
66	TYR	48.32	15.41	-32.91		
67	CYS	1.67	1.67	0.00		
68	GLU	59.70	43.27	-16.43	EPITOPE	
69	GLY	0.00	0.00	0.00		
70	GLU	35.82	35.82	0.00	EPITOPE	
71	CYS	0.00	0.00	0.00		
72	ALA	43.27	43.27	0.00	*	*
73	PHE	39.54	39.54	0.00	EPITOPE	
74	PRO	96.68	96.68	0.00	EPITOPE	
75	LEU	1.72	1.72	0.00		
76	ASN	60.54	60.54	0.00	EPITOPE	
77	SER	73.24	73.24	0.00	EPITOPE	
78	TYR	104.34	104.34	0.00	EPITOPE	
79	MET	12.40	12.40	0.00		
80	ASN	46.31	46.31	0.00	*	
81	ALA	32.45	32.45	0.00	*	*
82	THR	34.63	5.99	-28.64		
83	ASN	84.54	38.00	-46.54		
84	HIS	71.01	0.26	-70.75		
85	ALA	0.00	0.00	0.00		
86	ILE	46.99	46.93	-0.06	*	*
87	VAL	64.29	1.95	-62.34		
88	GLN	18.05	4.31	-13.74		
89	THR	4.29	4.29	0.00		
90	LEU	50.95	29.43	-21.52		
91	VAL	39.39	8.51	-30.88		
92	HIS	26.42	26.42	0.00	*	

14/98
Fig. 8.3

93	PHE	73.77	73.77	0.00	EPITOPE
94	ILE	57.23	32.03	-25.20	EPITOPE
95	ASN	43.23	43.23	0.00	EPITOPE
96	PRO	66.64	66.64	0.00	EPITOPE
97	GLU	88.25	88.25	0.00	EPITOPE
98	THR	52.59	48.71	-3.88	EPITOPE
99	VAL	25.83	0.00	-25.83	
100	PRO	89.22	30.78	-58.44	
101	LYS	35.15	35.15	0.00	*
102	PRO	0.00	0.00	0.00	
103	CYS	79.14	27.13	-52.01	
104	CYS	5.39	5.39	0.00	
105	ALA	44.46	5.15	-39.31	
106	PRO	11.24	2.30	-8.94	
107	THR	21.76	21.76	0.00	*
108	GLN	53.40	53.40	0.00	EPITOPE
109	LEU	29.98	7.79	-22.19	
110	ASN	35.00	35.00	0.00	*
111	ALA	23.61	23.61	0.00	* *
112	ILE	22.72	22.72	0.00	* *
113	SER	38.55	38.55	0.00	*
114	VAL	1.15	1.15	0.00	
115	LEU	36.05	36.05	0.00	EPITOPE
116	TYR	18.62	18.62	0.00	
117	PHE	46.55	46.55	0.00	EPITOPE
118	ASP	32.53	32.53	0.00	EPITOPE
119	ASP	84.02	84.02	0.00	EPITOPE
120	SER	48.35	48.35	0.00	EPITOPE
121	SER	68.39	68.39	0.00	EPITOPE

15/98


Fig. 8.4

122	ASN	63.15	63.15	0.00	EPITOPE
123	VAL	41.27	41.27	0.00	EPITOPE
124	ILE	34.51	34.51	0.00	EPITOPE
125	LEU	63.34	63.34	0.00	EPITOPE
126	LYS	54.81	54.81	0.00	EPITOPE
127	LYS	48.78	48.78	0.00	EPITOPE
128	TYR	34.23	32.55	-1.68	
129	ARG	63.25	62.85	-0.40	EPITOPE
130	ASN	62.31	40.62	-21.69	
131	MET	32.35	7.44	-24.91	
132	VAL	16.38	16.38	0.00	
133	VAL	7.50	0.07	-7.43	
134	ARG	65.10	65.10	0.00	
135	ALA	47.10	47.10	0.00	
136	CYS	0.29	0.29	0.00	
137	GLY	0.00	0.00	0.00	
138	CYS	0.00	0.00	0.00	
139	HIS	47.68	18.94	-28.74	

16/98

Fig. 9

Ridge Residues		Receptor Sites	
B90	Leu	Heel	
B91	Val	Heel	
B92	His	Heel	
B93	Phe	Heel	*
B94	Ile	Heel	
B95	Asn	Heel	*
B96	Pro	Heel	*
B97	Glu	Heel	*
B98	Thr	Heel	*
A48	Arg	Finger 1	*
A49	Asp	Finger 1	
A50	Leu	Finger 1	
A51	Gly	Finger 1	
A52	Trp	Finger 1	
A53	Gln	Finger 1	*
A54	Asp	Finger 1	
A55	Trp	Finger 1	
A56	Ile	Finger 1	
A57	Ile	Finger 1	
A58	Ala	Finger 1	
A59	Pro	Finger 1	
A60	Glu	Finger 1	*
A116	Tyr	Finger 2	
A117	Phe	Finger 2	*
A118	Asp	Finger 2	
A119	Asp	Finger 2	*
A120	Ser	Finger 2	*
A121	Ser	Finger 2	*
A122	Asn	Finger 2	*
A123	Val	Finger 2	
A124	Ile	Finger 2	
A125	Leu	Finger 2	
A126	Lys	Finger 2	
A127	Lys	Finger 2	*
A128	Tyr	Finger 2	
A129	Arg	Finger 2	*

^{17/98}
Fig. 10

18/98

Fig. 11A

Residue in 1st Chain	Residue in 2nd Chain	Distance (Å)
Ala-105	Ala-105	3.61
Cys-103	Cys-103	3.95
Asn-83	Asn-130	4.01
Thr-82	Asn-130	4.20

Fig. 11B

Residue in Finger 1	Residue in Finger 2	Distance (Å)
Ala-58	Val-114	3.30
Tyr-65	Val-133	3.93
Ala-58	Leu-115	3.98
Ile-57	Leu-115	4.62
Ile-56	Tyr-116	4.54
Trp-55	Tyr-116	4.74

FIGURE 12

OP-1: CKKHLYVSFRDLGWQDWIIAPEGYAAYCEGECAPPLNSYMNATNHAIVQTLVHFINPETVPKPCCAPTQLNAISVLVFDSSNVILKKYRNVVRACGCH
 |<-----Finger 1----->| |<-----Heel----->|<-----Finger 2----->|

Finger 1 Peptides:

F1-2: CFRDLGWQDWIIAPC

Heel Peptides:

H-1

H-n2:

H-c2:

CAFP LNSYMNATNHAIVQTLVHFINPETVPKC

CLNSYMNATNHAC

CCFINPETVCC

Finger 2 Peptides



F2-2

F2-3



CYFDDSSNVIC

CYFDDSSNVICKKYRS



20/98

Fig. 13A*Fig. 13B*

21/98

Fig. 13C*Fig. 13D*

22/98

Fig. 14A*Fig. 14B*

23/98

Fig. 15.1

Atom	Residue	Chain	X	Y	Z
1 N	GLN 36	A	3.543	-13.365	11.899
2 CA	GLN 36	A	2.246	-13.227	11.283
3 C	GLN 36	A	2.601	-12.220	10.237
4 O	GLN 36	A	2.736	-11.030	10.439
5 CB	GLN 36	A	1.233	-12.644	12.291
6 CG	GLN 36	A	1.403	-11.130	12.646
7 CD	GLN 36	A	2.684	-10.667	13.351
8 OE1	GLN 36	A	3.651	-11.405	13.452
9 NE2	GLN 36	A	2.743	-9.471	13.880
10 N	ALA 37	A	2.753	-12.854	9.097
11 CA	ALA 37	A	3.663	-12.466	8.044
12 C	ALA 37	A	2.836	-11.639	7.174
13 O	ALA 37	A	1.660	-11.899	7.227
14 CB	ALA 37	A	4.126	-13.708	7.294
15 N	CYS 38	A	3.205	-10.694	6.369
16 CA	CYS 38	A	2.177	-9.971	5.682
17 C	CYS 38	A	1.538	-10.860	4.675
18 O	CYS 38	A	2.249	-11.266	3.753
19 CB	CYS 38	A	2.821	-8.791	5.057
20 SG	CYS 38	A	1.945	-8.415	3.597
21 N	LYS 39	A	0.235	-11.141	4.862
22 CA	LYS 39	A	-0.586	-11.883	3.903
23 C	LYS 39	A	-1.990	-11.277	3.660
24 O	LYS 39	A	-2.443	-10.382	4.358
25 CB	LYS 39	A	-0.753	-13.348	4.365
26 CG	LYS 39	A	-0.858	-13.470	5.883
27 CD	LYS 39	A	-1.201	-14.824	6.505
28 CE	LYS 39	A	-0.991	-14.585	8.025
29 NZ	LYS 39	A	-1.199	-15.799	8.829
30 N	LYS 40	A	-2.728	-11.748	2.658
31 CA	LYS 40	A	-4.076	-11.299	2.397
32 C	LYS 40	A	-4.987	-11.916	3.394
33 O	LYS 40	A	-4.652	-12.991	3.851
34 CB	LYS 40	A	-4.574	-11.743	1.058
35 CG	LYS 40	A	-5.826	-11.024	0.591
36 CD	LYS 40	A	-6.625	-11.965	-0.285
37 CE	LYS 40	A	-5.763	-12.585	-1.352
38 NZ	LYS 40	A	-6.552	-13.536	-2.104
39 N	HIS 41	A	-6.124	-11.330	3.782
40 CA	HIS 41	A	-7.047	-11.957	4.717
41 C	HIS 41	A	-8.465	-11.660	4.308
42 O	HIS 41	A	-8.676	-10.901	3.384
43 CB	HIS 41	A	-6.945	-11.406	6.101
44 CG	HIS 41	A	-5.626	-11.606	6.758
45 ND1	HIS 41	A	-5.353	-12.221	7.907

24/98

Fig. 15.2

Atom	Residue	Chain	X	Y	Z	
46	CD2	HIS 41	A	-4.544	-10.869	6.408
47	CE1	HIS 41	A	-4.173	-11.833	8.293
48	NE2	HIS 41	A	-3.705	-11.017	7.380
49	N	GLU 42	A	-9.516	-12.183	4.920
50	CA	GLU 42	A	-10.858	-12.077	4.368
51	C	GLU 42	A	-11.642	-10.923	4.856
52	O	GLU 42	A	-11.379	-10.417	5.925
53	CB	GLU 42	A	-11.639	-13.320	4.667
54	CG	GLU 42	A	-11.132	-14.387	3.691
55	CD	GLU 42	A	-12.251	-15.373	3.342
56	OE1	GLU 42	A	-11.923	-16.555	3.062
57	OE2	GLU 42	A	-13.443	-14.955	3.347
58	N	LEU 43	A	-12.631	-10.414	4.163
59	CA	LEU 43	A	-13.426	-9.333	4.751
60	C	LEU 43	A	-14.554	-9.413	3.765
61	O	LEU 43	A	-14.352	-9.107	2.600
62	CB	LEU 43	A	-12.846	-7.946	4.589
63	CG	LEU 43	A	-12.886	-6.923	5.690
64	CD1	LEU 43	A	-13.082	-5.561	5.031
65	CD2	LEU 43	A	-13.968	-7.216	6.695
66	N	TYR 44	A	-15.723	-9.818	4.196
67	CA	TYR 44	A	-16.879	-9.800	3.365
68	C	TYR 44	A	-17.493	-8.484	3.638
69	O	TYR 44	A	-17.838	-8.320	4.790
70	CB	TYR 44	A	-17.749	-10.903	3.800
71	CG	TYR 44	A	-18.836	-11.049	2.820
72	CD1	TYR 44	A	-20.055	-10.566	3.189
73	CD2	TYR 44	A	-18.610	-11.580	1.592
74	CE1	TYR 44	A	-21.079	-10.473	2.308
75	CE2	TYR 44	A	-19.633	-11.489	0.699
76	CZ	TYR 44	A	-20.825	-10.877	1.042
77	OH	TYR 44	A	-21.774	-10.553	0.068
78	N	VAL 45	A	-17.673	-7.514	2.740
79	CA	VAL 45	A	-18.427	-6.338	3.122
80	C	VAL 45	A	-19.847	-6.545	2.610
81	O	VAL 45	A	-20.040	-6.909	1.461
82	CB	VAL 45	A	-17.796	-5.139	2.507
83	CG1	VAL 45	A	-18.457	-3.857	2.960
84	CG2	VAL 45	A	-16.348	-5.147	2.939
85	N	SER 46	A	-20.877	-6.334	3.431
86	CA	SER 46	A	-22.255	-6.548	3.059
87	C	SER 46	A	-22.799	-5.210	2.818
88	O	SER 46	A	-22.704	-4.398	3.707
89	CB	SER 46	A	-23.054	-7.160	4.179
90	OG	SER 46	A	-24.417	-7.335	3.799

25/98

Fig. 15.3

Atom	Residuc	Chain	X	Y	Z	
91	N	PHE 47	A	-23.389	-4.851	1.705
92	CA	PHE 47	A	-23.751	-3.481	1.472
93	C	PHE 47	A	-24.740	-2.959	2.450
94	O	PHE 47	A	-24.979	-1.770	2.498
95	CB	PHE 47	A	-24.352	-3.321	0.142
96	CG	PHE 47	A	-23.337	-3.619	-0.910
97	CD1	PHE 47	A	-22.233	-2.845	-1.022
98	CD2	PHE 47	A	-23.634	-4.545	-1.868
99	CE1	PHE 47	A	-21.486	-2.920	-2.152
100	CE2	PHE 47	A	-22.889	-4.598	-3.018
101	CZ	PHE 47	A	-21.825	-3.765	-3.173
102	N	ARG 48	A	-25.371	-3.781	3.275
103	CA	ARG 48	A	-26.102	-3.279	4.414
104	C	ARG 48	A	-25.162	-2.527	5.278
105	O	ARG 48	A	-25.411	-1.399	5.572
106	CB	ARG 48	A	-26.719	-4.406	5.231
107	CG	ARG 48	A	-27.809	-5.147	4.442
108	CD	ARG 48	A	-28.661	-6.042	5.341
109	NE	ARG 48	A	-29.918	-6.386	4.698
110	CZ	ARG 48	A	-30.415	-7.627	4.795
111	NH1	ARG 48	A	-29.715	-8.597	5.456
112	NH2	ARG 48	A	-31.623	-7.961	4.241
113	N	ASP 49	A	-24.037	-3.036	5.731
114	CA	ASP 49	A	-23.280	-2.378	6.778
115	C	ASP 49	A	-22.753	-1.006	6.372
116	O	ASP 49	A	-22.313	-0.188	7.168
117	CB	ASP 49	A	-22.095	-3.224	7.190
118	CG	ASP 49	A	-22.486	-4.646	7.506
119	OD1	ASP 49	A	-21.605	-5.482	7.303
120	OD2	ASP 49	A	-23.619	-4.962	7.924
121	N	LEU 50	A	-22.770	-0.677	5.092
122	CA	LEU 50	A	-22.319	0.638	4.686
123	C	LEU 50	A	-23.510	1.477	4.402
124	O	LEU 50	A	-23.448	2.588	3.870
125	CB	LEU 50	A	-21.468	0.539	3.433
126	CG	LEU 50	A	-20.287	-0.404	3.582
127	CD1	LEU 50	A	-20.061	-1.030	2.227
128	CD2	LEU 50	A	-19.052	0.319	4.121
129	N	GLY 51	A	-24.651	0.915	4.749
130	CA	GLY 51	A	-25.907	1.551	4.454
131	C	GLY 51	A	-26.245	1.497	2.985
132	O	GLY 51	A	-27.355	1.857	2.609
133	N	TRP 52	A	-25.389	1.072	2.054
134	CA	TRP 52	A	-25.828	1.011	0.660
135	C	TRP 52	A	-27.045	0.114	0.381

26/98
Fig. 15.4

Atom		Residue		Chain	X	Y	Z
136	O	TRP	52	A	-27.129	-0.483	-0.668
137	CB	TRP	52	A	-24.614	0.556	-0.189
138	CG	TRP	52	A	-23.484	1.602	-0.152
139	CD1	TRP	52	A	-23.615	2.896	0.315
140	CD2	TRP	52	A	-22.092	1.295	-0.610
141	NE1	TRP	52	A	-22.430	3.454	0.213
142	CE2	TRP	52	A	-21.489	2.641	-0.293
143	CE3	TRP	52	A	-21.262	0.304	-1.105
144	CZ2	TRP	52	A	-20.133	2.852	-0.442
145	CZ3	TRP	52	A	-19.913	0.575	-1.247
146	CH2	TRP	52	A	-19.360	1.810	-0.907
147	N	GLN	53	A	-28.072	-0.088	1.197
148	CA	GLN	53	A	-28.758	-1.377	1.098
149	C	GLN	53	A	-29.798	-1.458	0.072
150	O	GLN	53	A	-30.460	-2.475	-0.062
151	CB	GLN	53	A	-29.504	-1.835	2.360
152	CG	GLN	53	A	-29.756	-0.759	3.403
153	CD	GLN	53	A	-31.038	-1.078	4.192
154	OE1	GLN	53	A	-31.953	-0.232	4.230
155	NE2	GLN	53	A	-31.163	-2.269	4.840
156	N	ASP	54	A	-29.992	-0.405	-0.689
157	CA	ASP	54	A	-31.357	0.003	-0.992
158	C	ASP	54	A	-31.136	0.729	-2.255
159	O	ASP	54	A	-31.000	1.942	-2.119
160	CB	ASP	54	A	-31.873	0.982	0.073
161	CG	ASP	54	A	-30.722	1.807	0.710
162	OD1	ASP	54	A	-30.861	2.211	1.892
163	OD2	ASP	54	A	-29.698	2.012	0.027
164	N	TRP	55	A	-31.067	0.175	-3.470
165	CA	TRP	55	A	-30.322	0.915	-4.515
166	C	TRP	55	A	-29.770	-0.083	-5.462
167	O	TRP	55	A	-29.915	0.104	-6.662
168	CB	TRP	55	A	-29.011	1.599	-4.158
169	CG	TRP	55	A	-29.091	3.035	-3.787
170	CD1	TRP	55	A	-30.088	3.925	-4.101
171	CD2	TRP	55	A	-27.886	3.623	-3.134
172	NE1	TRP	55	A	-29.608	5.108	-3.722
173	CE2	TRP	55	A	-28.348	5.080	-3.173
174	CE3	TRP	55	A	-26.706	3.288	-2.489
175	CZ2	TRP	55	A	-27.508	6.047	-2.602
176	CZ3	TRP	55	A	-25.932	4.297	-1.921
177	CH2	TRP	55	A	-26.304	5.642	-1.988
178	N	ILE	56	A	-29.151	-1.108	-4.812
179	CA	ILE	56	A	-28.323	-2.142	-5.400
180	C	ILE	56	A	-29.209	-3.352	-5.699

27/98

Fig. 15.5

Atom	Residuc	Chain	X	Y	Z	
181	O	ILE 56	A	-29.900	-3.795	-4.792
182	CB	ILE 56	A	-27.221	-2.452	-4.390
183	CG1	ILE 56	A	-25.998	-1.743	-4.815
184	CG2	ILE 56	A	-26.917	-3.924	-4.291
185	CD1	ILE 56	A	-25.964	-0.422	-4.122
186	N	ILE 57	A	-29.260	-3.924	-6.907
187	CA	ILE 57	A	-29.961	-5.182	-7.093
188	C	ILE 57	A	-28.927	-6.288	-6.809
189	O	ILE 57	A	-29.163	-7.475	-6.713
190	CB	ILE 57	A	-30.481	-5.241	-8.549
191	CG1	ILE 57	A	-31.405	-4.079	-8.780
192	CG2	ILE 57	A	-31.273	-6.511	-8.844
193	CD1	ILE 57	A	-31.745	-4.110	-10.274
194	N	ALA 58	A	-27.684	-5.914	-6.623
195	CA	ALA 58	A	-26.550	-6.030	-7.556
196	C	ALA 58	A	-25.789	-7.248	-7.200
197	O	ALA 58	A	-26.272	-8.230	-7.747
198	CB	ALA 58	A	-25.682	-4.813	-7.386
199	N	PRO 59	A	-24.748	-7.272	-6.411
200	CA	PRO 59	A	-24.570	-8.349	-5.438
201	C	PRO 59	A	-25.086	-7.952	-4.063
202	O	PRO 59	A	-25.265	-6.789	-3.797
203	CB	PRO 59	A	-23.068	-8.697	-5.439
204	CG	PRO 59	A	-22.447	-7.366	-5.730
205	CD	PRO 59	A	-23.408	-6.817	-6.757
206	N	GLU 60	A	-25.376	-8.795	-3.088
207	CA	GLU 60	A	-25.696	-8.242	-1.777
208	C	GLU 60	A	-24.492	-7.832	-0.980
209	O	GLU 60	A	-24.571	-7.233	0.078
210	CB	GLU 60	A	-26.502	-9.249	-0.927
211	CG	GLU 60	A	-25.753	-10.380	-0.180
212	CD	GLU 60	A	-24.990	-9.971	1.084
213	OE1	GLU 60	A	-25.319	-8.939	1.717
214	OE2	GLU 60	A	-24.089	-10.738	1.457
215	N	GLY 61	A	-23.307	-8.146	-1.448
216	CA	GLY 61	A	-22.099	-7.722	-0.784
217	C	GLY 61	A	-21.009	-8.349	-1.571
218	O	GLY 61	A	-21.289	-8.862	-2.648
219	N	TYR 62	A	-19.777	-8.343	-1.088
220	CA	TYR 62	A	-18.689	-8.872	-1.879
221	C	TYR 62	A	-17.524	-9.144	-0.981
222	O	TYR 62	A	-17.554	-8.747	0.169
223	CB	TYR 62	A	-18.362	-7.838	-2.934
224	CG	TYR 62	A	-17.588	-6.667	-2.413
225	CD1	TYR 62	A	-18.181	-5.490	-1.956

28/98

Fig. 15.6

Atom	Residue	Chain	X	Y	Z	
226	CD2	TYR 62	A	-16.236	-6.785	-2.513
227	CE1	TYR 62	A	-17.379	-4.412	-1.599
228	CE2	TYR 62	A	-15.447	-5.739	-2.169
229	CZ	TYR 62	A	-16.004	-4.571	-1.715
230	OH	TYR 62	A	-15.097	-3.565	-1.383
231	N	ALA 63	A	-16.476	-9.808	-1.456
232	CA	ALA 63	A	-15.231	-9.961	-0.740
233	C	ALA 63	A	-14.242	-8.820	-0.961
234	O	ALA 63	A	-13.659	-8.712	-2.023
235	CB	ALA 63	A	-14.655	-11.248	-1.205
236	N	ALA 64	A	-14.020	-7.948	0.007
237	CA	ALA 64	A	-13.051	-6.893	-0.114
238	C	ALA 64	A	-11.606	-7.304	0.192
239	O	ALA 64	A	-10.608	-6.800	-0.298
240	CB	ALA 64	A	-13.454	-5.781	0.832
241	N	TYR 65	A	-11.469	-8.294	1.060
242	CA	TYR 65	A	-10.215	-8.675	1.672
243	C	TYR 65	A	-9.503	-7.544	2.281
244	O	TYR 65	A	-9.863	-6.400	2.147
245	CB	TYR 65	A	-9.290	-9.333	0.678
246	CG	TYR 65	A	-9.846	-10.614	0.126
247	CD1	TYR 65	A	-10.300	-11.608	0.939
248	CD2	TYR 65	A	-9.927	-10.753	-1.219
249	CE1	TYR 65	A	-10.905	-12.716	0.405
250	CE2	TYR 65	A	-10.526	-11.847	-1.767
251	CZ	TYR 65	A	-11.039	-12.799	-0.951
252	OH	TYR 65	A	-11.729	-13.859	-1.515
253	N	TYR 66	A	-8.459	-7.830	2.998
254	CA	TYR 66	A	-7.615	-6.771	3.442
255	C	TYR 66	A	-6.303	-7.401	3.710
256	O	TYR 66	A	-6.198	-8.619	3.836
257	CB	TYR 66	A	-8.154	-6.109	4.712
258	CG	TYR 66	A	-8.091	-6.906	5.984
259	CD1	TYR 66	A	-8.948	-7.941	6.159
260	CD2	TYR 66	A	-7.215	-6.546	6.978
261	CE1	TYR 66	A	-8.929	-8.634	7.333
262	CE2	TYR 66	A	-7.183	-7.227	8.155
263	CZ	TYR 66	A	-8.041	-8.273	8.310
264	OH	TYR 66	A	-8.024	-9.037	9.472
265	N	CYS 67	A	-5.286	-6.571	3.812
266	CA	CYS 67	A	-3.921	-7.055	3.949
267	C	CYS 67	A	-3.364	-6.825	5.358
268	O	CYS 67	A	-3.426	-5.705	5.888
269	CB	CYS 67	A	-2.971	-6.328	3.056
270	SG	CYS 67	A	-3.158	-6.344	1.310

29/98

Fig. 15.7

Atom	Residue	Chain	X	Y	Z	
271	N	GLU 68	A	-2.792	-7.828	6.031
272	CA	GLU 68	A	-2.322	-7.457	7.318
273	C	GLU 68	A	-1.275	-8.383	7.726
274	O	GLU 68	A	-1.438	-9.577	7.481
275	CB	GLU 68	A	-3.408	-7.534	8.334
276	CG	GLU 68	A	-3.092	-6.620	9.488
277	CD	GLU 68	A	-3.797	-7.134	10.693
278	OE1	GLU 68	A	-3.899	-6.397	11.674
279	OE2	GLU 68	A	-4.246	-8.282	10.661
280	N	GLY 69	A	-0.219	-7.857	8.348
281	CA	GLY 69	A	0.772	-8.701	8.979
282	C	GLY 69	A	2.107	-7.997	8.920
283	O	GLY 69	A	2.212	-6.844	8.501
284	N	GLU 70	A	3.192	-8.624	9.316
285	CA	GLU 70	A	4.349	-7.810	9.600
286	C	GLU 70	A	5.364	-7.795	8.496
287	O	GLU 70	A	5.603	-8.818	7.881
288	CB	GLU 70	A	4.996	-8.309	10.865
289	CG	GLU 70	A	5.202	-7.080	11.735
290	CD	GLU 70	A	6.392	-7.339	12.618
291	OE1	GLU 70	A	7.092	-6.368	12.941
292	OE2	GLU 70	A	6.637	-8.516	12.968
293	N	CYS 71	A	5.988	-6.666	8.222
294	CA	CYS 71	A	7.049	-6.657	7.253
295	C	CYS 71	A	8.436	-6.637	7.832
296	O	CYS 71	A	8.961	-5.627	8.241
297	CB	CYS 71	A	6.787	-5.467	6.371
298	SG	CYS 71	A	5.598	-5.970	5.102
299	N	ALA 72	A	9.141	-7.740	7.914
300	CA	ALA 72	A	10.322	-7.789	8.729
301	C	ALA 72	A	11.166	-8.899	8.174
302	O	ALA 72	A	10.670	-9.710	7.406
303	CB	ALA 72	A	9.874	-8.066	10.139
304	N	PHE 73	A	12.457	-9.057	8.463
305	CA	PHE 73	A	13.195	-10.119	7.773
306	C	PHE 73	A	12.738	-11.435	8.328
307	O	PHE 73	A	12.521	-11.454	9.530
308	CB	PHE 73	A	14.723	-9.995	7.990
309	CG	PHE 73	A	15.255	-8.809	7.230
310	CD1	PHE 73	A	15.243	-8.804	5.868
311	CD2	PHE 73	A	15.746	-7.737	7.919
312	CE1	PHE 73	A	15.709	-7.716	5.190
313	CE2	PHE 73	A	16.225	-6.647	7.217
314	CZ	PHE 73	A	16.203	-6.634	5.857
315	N	PRO 74	A	12.570	-12.531	7.695

30/98

Fig. 15.8

Atom	Residue	Chain	X	Y	Z	
316	CA	PRO 74	A	12.982	-12.802	6.339
317	C	PRO 74	A	11.990	-12.148	5.485
318	O	PRO 74	A	10.891	-12.643	5.423
319	CB	PRO 74	A	12.945	-14.284	6.176
320	CG	PRO 74	A	12.664	-14.772	7.587
321	CD	PRO 74	A	11.893	-13.668	8.292
322	N	LEU 75	A	12.254	-11.051	4.786
323	CA	LEU 75	A	11.328	-10.466	3.831
324	C	LEU 75	A	11.299	-11.508	2.762
325	O	LEU 75	A	11.992	-11.532	1.756
326	CB	LEU 75	A	11.912	-9.144	3.346
327	CG	LEU 75	A	10.922	-8.023	3.399
328	CD1	LEU 75	A	9.924	-8.232	4.514
329	CD2	LEU 75	A	11.680	-6.741	3.622
330	N	ASN 76	A	10.419	-12.442	3.038
331	CA	ASN 76	A	10.337	-13.665	2.288
332	C	ASN 76	A	9.758	-13.535	0.863
333	O	ASN 76	A	8.766	-14.166	0.569
334	CB	ASN 76	A	9.548	-14.511	3.229
335	CG	ASN 76	A	9.704	-15.945	2.935
336	OD1	ASN 76	A	9.686	-16.766	3.862
337	ND2	ASN 76	A	9.760	-16.324	1.643
338	N	SER 77	A	10.263	-12.752	-0.109
339	CA	SER 77	A	9.917	-12.843	-1.545
340	C	SER 77	A	8.445	-12.920	-1.915
341	O	SER 77	A	7.832	-11.897	-2.284
342	CB	SER 77	A	10.599	-14.059	-2.195
343	OG	SER 77	A	10.201	-15.273	-1.560
344	N	TYR 78	A	7.776	-14.081	-1.847
345	CA	TYR 78	A	6.343	-13.977	-1.994
346	C	TYR 78	A	5.734	-13.258	-0.792
347	O	TYR 78	A	4.827	-13.833	-0.216
348	CB	TYR 78	A	5.683	-15.337	-2.079
349	CG	TYR 78	A	5.162	-15.723	-3.465
350	CD1	TYR 78	A	4.144	-16.681	-3.432
351	CD2	TYR 78	A	5.865	-15.473	-4.665
352	CE1	TYR 78	A	3.985	-17.578	-4.480
353	CE2	TYR 78	A	5.706	-16.377	-5.729
354	CZ	TYR 78	A	4.859	-17.489	-5.557
355	OH	TYR 78	A	4.965	-18.610	-6.363
356	N	MET 79	A	6.220	-12.058	-0.418
357	CA	MET 79	A	5.629	-11.035	0.425
358	C	MET 79	A	5.931	-9.703	-0.270
359	O	MET 79	A	5.976	-8.606	0.269
360	CB	MET 79	A	6.271	-10.880	1.754

31/98
Fig. 15.9

Atom	Residue	Chain	X	Y	Z	
361	CG	MET 79	A	6.288	-12.036	2.716
362	SD	MET 79	A	6.974	-11.247	4.193
363	CE	MET 79	A	7.145	-12.664	5.230
364	N	ASN 80	A	6.171	-9.795	-1.562
365	CA	ASN 80	A	6.444	-8.675	-2.417
366	C	ASN 80	A	7.324	-7.534	-1.973
367	O	ASN 80	A	7.169	-6.458	-2.499
368	CB	ASN 80	A	5.129	-8.059	-2.909
369	CG	ASN 80	A	5.403	-7.179	-4.135
370	OD1	ASN 80	A	5.383	-5.934	-4.172
371	ND2	ASN 80	A	5.792	-7.856	-5.212
372	N	ALA 81	A	8.295	-7.505	-1.068
373	CA	ALA 81	A	9.543	-6.854	-1.431
374	C	ALA 81	A	9.731	-5.440	-1.931
375	O	ALA 81	A	10.248	-4.546	-1.301
376	CB	ALA 81	A	10.249	-7.718	-2.461
377	N	THR 82	A	9.352	-5.062	-3.108
378	CA	THR 82	A	10.122	-4.047	-3.818
379	C	THR 82	A	11.649	-4.060	-3.652
380	O	THR 82	A	12.323	-4.730	-4.441
381	CB	THR 82	A	9.752	-2.568	-3.534
382	OG1	THR 82	A	9.779	-2.312	-2.158
383	CG2	THR 82	A	8.458	-2.245	-4.220
384	N	ASN 83	A	12.288	-3.380	-2.692
385	CA	ASN 83	A	13.708	-3.070	-2.868
386	C	ASN 83	A	13.915	-1.957	-1.951
387	O	ASN 83	A	14.715	-1.956	-1.057
388	CB	ASN 83	A	14.082	-2.493	-4.184
389	CG	ASN 83	A	14.958	-3.390	-4.974
390	OD1	ASN 83	A	15.593	-2.957	-5.919
391	ND2	ASN 83	A	14.990	-4.682	-4.709
392	N	HIS 84	A	13.196	-0.870	-2.081
393	CA	HIS 84	A	13.067	0.094	-1.008
394	C	HIS 84	A	12.526	-0.632	0.181
395	O	HIS 84	A	12.864	-0.201	1.261
396	CB	HIS 84	A	12.095	1.187	-1.379
397	CG	HIS 84	A	12.020	2.264	-0.350
398	ND1	HIS 84	A	11.255	2.299	0.730
399	CD2	HIS 84	A	12.640	3.478	-0.465
400	CE1	HIS 84	A	11.393	3.487	1.258
401	NE2	HIS 84	A	12.201	4.192	0.533
402	N	ALA 85	A	11.706	-1.690	0.180
403	CA	ALA 85	A	11.398	-2.240	1.463
404	C	ALA 85	A	12.553	-3.020	1.972
405	O	ALA 85	A	12.626	-3.221	3.161

32/98

Fig. 15.10

Atom		Residue		Chain	X	Y	Z
406	CB	ALA	85	A	10.199	-3.130	1.402
407	N	ILE	86	A	13.538	-3.522	1.249
408	CA	ILE	86	A	14.606	-4.201	1.925
409	C	ILE	86	A	15.457	-3.071	2.472
410	O	ILE	86	A	15.835	-3.146	3.602
411	CB	ILE	86	A	15.300	-5.078	0.903
412	CG1	ILE	86	A	14.276	-6.058	0.301
413	CG2	ILE	86	A	16.470	-5.804	1.570
414	CD1	ILE	86	A	14.811	-7.212	-0.547
415	N	VAL	87	A	15.835	-1.962	1.844
416	CA	VAL	87	A	16.554	-0.892	2.511
417	C	VAL	87	A	15.798	-0.536	3.762
418	O	VAL	87	A	16.284	-0.769	4.847
419	CB	VAL	87	A	16.672	0.325	1.574
420	CG1	VAL	87	A	16.890	1.606	2.333
421	CG2	VAL	87	A	17.858	0.133	0.660
422	N	GLN	88	A	14.596	0.026	3.752
423	CA	GLN	88	A	13.942	0.411	4.993
424	C	GLN	88	A	13.900	-0.690	6.037
425	O	GLN	88	A	14.176	-0.400	7.186
426	CB	GLN	88	A	12.531	0.829	4.715
427	CG	GLN	88	A	12.099	2.208	5.146
428	CD	GLN	88	A	10.634	2.123	5.038
429	OE1	GLN	88	A	9.852	2.201	5.962
430	NE2	GLN	88	A	10.159	1.742	3.868
431	N	THR	89	A	13.582	-1.949	5.766
432	CA	THR	89	A	13.633	-2.920	6.799
433	C	THR	89	A	15.038	-3.073	7.309
434	O	THR	89	A	15.197	-3.213	8.509
435	CB	THR	89	A	13.156	-4.198	6.275
436	OG1	THR	89	A	11.863	-3.908	5.883
437	CG2	THR	89	A	12.973	-5.302	7.272
438	N	LEU	90	A	16.122	-3.061	6.519
439	CA	LEU	90	A	17.492	-3.003	7.017
440	C	LEU	90	A	17.782	-1.770	7.832
441	O	LEU	90	A	18.260	-1.850	8.947
442	CB	LEU	90	A	18.454	-3.009	5.868
443	CG	LEU	90	A	19.904	-3.128	6.174
444	CD1	LEU	90	A	20.166	-4.389	6.967
445	CD2	LEU	90	A	20.641	-3.116	4.854
446	N	VAL	91	A	17.528	-0.569	7.354
447	CA	VAL	91	A	17.690	0.623	8.155
448	C	VAL	91	A	16.866	0.546	9.409
449	O	VAL	91	A	17.188	1.150	10.422
450	CB	VAL	91	A	17.282	1.803	7.334

33/98

Fig. 15.11

Atom	Residue	Chain	X	Y	Z	
451	CG1	VAL 91	A	17.149	3.078	8.140
452	CG2	VAL 91	A	18.356	1.954	6.278
453	N	HIS 92	A	15.762	-0.193	9.453
454	CA	HIS 92	A	15.046	-0.309	10.719
455	C	HIS 92	A	15.946	-0.915	11.723
456	O	HIS 92	A	16.085	-0.444	12.816
457	CB	HIS 92	A	13.805	-1.224	10.633
458	CG	HIS 92	A	12.902	-1.039	11.826
459	ND1	HIS 92	A	12.332	-1.848	12.711
460	CD2	HIS 92	A	12.540	0.216	12.159
461	CE1	HIS 92	A	11.678	-1.086	13.538
462	NE2	HIS 92	A	11.826	0.165	13.212
463	N	PHE 93	A	16.591	-1.984	11.374
464	CA	PHE 93	A	17.374	-2.777	12.262
465	C	PHE 93	A	18.620	-2.032	12.652
466	O	PHE 93	A	19.005	-1.893	13.796
467	CB	PHE 93	A	17.592	-4.024	11.472
468	CG	PHE 93	A	18.741	-4.858	11.921
469	CD1	PHE 93	A	20.039	-4.461	11.662
470	CD2	PHE 93	A	18.474	-6.111	12.455
471	CE1	PHE 93	A	21.058	-5.341	11.903
472	CE2	PHE 93	A	19.507	-6.993	12.676
473	CZ	PHE 93	A	20.790	-6.600	12.394
474	N	ILE 94	A	19.399	-1.476	11.751
475	CA	ILE 94	A	20.477	-0.592	12.182
476	C	ILE 94	A	19.938	0.454	13.131
477	O	ILE 94	A	20.706	1.117	13.787
478	CB	ILE 94	A	21.036	-0.005	10.907
479	CG1	ILE 94	A	21.808	-1.156	10.367
480	CG2	ILE 94	A	21.876	1.246	11.008
481	CD1	ILE 94	A	22.103	-0.863	8.903
482	N	ASN 95	A	18.635	0.704	13.302
483	CA	ASN 95	A	18.237	1.955	13.901
484	C	ASN 95	A	16.753	2.185	14.183
485	O	ASN 95	A	16.082	3.112	13.702
486	CB	ASN 95	A	18.719	3.074	13.024
487	CG	ASN 95	A	18.887	4.221	13.977
488	OD1	ASN 95	A	18.243	4.291	15.040
489	ND2	ASN 95	A	19.771	5.172	13.602
490	N	PRO 96	A	16.204	1.357	15.006
491	CA	PRO 96	A	14.787	1.108	15.108
492	C	PRO 96	A	13.919	2.281	15.324
493	O	PRO 96	A	12.713	2.226	15.089
494	CB	PRO 96	A	14.647	0.084	16.217
495	CG	PRO 96	A	16.039	-0.490	16.300

34/98
Fig. 15.12

Atom	Residue		Chain	X	Y	Z
496	CD	PRO 96	A	16.846	0.801	16.173
497	N	GLU 97	A	14.559	3.336	15.774
498	CA	GLU 97	A	13.846	4.558	15.992
499	C	GLU 97	A	14.019	5.333	14.722
500	O	GLU 97	A	13.031	5.781	14.197
501	CB	GLU 97	A	14.429	5.329	17.184
502	CG	GLU 97	A	13.550	5.172	18.453
503	CD	GLU 97	A	13.769	3.807	19.163
504	OE1	GLU 97	A	14.723	3.749	19.999
505	OE2	GLU 97	A	12.998	2.824	18.896
506	N	THR 98	A	15.145	5.584	14.080
507	CA	THR 98	A	15.073	6.445	12.897
508	C	THR 98	A	14.148	6.043	11.715
509	O	THR 98	A	14.015	6.906	10.851
510	CB	THR 98	A	16.514	6.624	12.388
511	OG1	THR 98	A	17.067	5.331	12.373
512	CG2	THR 98	A	17.451	7.417	13.322
513	N	VAL 99	A	13.482	4.877	11.518
514	CA	VAL 99	A	12.654	4.635	10.334
515	C	VAL 99	A	11.675	3.508	10.660
516	O	VAL 99	A	12.183	2.611	11.310
517	CB	VAL 99	A	13.546	4.251	9.139
518	CG1	VAL 99	A	13.636	2.753	8.774
519	CG2	VAL 99	A	12.902	4.955	7.978
520	N	PRO 100	A	10.408	3.461	10.322
521	CA	PRO 100	A	9.442	2.427	10.687
522	C	PRO 100	A	9.639	1.132	10.021
523	O	PRO 100	A	10.611	1.033	9.342
524	CB	PRO 100	A	8.119	3.003	10.359
525	CG	PRO 100	A	8.348	4.485	10.464
526	CD	PRO 100	A	9.713	4.645	9.854
527	N	LYS 101	A	8.889	0.042	10.071
528	CA	LYS 101	A	9.245	-1.083	9.226
529	C	LYS 101	A	8.344	-0.893	8.022
530	O	LYS 101	A	7.361	-0.183	8.151
531	CB	LYS 101	A	8.901	-2.426	9.859
532	CG	LYS 101	A	8.737	-2.477	11.364
533	CD	LYS 101	A	8.039	-3.748	11.889
534	CE	LYS 101	A	6.454	-3.645	12.044
535	NZ	LYS 101	A	5.616	-3.935	10.845
536	N	PRO 102	A	8.452	-1.381	6.846
537	CA	PRO 102	A	7.556	-1.032	5.775
538	C	PRO 102	A	6.203	-1.549	6.093
539	O	PRO 102	A	6.087	-2.464	6.892
540	CB	PRO 102	A	8.171	-1.646	4.535

35/98
Fig. 15.13

Atom	Residue	Chain	X	Y	Z	
541	CG	PRO 102	A	8.812	-2.855	5.077
542	CD	PRO 102	A	9.333	-2.458	6.451
543	N	CYS 103	A	5.086	-1.078	5.549
544	CA	CYS 103	A	3.887	-1.780	5.921
545	C	CYS 103	A	3.229	-2.407	4.753
546	O	CYS 103	A	3.592	-2.289	3.598
547	CB	CYS 103	A	2.930	-0.840	6.599
548	SG	CYS 103	A	2.131	-0.050	5.233
549	N	CYS 104	A	2.191	-3.102	5.117
550	CA	CYS 104	A	1.736	-4.199	4.338
551	C	CYS 104	A	0.466	-3.786	3.641
552	O	CYS 104	A	-0.555	-3.520	4.240
553	CB	CYS 104	A	1.633	-5.221	5.372
554	SG	CYS 104	A	0.823	-6.586	4.674
555	N	ALA 105	A	0.490	-3.719	2.337
556	CA	ALA 105	A	-0.396	-2.886	1.605
557	C	ALA 105	A	-0.839	-3.742	0.482
558	O	ALA 105	A	-0.170	-4.745	0.282
559	CB	ALA 105	A	0.400	-1.708	1.149
560	N	PRO 106	A	-1.857	-3.510	-0.271
561	CA	PRO 106	A	-2.156	-4.293	-1.435
562	C	PRO 106	A	-1.352	-3.905	-2.675
563	O	PRO 106	A	-1.065	-2.765	-2.987
564	CB	PRO 106	A	-3.647	-4.136	-1.574
565	CG	PRO 106	A	-3.879	-2.759	-1.095
566	CD	PRO 106	A	-3.107	-2.927	0.186
567	N	THR 107	A	-0.999	-4.955	-3.391
568	CA	THR 107	A	-0.073	-4.974	-4.479
569	C	THR 107	A	-1.001	-4.913	-5.659
570	O	THR 107	A	-0.955	-3.955	-6.424
571	CB	THR 107	A	0.666	-6.265	-4.172
572	OG1	THR 107	A	1.728	-5.736	-3.391
573	CG2	THR 107	A	1.115	-7.153	-5.311
574	N	GLN 108	A	-1.900	-5.847	-5.940
575	CA	GLN 108	A	-2.913	-5.442	-6.865
576	C	GLN 108	A	-4.352	-5.682	-6.500
577	O	GLN 108	A	-4.739	-6.674	-5.905
578	CB	GLN 108	A	-2.460	-6.080	-8.189
579	CG	GLN 108	A	-2.948	-7.440	-8.663
580	CD	GLN 108	A	-2.234	-8.459	-7.886
581	OE1	GLN 108	A	-1.660	-8.222	-6.832
582	NE2	GLN 108	A	-2.298	-9.656	-8.471
583	N	LEU 109	A	-5.100	-4.662	-6.917
584	CA	LEU 109	A	-6.522	-4.430	-6.644
585	C	LEU 109	A	-7.495	-4.780	-7.798

36/98

Fig. 15.14

Atom	Residue	Chain	X	Y	Z	
586	O	LEU 109	A	-7.256	-4.573	-8.986
587	CB	LEU 109	A	-6.694	-2.932	-6.225
588	CG	LEU 109	A	-6.102	-2.671	-4.830
589	CD1	LEU 109	A	-6.012	-1.216	-4.502
590	CD2	LEU 109	A	-7.022	-3.304	-3.811
591	N	ASN 110	A	-8.659	-5.343	-7.521
592	CA	ASN 110	A	-9.570	-5.629	-8.584
593	C	ASN 110	A	-10.824	-4.870	-8.418
594	O	ASN 110	A	-10.985	-4.117	-7.471
595	CB	ASN 110	A	-9.945	-7.054	-8.620
596	CG	ASN 110	A	-8.771	-7.863	-9.017
597	OD1	ASN 110	A	-8.810	-9.080	-8.781
598	ND2	ASN 110	A	-7.687	-7.281	-9.560
599	N	ALA 111	A	-11.767	-5.052	-9.338
600	CA	ALA 111	A	-12.876	-4.131	-9.476
601	C	ALA 111	A	-14.026	-4.974	-9.121
602	O	ALA 111	A	-13.880	-6.178	-9.058
603	CB	ALA 111	A	-12.993	-3.615	-10.928
604	N	ILE 112	A	-15.182	-4.401	-8.876
605	CA	ILE 112	A	-16.371	-5.172	-8.539
606	C	ILE 112	A	-17.371	-4.462	-9.404
607	O	ILE 112	A	-17.265	-3.251	-9.506
608	CB	ILE 112	A	-16.583	-5.062	-6.980
609	CG1	ILE 112	A	-16.477	-6.535	-6.632
610	CG2	ILE 112	A	-17.869	-4.480	-6.379
611	CD1	ILE 112	A	-17.762	-7.334	-6.966
612	N	SER 113	A	-18.320	-5.126	-10.037
613	CA	SER 113	A	-19.389	-4.403	-10.673
614	C	SER 113	A	-20.670	-4.529	-9.920
615	O	SER 113	A	-20.965	-5.655	-9.564
616	CB	SER 113	A	-19.576	-4.919	-12.074
617	OG	SER 113	A	-18.826	-4.052	-12.946
618	N	VAL 114	A	-21.472	-3.484	-9.631
619	CA	VAL 114	A	-22.751	-3.722	-8.970
620	C	VAL 114	A	-23.830	-3.148	-9.882
621	O	VAL 114	A	-23.658	-2.119	-10.522
622	CB	VAL 114	A	-22.897	-3.057	-7.536
623	CG1	VAL 114	A	-21.637	-3.226	-6.742
624	CG2	VAL 114	A	-23.194	-1.588	-7.611
625	N	LEU 115	A	-24.960	-3.847	-9.955
626	CA	LEU 115	A	-26.160	-3.492	-10.709
627	C	LEU 115	A	-27.151	-2.699	-9.903
628	O	LEU 115	A	-27.603	-3.173	-8.860
629	CB	LEU 115	A	-26.840	-4.789	-11.194
630	CG	LEU 115	A	-28.187	-4.790	-11.902

37/98

Fig. 15.15

Atom	Residue	Chain	X	Y	Z	
631	CD1	LEU 115	A	-28.011	-4.403	-13.339
632	CD2	LEU 115	A	-28.797	-6.184	-11.806
633	N	TYR 116	A	-27.598	-1.505	-10.254
634	CA	TYR 116	A	-28.495	-0.859	-9.329
635	C	TYR 116	A	-29.472	0.124	-9.908
636	O	TYR 116	A	-29.241	0.596	-11.014
637	CB	TYR 116	A	-27.619	-0.201	-8.316
638	CG	TYR 116	A	-26.890	0.935	-8.941
639	CD1	TYR 116	A	-25.664	0.712	-9.513
640	CD2	TYR 116	A	-27.474	2.166	-8.888
641	CE1	TYR 116	A	-25.034	1.750	-10.131
642	CE2	TYR 116	A	-26.850	3.205	-9.502
643	CZ	TYR 116	A	-25.669	2.961	-10.141
644	OH	TYR 116	A	-25.140	3.965	-10.902
645	N	PHE 117	A	-30.566	0.506	-9.242
646	CA	PHE 117	A	-31.478	1.445	-9.867
647	C	PHE 117	A	-30.914	2.785	-9.556
648	O	PHE 117	A	-30.775	3.080	-8.381
649	CB	PHE 117	A	-32.882	1.514	-9.284
650	CG	PHE 117	A	-33.447	0.159	-8.982
651	CD1	PHE 117	A	-33.259	-0.395	-7.738
652	CD2	PHE 117	A	-34.137	-0.534	-9.952
653	CE1	PHE 117	A	-33.750	-1.660	-7.482
654	CE2	PHE 117	A	-34.618	-1.807	-9.709
655	CZ	PHE 117	A	-34.419	-2.373	-8.466
656	N	ASP 118	A	-30.555	3.683	-10.458
657	CA	ASP 118	A	-30.375	5.050	-9.964
658	C	ASP 118	A	-31.794	5.566	-9.803
659	O	ASP 118	A	-32.746	4.804	-9.876
660	CB	ASP 118	A	-29.612	5.972	-10.973
661	CG	ASP 118	A	-30.317	5.985	-12.316
662	OD1	ASP 118	A	-29.855	6.724	-13.211
663	OD2	ASP 118	A	-31.322	5.245	-12.467
664	N	ASP 119	A	-31.939	6.868	-9.594
665	CA	ASP 119	A	-33.218	7.509	-9.718
666	C	ASP 119	A	-33.915	7.160	-11.070
667	O	ASP 119	A	-33.359	6.483	-11.946
668	CB	ASP 119	A	-33.033	9.030	-9.618
669	CG	ASP 119	A	-31.942	9.455	-10.598
670	OD1	ASP 119	A	-31.911	8.877	-11.719
671	OD2	ASP 119	A	-31.115	10.325	-10.232
672	N	SER 120	A	-35.161	7.670	-11.218
673	CA	SER 120	A	-36.125	7.119	-12.146
674	C	SER 120	A	-36.295	5.699	-11.621
675	O	SER 120	A	-36.515	5.484	-10.431

38/98

Fig. 15.16

Atom	Residue	Chain	X	Y	Z
676	CB	SER 120	A	-35.506	7.257 -13.534
677	OG	SER 120	A	-35.100	6.005 -14.079
678	N	SER 121	A	-36.217	4.669 -12.388
679	CA	SER 121	A	-35.537	3.572 -11.785
680	C	SER 121	A	-34.705	3.104 -12.949
681	O	SER 121	A	-34.958	2.007 -13.464
682	CB	SER 121	A	-36.551	2.525 -11.340
683	OG	SER 121	A	-37.674	3.246 -10.810
684	N	ASN 122	A	-33.701	3.800 -13.473
685	CA	ASN 122	A	-32.990	3.100 -14.519
686	C	ASN 122	A	-32.307	1.922 -13.858
687	O	ASN 122	A	-31.660	2.138 -12.857
688	CB	ASN 122	A	-31.913	3.941 -15.157
689	CG	ASN 122	A	-32.456	4.847 -16.232
690	OD1	ASN 122	A	-33.543	5.397 -16.193
691	ND2	ASN 122	A	-31.577	5.040 -17.223
692	N	VAL 123	A	-32.362	0.675 -14.271
693	CA	VAL 123	A	-31.447	-0.312 -13.748
694	C	VAL 123	A	-30.061	-0.155 -14.305
695	O	VAL 123	A	-29.858	-0.450 -15.460
696	CB	VAL 123	A	-31.992	-1.647 -14.081
697	CG1	VAL 123	A	-31.033	-2.769 -13.818
698	CG2	VAL 123	A	-33.230	-1.785 -13.234
699	N	ILE 124	A	-29.010	0.291 -13.637
700	CA	ILE 124	A	-27.704	0.244 -14.288
701	C	ILE 124	A	-26.583	-0.682 -13.814
702	O	ILE 124	A	-26.689	-1.441 -12.866
703	CB	ILE 124	A	-27.187	1.630 -14.333
704	CG1	ILE 124	A	-27.086	2.273 -13.025
705	CG2	ILE 124	A	-28.198	2.409 -15.116
706	CD1	ILE 124	A	-26.681	3.705 -13.362
707	N	LEU 125	A	-25.411	-0.711 -14.431
708	CA	LEU 125	A	-24.372	-1.639 -13.989
709	C	LEU 125	A	-23.210	-0.681 -13.840
710	O	LEU 125	A	-22.730	-0.263 -14.888
711	CB	LEU 125	A	-24.075	-2.659 -15.069
712	CG	LEU 125	A	-23.261	-3.900 -14.761
713	CD1	LEU 125	A	-23.927	-4.875 -13.791
714	CD2	LEU 125	A	-23.108	-4.618 -16.093
715	N	LYS 126	A	-22.754	-0.305 -12.615
716	CA	LYS 126	A	-21.529	0.463 -12.487
717	C	LYS 126	A	-20.416	-0.467 -12.106
718	O	LYS 126	A	-20.595	-1.572 -11.608
719	CB	LYS 126	A	-21.624	1.556 -11.437
720	CG	LYS 126	A	-20.629	2.603 -12.021

39/98

Fig. 15.17

Atom	Residuc	Chain	X	Y	Z
721	CD	LYS 126	A	-20.367	3.908 -11.203
722	CE	LYS 126	A	-19.079	4.686 -11.635
723	NZ	LYS 126	A	-17.843	3.882 -11.529
724	N	LYS 127	A	-19.199	-0.029 -12.352
725	CA	LYS 127	A	-18.017	-0.835 -12.143
726	C	LYS 127	A	-17.282	0.026 -11.185
727	O	LYS 127	A	-17.184	1.224 -11.370
728	CB	LYS 127	A	-17.097	-0.968 -13.339
729	CG	LYS 127	A	-16.235	-2.196 -13.072
730	CD	LYS 127	A	-15.334	-2.558 -14.231
731	CE	LYS 127	A	-14.497	-1.365 -14.645
732	NZ	LYS 127	A	-13.360	-1.909 -15.350
733	N	TYR 128	A	-16.763	-0.565 -10.145
734	CA	TYR 128	A	-16.120	0.206 -9.143
735	C	TYR 128	A	-14.784	-0.346 -9.316
736	O	TYR 128	A	-14.664	-1.559 -9.483
737	CB	TYR 128	A	-16.692	-0.131 -7.789
738	CG	TYR 128	A	-17.935	0.643 -7.636
739	CD1	TYR 128	A	-19.083	0.008 -7.229
740	CD2	TYR 128	A	-17.936	1.966 -7.982
741	CE1	TYR 128	A	-20.272	0.703 -7.216
742	CE2	TYR 128	A	-19.101	2.672 -7.977
743	CZ	TYR 128	A	-20.270	2.038 -7.613
744	OH	TYR 128	A	-21.474	2.754 -7.674
745	N	ARG 129	A	-13.741	0.463 -9.292
746	CA	ARG 129	A	-12.483	-0.169 -9.488
747	C	ARG 129	A	-11.895	-0.158 -8.145
748	O	ARG 129	A	-12.287	0.615 -7.288
749	CB	ARG 129	A	-11.616	0.626 -10.404
750	CG	ARG 129	A	-12.191	1.806 -11.208
751	CD	ARG 129	A	-11.134	2.214 -12.267
752	NE	ARG 129	A	-10.607	1.001 -12.893
753	CZ	ARG 129	A	-11.262	0.401 -13.888
754	NH1	ARG 129	A	-12.339	0.985 -14.529
755	NH2	ARG 129	A	-10.849	-0.846 -14.242
756	N	ASN 130	A	-10.935	-1.039 -7.949
757	CA	ASN 130	A	-10.085	-1.012 -6.790
758	C	ASN 130	A	-10.851	-1.277 -5.599
759	O	ASN 130	A	-10.708	-0.669 -4.572
760	CB	ASN 130	A	-9.393	0.323 -6.609
761	CG	ASN 130	A	-8.261	0.341 -7.600
762	OD1	ASN 130	A	-7.739	-0.694 -7.981
763	ND2	ASN 130	A	-7.853	1.492 -8.122
764	N	MET 131	A	-11.728	-2.248 -5.704
765	CA	MET 131	A	-12.531	-2.586 -4.577

40/98

Fig. 15.18

Atom	Residue	Chain	X	Y	Z	
766	C	MET 131	A	-12.155	-3.895	-3.926
767	O	MET 131	A	-12.642	-4.228	-2.858
768	CB	MET 131	A	-13.944	-2.576	-5.083
769	CG	MET 131	A	-14.347	-1.149	-5.256
770	SD	MET 131	A	-15.527	-0.836	-3.944
771	CE	MET 131	A	-14.884	0.789	-3.671
772	N	VAL 132	A	-11.291	-4.690	-4.521
773	CA	VAL 132	A	-11.025	-6.042	-4.048
774	C	VAL 132	A	-9.531	-6.189	-3.925
775	O	VAL 132	A	-8.827	-5.797	-4.819
776	CB	VAL 132	A	-11.556	-7.015	-5.070
777	CG1	VAL 132	A	-11.478	-8.446	-4.622
778	CG2	VAL 132	A	-13.002	-6.636	-5.321
779	N	VAL 133	A	-8.890	-6.710	-2.915
780	CA	VAL 133	A	-7.469	-6.920	-2.922
781	C	VAL 133	A	-7.287	-8.237	-3.624
782	O	VAL 133	A	-8.076	-9.130	-3.369
783	CB	VAL 133	A	-7.014	-6.965	-1.472
784	CG1	VAL 133	A	-5.786	-7.825	-1.195
785	CG2	VAL 133	A	-6.799	-5.522	-1.123
786	N	ARG 134	A	-6.278	-8.372	-4.493
787	CA	ARG 134	A	-5.826	-9.648	-5.013
788	C	ARG 134	A	-4.558	-10.087	-4.286
789	O	ARG 134	A	-4.366	-11.235	-3.880
790	CB	ARG 134	A	-5.517	-9.521	-6.505
791	CG	ARG 134	A	-6.403	-10.251	-7.540
792	CD	ARG 134	A	-5.677	-10.152	-8.920
793	NE	ARG 134	A	-6.330	-10.776	-10.106
794	CZ	ARG 134	A	-6.819	-12.047	-10.211
795	NH1	ARG 134	A	-7.257	-12.490	-11.438
796	NH2	ARG 134	A	-6.948	-12.890	-9.134
797	N	ALA 135	A	-3.591	-9.198	-4.086
798	CA	ALA 135	A	-2.487	-9.628	-3.277
799	C	ALA 135	A	-1.918	-8.469	-2.523
800	O	ALA 135	A	-2.177	-7.321	-2.854
801	CB	ALA 135	A	-1.466	-10.268	-4.196
802	N	CYS 136	A	-1.126	-8.787	-1.485
803	CA	CYS 136	A	-0.495	-7.839	-0.552
804	C	CYS 136	A	1.009	-8.034	-0.432
805	O	CYS 136	A	1.465	-9.160	-0.589
806	CB	CYS 136	A	-0.883	-7.996	0.872
807	SG	CYS 136	A	-2.530	-8.641	1.107
808	N	GLY 137	A	1.790	-7.010	-0.149
809	CA	GLY 137	A	3.125	-7.303	0.230
810	C	GLY 137	A	3.804	-6.043	0.544

41/98
Fig. 15.19

Atom	Residue	Chain	X	Y	Z	
811	O	GLY 137	A	3.166	-5.012	0.413
812	N	CYS 138	A	5.077	-6.032	0.947
813	CA	CYS 138	A	5.617	-4.923	1.740
814	C	CYS 138	A	6.031	-3.785	0.894
815	O	CYS 138	A	6.745	-4.136	-0.019
816	CB	CYS 138	A	6.815	-5.368	2.504
817	SG	CYS 138	A	6.442	-7.019	3.133
818	N	HIS 139	A	5.732	-2.511	1.016
819	CA	HIS 139	A	6.451	-1.586	0.176
820	C	HIS 139	A	6.716	-0.472	1.145
821	O	HIS 139	A	5.993	-0.333	2.154
822	CB	HIS 139	A	5.651	-1.066	-1.031
823	CG	HIS 139	A	5.126	-2.180	-1.889
824	ND1	HIS 139	A	4.592	-3.381	-1.428
825	CD2	HIS 139	A	5.565	-2.289	-3.201
826	CE1	HIS 139	A	4.749	-4.140	-2.529
827	NE2	HIS 139	A	5.322	-3.570	-3.624
828	OXT	HIS 139	A	7.321	0.362	0.777
829	N	GLN 36	B	-1.316	13.234	11.930
830	CA	GLN 36	B	-0.019	13.097	11.315
831	C	GLN 36	B	-0.372	12.092	10.267
832	O	GLN 36	B	-0.507	10.903	10.466
833	CB	GLN 36	B	0.994	12.511	12.324
834	CG	GLN 36	B	0.822	10.996	12.674
835	CD	GLN 36	B	-0.460	10.531	13.376
836	OE1	GLN 36	B	-1.426	11.270	13.478
837	NE2	GLN 36	B	-0.520	9.334	13.901
838	N	ALA 37	B	-0.523	12.730	9.128
839	CA	ALA 37	B	-1.432	12.344	8.073
840	C	ALA 37	B	-0.603	11.520	7.202
841	O	ALA 37	B	0.573	11.779	7.257
842	CB	ALA 37	B	-1.894	13.588	7.326
843	N	CYS 38	B	-0.971	10.577	6.394
844	CA	CYS 38	B	0.057	9.856	5.706
845	C	CYS 38	B	0.697	10.748	4.703
846	O	CYS 38	B	-0.011	11.155	3.781
847	CB	CYS 38	B	-0.585	8.677	5.078
848	SG	CYS 38	B	0.292	8.306	3.618
849	N	LYS 39	B	2.000	11.027	4.892
850	CA	LYS 39	B	2.823	11.772	3.937
851	C	LYS 39	B	4.228	11.167	3.694
852	O	LYS 39	B	4.679	10.270	4.389
853	CB	LYS 39	B	2.990	13.236	4.403
854	CG	LYS 39	B	3.093	13.354	5.921
855	CD	LYS 39	B	3.434	14.707	6.546

42/98
Fig. 15.20

Atom	Residue		Chain	X	Y	Z
856	CE	LYS 39	B	3.222	14.463	8.066
857	NZ	LYS 39	B	3.429	15.675	8.873
858	N	LYS 40	B	4.966	11.640	2.693
859	CA	LYS 40	B	6.314	11.192	2.434
860	C	LYS 40	B	7.224	11.807	3.433
861	O	LYS 40	B	6.889	12.880	3.892
862	CB	LYS 40	B	6.814	11.640	1.096
863	CG	LYS 40	B	8.068	10.922	0.629
864	CD	LYS 40	B	8.867	11.865	-0.243
865	CE	LYS 40	B	8.006	12.488	-1.310
866	NZ	LYS 40	B	8.797	13.441	-2.059
867	N	HIS 41	B	8.361	11.220	3.821
868	CA	HIS 41	B	9.283	11.844	4.760
869	C	HIS 41	B	10.701	11.549	4.352
870	O	HIS 41	B	10.914	10.791	3.425
871	CB	HIS 41	B	9.179	11.290	6.141
872	CG	HIS 41	B	7.860	11.488	6.797
873	ND1	HIS 41	B	7.584	12.100	7.948
874	CD2	HIS 41	B	6.778	10.752	6.444
875	CE1	HIS 41	B	6.404	11.711	8.330
876	NE2	HIS 41	B	5.937	10.898	7.415
877	N	GLU 42	B	11.751	12.069	4.966
878	CA	GLU 42	B	13.094	11.965	4.415
879	C	GLU 42	B	13.878	10.810	4.901
880	O	GLU 42	B	13.613	10.301	5.969
881	CB	GLU 42	B	13.875	13.207	4.718
882	CG	GLU 42	B	13.369	14.277	3.745
883	CD	GLU 42	B	14.490	15.264	3.400
884	OE1	GLU 42	B	14.161	16.446	3.123
885	OE2	GLU 42	B	15.681	14.846	3.405
886	N	LEU 43	B	14.868	10.303	4.208
887	CA	LEU 43	B	15.662	9.220	4.794
888	C	LEU 43	B	16.791	9.303	3.810
889	O	LEU 43	B	16.590	9.000	2.644
890	CB	LEU 43	B	15.082	7.834	4.628
891	CG	LEU 43	B	15.121	6.808	5.726
892	CD1	LEU 43	B	15.317	5.447	5.064
893	CD2	LEU 43	B	16.201	7.098	6.734
894	N	TYR 44	B	17.960	9.706	4.244
895	CA	TYR 44	B	19.116	9.691	3.414
896	C	TYR 44	B	19.731	8.374	3.684
897	O	TYR 44	B	20.074	8.206	4.838
898	CB	TYR 44	B	19.987	10.792	3.853
899	CG	TYR 44	B	21.075	10.941	2.875
900	CD1	TYR 44	B	22.293	10.457	3.244

43/98

Fig. 15.21

Atom	Residue	Chain	X	Y	Z	
901	CD2	TYR 44	B	20.850	11.476	1.648
902	CE1	TYR 44	B	23.318	10.366	2.365
903	CE2	TYR 44	B	21.874	11.387	0.757
904	CZ	TYR 44	B	23.066	10.774	1.099
905	OH	TYR 44	B	24.015	10.452	0.125
906	N	VAL 45	B	19.912	7.407	2.784
907	CA	VAL 45	B	20.665	6.229	3.164
908	C	VAL 45	B	22.087	6.437	2.654
909	O	VAL 45	B	22.279	6.805	1.507
910	CB	VAL 45	B	20.035	5.032	2.545
911	CG1	VAL 45	B	20.695	3.749	2.995
912	CG2	VAL 45	B	18.586	5.039	2.976
913	N	SER 46	B	23.114	6.225	3.477
914	CA	SER 46	B	24.493	6.439	3.107
915	C	SER 46	B	25.038	5.102	2.862
916	O	SER 46	B	24.941	4.287	3.750
917	CB	SER 46	B	25.291	7.048	4.229
918	OG	SER 46	B	26.655	7.224	3.852
919	N	PHE 47	B	25.629	4.746	1.749
920	CA	PHE 47	B	25.991	3.376	1.514
921	C	PHE 47	B	26.978	2.852	2.491
922	O	PHE 47	B	27.217	1.663	2.537
923	CB	PHE 47	B	26.594	3.221	0.184
924	CG	PHE 47	B	25.580	3.521	-0.868
925	CD1	PHE 47	B	24.477	2.747	-0.984
926	CD2	PHE 47	B	25.878	4.449	-1.823
927	CE1	PHE 47	B	23.731	2.826	-2.115
928	CE2	PHE 47	B	25.135	4.505	-2.974
929	CZ	PHE 47	B	24.071	3.673	-3.132
930	N	ARG 48	B	27.608	3.672	3.320
931	CA	ARG 48	B	28.339	3.167	4.458
932	C	ARG 48	B	27.398	2.413	5.320
933	O	ARG 48	B	27.646	1.284	5.611
934	CB	ARG 48	B	28.954	4.292	5.278
935	CG	ARG 48	B	30.046	5.035	4.494
936	CD	ARG 48	B	30.896	5.928	5.396
937	NE	ARG 48	B	32.154	6.274	4.756
938	CZ	ARG 48	B	32.651	7.514	4.856
939	NH1	ARG 48	B	31.950	8.482	5.519
940	NH2	ARG 48	B	33.859	7.849	4.304
941	N	ASP 49	B	26.272	2.920	5.771
942	CA	ASP 49	B	25.514	2.260	6.816
943	C	ASP 49	B	24.987	0.889	6.405
944	O	ASP 49	B	24.545	0.069	7.199
945	CB	ASP 49	B	24.328	3.105	7.229

44/98

Fig. 15.22

Atom	Residuc	Chain	X	Y	Z	
946	CG	ASP 49	B	24.718	4.526	7.549
947	OD1	ASP 49	B	23.837	5.362	7.347
948	OD2	ASP 49	B	25.851	4.840	7.969
949	N	LEU 50	B	25.005	0.563	5.125
950	CA	LEU 50	B	24.555	-0.751	4.716
951	C	LEU 50	B	25.746	-1.589	4.430
952	O	LEU 50	B	25.685	-2.699	3.895
953	CB	LEU 50	B	23.706	-0.649	3.461
954	CG	LEU 50	B	22.524	0.294	3.611
955	CD1	LEU 50	B	22.301	0.924	2.257
956	CD2	LEU 50	B	21.289	-0.430	4.147
957	N	GLY 51	B	26.887	-1.028	4.780
958	CA	GLY 51	B	28.143	-1.664	4.485
959	C	GLY 51	B	28.483	-1.605	3.017
960	O	GLY 51	B	29.593	-1.964	2.642
961	N	TRP 52	B	27.628	-1.178	2.086
962	CA	TRP 52	B	28.069	-1.113	0.692
963	C	TRP 52	B	29.287	-0.215	0.418
964	O	TRP 52	B	29.372	0.385	-0.629
965	CB	TRP 52	B	26.856	-0.656	-0.156
966	CG	TRP 52	B	25.726	-1.702	-0.123
967	CD1	TRP 52	B	25.857	-2.997	0.341
968	CD2	TRP 52	B	24.335	-1.394	-0.582
969	NE1	TRP 52	B	24.672	-3.554	0.236
970	CE2	TRP 52	B	23.731	-2.740	-0.270
971	CE3	TRP 52	B	23.505	-0.402	-1.076
972	CZ2	TRP 52	B	22.376	-2.951	-0.421
973	CZ3	TRP 52	B	22.157	-0.672	-1.220
974	CH2	TRP 52	B	21.604	-1.908	-0.885
975	N	GLN 53	B	30.313	-0.015	1.236
976	CA	GLN 53	B	30.998	1.273	1.141
977	C	GLN 53	B	32.040	1.358	0.116
978	O	GLN 53	B	32.702	2.374	-0.014
979	CB	GLN 53	B	31.742	1.728	2.405
980	CG	GLN 53	B	31.993	0.649	3.446
981	CD	GLN 53	B	33.274	0.967	4.237
982	OE1	GLN 53	B	34.189	0.121	4.274
983	NE2	GLN 53	B	33.399	2.156	4.889
984	N	ASP 54	B	32.235	0.306	-0.647
985	CA	ASP 54	B	33.600	-0.101	-0.948
986	C	ASP 54	B	33.381	-0.824	-2.214
987	O	ASP 54	B	33.245	-2.036	-2.082
988	CB	ASP 54	B	34.116	-1.083	0.114
989	CG	ASP 54	B	32.963	-1.909	0.747
990	OD1	ASP 54	B	33.101	-2.317	1.928

45/98

Fig. 15.23

Atom	Residue	Chain	X	Y	Z	
991	OD2	ASP 54	B	31.940	-2.112	0.062
992	N	TRP 55	B	33.314	-0.266	-3.427
993	CA	TRP 55	B	32.570	-1.004	-4.476
994	C	TRP 55	B	32.019	-0.003	-5.421
995	O	TRP 55	B	32.165	-0.187	-6.621
996	CB	TRP 55	B	31.258	-1.689	-4.123
997	CG	TRP 55	B	31.339	-3.126	-3.756
998	CD1	TRP 55	B	32.336	-4.015	-4.070
999	CD2	TRP 55	B	30.132	-3.715	-3.105
1000	NE1	TRP 55	B	31.855	-5.199	-3.695
1001	CE2	TRP 55	B	30.594	-5.172	-3.148
1002	CE3	TRP 55	B	28.951	-3.382	-2.460
1003	CZ2	TRP 55	B	29.753	-6.141	-2.581
1004	CZ3	TRP 55	B	28.177	-4.393	-1.896
1005	CH2	TRP 55	B	28.549	-5.738	-1.967
1006	N	ILE 56	B	31.399	1.020	-4.769
1007	CA	ILE 56	B	30.572	2.056	-5.356
1008	C	ILE 56	B	31.459	3.266	-5.650
1009	O	ILE 56	B	32.148	3.707	-4.741
1010	CB	ILE 56	B	29.469	2.363	-4.346
1011	CG1	ILE 56	B	28.247	1.655	-4.774
1012	CG2	ILE 56	B	29.164	3.835	-4.244
1013	CD1	ILE 56	B	28.212	0.333	-4.085
1014	N	ILE 57	B	31.511	3.841	-6.857
1015	CA	ILE 57	B	32.213	5.100	-7.039
1016	C	ILE 57	B	31.178	6.206	-6.753
1017	O	ILE 57	B	31.414	7.392	-6.654
1018	CB	ILE 57	B	32.735	5.162	-8.493
1019	CG1	ILE 57	B	33.659	4.001	-8.726
1020	CG2	ILE 57	B	33.527	6.433	-8.784
1021	CD1	ILE 57	B	34.001	4.036	-10.220
1022	N	ALA 58	B	29.934	5.831	-6.570
1023	CA	ALA 58	B	28.802	5.950	-7.504
1024	C	ALA 58	B	28.040	7.166	-7.145
1025	O	ALA 58	B	28.524	8.150	-7.690
1026	CB	ALA 58	B	27.934	4.732	-7.338
1027	N	PRO 59	B	26.999	7.188	-6.358
1028	CA	PRO 59	B	26.820	8.263	-5.382
1029	C	PRO 59	B	27.333	7.862	-4.008
1030	O	PRO 59	B	27.512	6.698	-3.744
1031	CB	PRO 59	B	25.317	8.611	-5.384
1032	CG	PRO 59	B	24.697	7.280	-5.680
1033	CD	PRO 59	B	25.659	6.734	-6.707
1034	N	GLU 60	B	27.622	8.703	-3.030
1035	CA	GLU 60	B	27.941	8.146	-1.719

46/98
Fig. 15.24

Atom	Residue			Chain	X	Y	Z
1036	C	GLU	60	B	26.735	7.734	-0.926
1037	O	GLU	60	B	26.813	7.132	0.131
1038	CB	GLU	60	B	28.745	9.151	-0.867
1039	CG	GLU	60	B	27.995	10.280	-0.117
1040	CD	GLU	60	B	27.231	9.868	1.144
1041	OE1	GLU	60	B	27.559	8.834	1.775
1042	OE2	GLU	60	B	26.329	10.634	1.518
1043	N	GLY	61	B	25.552	8.049	-1.395
1044	CA	GLY	61	B	24.342	7.623	-0.733
1045	C	GLY	61	B	23.253	8.252	-1.519
1046	O	GLY	61	B	23.535	8.769	-2.595
1047	N	TYR	62	B	22.021	8.246	-1.038
1048	CA	TYR	62	B	20.934	8.777	-1.830
1049	C	TYR	62	B	19.767	9.046	-0.932
1050	O	TYR	62	B	19.796	8.647	0.217
1051	CB	TYR	62	B	20.608	7.745	-2.889
1052	CG	TYR	62	B	19.833	6.572	-2.370
1053	CD1	TYR	62	B	20.425	5.395	-1.917
1054	CD2	TYR	62	B	18.481	6.692	-2.472
1055	CE1	TYR	62	B	19.624	4.316	-1.563
1056	CE2	TYR	62	B	17.692	5.644	-2.132
1057	CZ	TYR	62	B	18.249	4.475	-1.681
1058	OH	TYR	62	B	17.341	3.468	-1.352
1059	N	ALA	63	B	18.720	9.711	-1.407
1060	CA	ALA	63	B	17.474	9.863	-0.693
1061	C	ALA	63	B	16.485	8.722	-0.918
1062	O	ALA	63	B	15.903	8.617	-1.981
1063	CB	ALA	63	B	16.899	11.151	-1.155
1064	N	ALA	64	B	16.262	7.847	0.047
1065	CA	ALA	64	B	15.293	6.793	-0.077
1066	C	ALA	64	B	13.847	7.203	0.227
1067	O	ALA	64	B	12.851	6.701	-0.265
1068	CB	ALA	64	B	15.695	5.679	0.866
1069	N	TYR	65	B	13.710	8.191	1.098
1070	CA	TYR	65	B	12.454	8.570	1.709
1071	C	TYR	65	B	11.742	7.437	2.314
1072	O	TYR	65	B	12.101	6.294	2.178
1073	CB	TYR	65	B	11.531	9.231	0.716
1074	CG	TYR	65	B	12.088	10.513	0.168
1075	CD1	TYR	65	B	12.540	11.505	0.984
1076	CD2	TYR	65	B	12.170	10.655	-1.176
1077	CE1	TYR	65	B	13.147	12.614	0.454
1078	CE2	TYR	65	B	12.771	11.751	-1.721
1079	CZ	TYR	65	B	13.282	12.701	-0.902
1080	OH	TYR	65	B	13.974	13.762	-1.462

47/98
Fig. 15.25

Atom	Residuc	Chain	X	Y	Z	
1081	N	TYR 66	B	10.697	7.722	3.031
1082	CA	TYR 66	B	9.852	6.661	3.471
1083	C	TYR 66	B	8.541	7.291	3.740
1084	O	TYR 66	B	8.435	8.508	3.867
1085	CB	TYR 66	B	10.389	5.996	4.739
1086	CG	TYR 66	B	10.325	6.790	6.014
1087	CD1	TYR 66	B	11.182	7.824	6.193
1088	CD2	TYR 66	B	9.448	6.427	7.006
1089	CE1	TYR 66	B	11.162	8.514	7.369
1090	CE2	TYR 66	B	9.415	7.105	8.185
1091	CZ	TYR 66	B	10.273	8.151	8.343
1092	OH	TYR 66	B	10.254	8.912	9.507
1093	N	CYS 67	B	7.523	6.461	3.838
1094	CA	CYS 67	B	6.157	6.944	3.974
1095	C	CYS 67	B	5.600	6.711	5.382
1096	O	CYS 67	B	5.660	5.589	5.909
1097	CB	CYS 67	B	5.210	6.219	3.078
1098	SG	CYS 67	B	5.398	6.240	1.331
1099	N	GLU 68	B	5.026	7.712	6.056
1100	CA	GLU 68	B	4.554	7.337	7.342
1101	C	GLU 68	B	3.507	8.263	7.751
1102	O	GLU 68	B	3.670	9.457	7.509
1103	CB	GLU 68	B	5.639	7.411	8.360
1104	CG	GLU 68	B	5.322	6.494	9.510
1105	CD	GLU 68	B	6.024	7.006	10.718
1106	OE1	GLU 68	B	6.127	6.266	11.697
1107	OE2	GLU 68	B	6.475	8.154	10.689
1108	N	GLY 69	B	2.450	7.734	8.369
1109	CA	GLY 69	B	1.458	8.577	9.002
1110	C	GLY 69	B	0.123	7.873	8.939
1111	O	GLY 69	B	0.019	6.722	8.517
1112	N	GLU 70	B	-0.962	8.499	9.336
1113	CA	GLU 70	B	-2.120	7.685	9.616
1114	C	GLU 70	B	-3.133	7.672	8.511
1115	O	GLU 70	B	-3.372	8.697	7.898
1116	CB	GLU 70	B	-2.769	8.180	10.882
1117	CG	GLU 70	B	-2.975	6.949	11.748
1118	CD	GLU 70	B	-4.167	7.206	12.630
1119	OE1	GLU 70	B	-4.866	6.234	12.949
1120	OE2	GLU 70	B	-4.412	8.381	12.984
1121	N	CYS 71	B	-3.757	6.544	8.233
1122	CA	CYS 71	B	-4.817	6.538	7.262
1123	C	CYS 71	B	-6.205	6.516	7.840
1124	O	CYS 71	B	-6.730	5.505	8.245
1125	CB	CYS 71	B	-4.553	5.350	6.377

48/98

Fig. 15.26

Atom	Residue	Chain	X	Y	Z	
1126	SG	CYS 71	B	-3.363	5.857	5.112
1127	N	ALA 72	B	-6.909	7.619	7.924
1128	CA	ALA 72	B	-8.091	7.666	8.737
1129	C	ALA 72	B	-8.935	8.777	8.184
1130	O	ALA 72	B	-8.438	9.590	7.418
1131	CB	ALA 72	B	-7.646	7.939	10.149
1132	N	PHE 73	B	-10.225	8.934	8.471
1133	CA	PHE 73	B	-10.963	9.998	7.782
1134	C	PHE 73	B	-10.507	11.313	8.342
1135	O	PHE 73	B	-10.291	11.328	9.544
1136	CB	PHE 73	B	-12.492	9.874	7.997
1137	CG	PHE 73	B	-13.023	8.690	7.235
1138	CD1	PHE 73	B	-13.009	8.688	5.873
1139	CD2	PHE 73	B	-13.514	7.616	7.919
1140	CE1	PHE 73	B	-13.473	7.602	5.190
1141	CE2	PHE 73	B	-13.992	6.528	7.214
1142	CZ	PHE 73	B	-13.969	6.518	5.854
1143	N	PRO 74	B	-10.337	12.410	7.712
1144	CA	PRO 74	B	-10.748	12.685	6.357
1145	C	PRO 74	B	-9.754	12.033	5.502
1146	O	PRO 74	B	-8.656	12.529	5.443
1147	CB	PRO 74	B	-10.712	14.168	6.197
1148	CG	PRO 74	B	-10.432	14.652	7.611
1149	CD	PRO 74	B	-9.662	13.546	8.313
1150	N	LEU 75	B	-10.018	10.938	4.799
1151	CA	LEU 75	B	-9.091	10.355	3.845
1152	C	LEU 75	B	-9.061	11.400	2.779
1153	O	LEU 75	B	-9.752	11.427	1.772
1154	CB	LEU 75	B	-9.674	9.034	3.355
1155	CG	LEU 75	B	-8.684	7.913	3.407
1156	CD1	LEU 75	B	-7.689	8.120	4.525
1157	CD2	LEU 75	B	-9.442	6.631	3.626
1158	N	ASN 76	B	-8.181	12.333	3.059
1159	CA	ASN 76	B	-8.098	13.558	2.311
1160	C	ASN 76	B	-7.517	13.432	0.887
1161	O	ASN 76	B	-6.525	14.065	0.596
1162	CB	ASN 76	B	-7.310	14.402	3.255
1163	CG	ASN 76	B	-7.466	15.837	2.965
1164	OD1	ASN 76	B	-7.449	16.655	3.894
1165	ND2	ASN 76	B	-7.521	16.219	1.674
1166	N	SER 77	B	-8.021	12.652	-0.088
1167	CA	SER 77	B	-7.673	12.747	-1.523
1168	C	SER 77	B	-6.200	12.824	-1.891
1169	O	SER 77	B	-5.587	11.803	-2.261
1170	CB	SER 77	B	-8.354	13.964	-2.171

49/98
Fig. 15.27

Atom	Residue	Chain	X	Y	Z	
1171	OG	SER 77	B	-7.957	15.177	-1.532
1172	N	TYR 78	B	-5.531	13.986	-1.819
1173	CA	TYR 78	B	-4.098	13.882	-1.965
1174	C	TYR 78	B	-3.490	13.159	-0.764
1175	O	TYR 78	B	-2.585	13.733	-0.185
1176	CB	TYR 78	B	-3.438	15.241	-2.045
1177	CG	TYR 78	B	-2.916	15.632	-3.429
1178	CD1	TYR 78	B	-1.897	16.590	-3.392
1179	CD2	TYR 78	B	-3.617	15.385	-4.630
1180	CE1	TYR 78	B	-1.737	17.490	-4.438
1181	CE2	TYR 78	B	-3.456	16.292	-5.692
1182	CZ	TYR 78	B	-2.610	17.403	-5.516
1183	OH	TYR 78	B	-2.715	18.527	-6.319
1184	N	MET 79	B	-3.978	11.959	-0.393
1185	CA	MET 79	B	-3.388	10.933	0.448
1186	C	MET 79	B	-3.689	9.603	-0.251
1187	O	MET 79	B	-3.734	8.505	0.285
1188	CB	MET 79	B	-4.031	10.775	1.776
1189	CG	MET 79	B	-4.050	11.928	2.741
1190	SD	MET 79	B	-4.737	11.135	4.215
1191	CE	MET 79	B	-4.910	12.550	5.255
1192	N	ASN 80	B	-3.926	9.699	-1.544
1193	CA	ASN 80	B	-4.199	8.581	-2.401
1194	C	ASN 80	B	-5.079	7.439	-1.962
1195	O	ASN 80	B	-4.924	6.364	-2.490
1196	CB	ASN 80	B	-2.883	7.967	-2.894
1197	CG	ASN 80	B	-3.155	7.090	-4.121
1198	OD1	ASN 80	B	-3.135	5.844	-4.162
1199	ND2	ASN 80	B	-3.543	7.769	-5.198
1200	N	ALA 81	B	-6.051	7.407	-1.058
1201	CA	ALA 81	B	-7.299	6.757	-1.424
1202	C	ALA 81	B	-7.486	5.344	-1.928
1203	O	ALA 81	B	-8.004	4.449	-1.302
1204	CB	ALA 81	B	-8.004	7.624	-2.452
1205	N	THR 82	B	-7.106	4.970	-3.106
1206	CA	THR 82	B	-7.875	3.957	-3.819
1207	C	THR 82	B	-9.402	3.970	-3.655
1208	O	THR 82	B	-10.075	4.641	-4.444
1209	CB	THR 82	B	-7.505	2.477	-3.539
1210	OG1	THR 82	B	-7.534	2.217	-2.163
1211	CG2	THR 82	B	-6.211	2.155	-4.224
1212	N	ASN 83	B	-10.043	3.286	-2.697
1213	CA	ASN 83	B	-11.462	2.977	-2.877
1214	C	ASN 83	B	-11.670	1.861	-1.963
1215	O	ASN 83	B	-12.472	1.858	-1.070

50/98
Fig. 15.28

Atom	Residue	Chain	X	Y	Z	
1216	CB	ASN 83	B	-11.835	2.404	-4.195
1217	CG	ASN 83	B	-12.710	3.302	-4.983
1218	OD1	ASN 83	B	-13.343	2.872	-5.931
1219	ND2	ASN 83	B	-12.741	4.594	-4.715
1220	N	HIS 84	B	-10.950	0.775	-2.095
1221	CA	HIS 84	B	-10.823	-0.192	-1.024
1222	C	HIS 84	B	-10.284	0.531	0.167
1223	O	HIS 84	B	-10.623	0.097	1.245
1224	CB	HIS 84	B	-9.851	-1.284	-1.397
1225	CG	HIS 84	B	-9.778	-2.364	-0.370
1226	ND1	HIS 84	B	-9.014	-2.402	0.711
1227	CD2	HIS 84	B	-10.397	-3.578	-0.489
1228	CE1	HIS 84	B	-9.153	-3.590	1.235
1229	NE2	HIS 84	B	-9.960	-4.294	0.507
1230	N	ALA 85	B	-9.464	1.589	0.169
1231	CA	ALA 85	B	-9.158	2.136	1.455
1232	C	ALA 85	B	-10.314	2.915	1.965
1233	O	ALA 85	B	-10.388	3.113	3.155
1234	CB	ALA 85	B	-7.958	3.026	1.398
1235	N	ILE 86	B	-11.298	3.418	1.242
1236	CA	ILE 86	B	-12.366	4.096	1.919
1237	C	ILE 86	B	-13.219	2.964	2.461
1238	O	ILE 86	B	-13.597	3.036	3.590
1239	CB	ILE 86	B	-13.058	4.975	0.897
1240	CG1	ILE 86	B	-12.035	5.957	0.299
1241	CG2	ILE 86	B	-14.230	5.699	1.565
1242	CD1	ILE 86	B	-12.568	7.113	-0.547
1243	N	VAL 87	B	-13.596	1.857	1.829
1244	CA	VAL 87	B	-14.315	0.785	2.493
1245	C	VAL 87	B	-13.560	0.425	3.744
1246	O	VAL 87	B	-14.049	0.656	4.828
1247	CB	VAL 87	B	-14.432	-0.429	1.553
1248	CG1	VAL 87	B	-14.652	-1.712	2.308
1249	CG2	VAL 87	B	-15.617	-0.236	0.637
1250	N	GLN 88	B	-12.358	-0.137	3.734
1251	CA	GLN 88	B	-11.706	-0.524	4.975
1252	C	GLN 88	B	-11.666	0.574	6.022
1253	O	GLN 88	B	-11.943	0.280	7.170
1254	CB	GLN 88	B	-10.295	-0.941	4.697
1255	CG	GLN 88	B	-9.864	-2.322	5.125
1256	CD	GLN 88	B	-8.398	-2.237	5.020
1257	OE1	GLN 88	B	-7.618	-2.317	5.944
1258	NE2	GLN 88	B	-7.922	-1.853	3.852
1259	N	THR 89	B	-11.347	1.833	5.754
1260	CA	THR 89	B	-11.401	2.802	6.790

51/98
Fig. 15.29

Atom		Residuc	Chain	X	Y	Z
1261	C	THR 89	B	-12.805	2.953	7.298
1262	O	THR 89	B	-12.966	3.090	8.499
1263	CB	THR 89	B	-10.923	4.081	6.270
1264	OG1	THR 89	B	-9.628	3.792	5.879
1265	CG2	THR 89	B	-10.741	5.183	7.270
1266	N	LEU 90	B	-13.888	2.943	6.506
1267	CA	LEU 90	B	-15.260	2.885	7.003
1268	C	LEU 90	B	-15.550	1.649	7.814
1269	O	LEU 90	B	-16.030	1.727	8.929
1270	CB	LEU 90	B	-16.220	2.893	5.853
1271	CG	LEU 90	B	-17.670	3.011	6.158
1272	CD1	LEU 90	B	-17.933	4.270	6.954
1273	CD2	LEU 90	B	-18.406	3.003	4.837
1274	N	VAL 91	B	-15.296	0.449	7.333
1275	CA	VAL 91	B	-15.459	-0.745	8.132
1276	C	VAL 91	B	-14.637	-0.671	9.387
1277	O	VAL 91	B	-14.959	-1.278	10.398
1278	CB	VAL 91	B	-15.049	-1.923	7.308
1279	CG1	VAL 91	B	-14.917	-3.200	8.111
1280	CG2	VAL 91	B	-16.122	-2.071	6.250
1281	N	HIS 92	B	-13.533	0.068	9.434
1282	CA	HIS 92	B	-12.818	0.180	10.702
1283	C	HIS 92	B	-13.719	0.784	11.705
1284	O	HIS 92	B	-13.859	0.310	12.797
1285	CB	HIS 92	B	-11.577	1.095	10.619
1286	CG	HIS 92	B	-10.675	0.908	11.812
1287	ND1	HIS 92	B	-10.106	1.714	12.701
1288	CD2	HIS 92	B	-10.314	-0.348	12.142
1289	CE1	HIS 92	B	-9.454	0.950	13.527
1290	NE2	HIS 92	B	-9.602	-0.300	13.197
1291	N	PHE 93	B	-14.364	1.854	11.358
1292	CA	PHE 93	B	-15.148	2.645	12.248
1293	C	PHE 93	B	-16.394	1.899	12.634
1294	O	PHE 93	B	-16.782	1.756	13.777
1295	CB	PHE 93	B	-15.365	3.894	11.461
1296	CG	PHE 93	B	-16.515	4.726	11.911
1297	CD1	PHE 93	B	-17.812	4.330	11.649
1298	CD2	PHE 93	B	-16.248	5.977	12.448
1299	CE1	PHE 93	B	-18.832	5.210	11.890
1300	CE2	PHE 93	B	-17.282	6.860	12.670
1301	CZ	PHE 93	B	-18.564	6.467	12.386
1302	N	ILE 94	B	-17.172	1.345	11.730
1303	CA	ILE 94	B	-18.251	0.460	12.158
1304	C	ILE 94	B	-17.714	-0.589	13.105
1305	O	ILE 94	B	-18.482	-1.254	13.758

52/98

Fig. 15.30

Atom	Residue	Chain	X	Y	Z	
1306	CB	ILE 94	B	-18.808	-0.124	10.880
1307	CG1	ILE 94	B	-19.580	1.028	10.342
1308	CG2	ILE 94	B	-19.648	-1.375	10.977
1309	CD1	ILE 94	B	-19.872	0.740	8.877
1310	N	ASN 95	B	-16.410	-0.839	13.277
1311	CA	ASN 95	B	-16.013	-2.092	13.873
1312	C	ASN 95	B	-14.529	-2.322	14.157
1313	O	ASN 95	B	-13.858	-3.248	13.673
1314	CB	ASN 95	B	-16.495	-3.208	12.993
1315	CG	ASN 95	B	-16.663	-4.358	13.942
1316	OD1	ASN 95	B	-16.021	-4.431	15.007
1317	ND2	ASN 95	B	-17.547	-5.309	13.564
1318	N	PRO 96	B	-13.981	-1.497	14.982
1319	CA	PRO 96	B	-12.565	-1.248	15.087
1320	C	PRO 96	B	-11.697	-2.421	15.300
1321	O	PRO 96	B	-10.491	-2.366	15.067
1322	CB	PRO 96	B	-12.426	-0.227	16.199
1323	CG	PRO 96	B	-13.819	0.347	16.282
1324	CD	PRO 96	B	-14.625	-0.944	16.151
1325	N	GLU 97	B	-12.338	-3.478	15.747
1326	CA	GLU 97	B	-11.625	-4.700	15.963
1327	C	GLU 97	B	-11.796	-5.472	14.691
1328	O	GLU 97	B	-10.808	-5.919	14.166
1329	CB	GLU 97	B	-12.209	-5.474	17.153
1330	CG	GLU 97	B	-11.332	-5.321	18.423
1331	CD	GLU 97	B	-11.552	-3.957	19.136
1332	OE1	GLU 97	B	-12.507	-3.902	19.971
1333	OE2	GLU 97	B	-10.781	-2.974	18.873
1334	N	THR 98	B	-12.921	-5.722	14.047
1335	CA	THR 98	B	-12.848	-6.579	12.862
1336	C	THR 98	B	-11.921	-6.174	11.682
1337	O	THR 98	B	-11.787	-7.035	10.815
1338	CB	THR 98	B	-14.289	-6.757	12.349
1339	OG1	THR 98	B	-14.841	-5.463	12.337
1340	CG2	THR 98	B	-15.226	-7.552	13.281
1341	N	VAL 99	B	-11.255	-5.008	11.488
1342	CA	VAL 99	B	-10.426	-4.763	10.307
1343	C	VAL 99	B	-9.447	-3.636	10.637
1344	O	VAL 99	B	-9.956	-2.741	11.289
1345	CB	VAL 99	B	-11.316	-4.375	9.112
1346	CG1	VAL 99	B	-11.405	-2.876	8.750
1347	CG2	VAL 99	B	-10.670	-5.076	7.950
1348	N	PRO 100	B	-8.179	-3.588	10.300
1349	CA	PRO 100	B	-7.213	-2.555	10.669
1350	C	PRO 100	B	-7.409	-1.258	10.007

53/98

Fig. 15.31

Atom	Residue	Chain	X	Y	Z	
1351	O	PRO 100	B	-8.381	-1.158	9.326
1352	CB	PRO 100	B	-5.890	-3.130	10.342
1353	CG	PRO 100	B	-6.120	-4.613	10.443
1354	CD	PRO 100	B	-7.484	-4.771	9.830
1355	N	LYS 101	B	-6.660	-0.169	10.060
1356	CA	LYS 101	B	-7.015	0.958	9.218
1357	C	LYS 101	B	-6.112	0.771	8.015
1358	O	LYS 101	B	-5.130	0.061	8.142
1359	CB	LYS 101	B	-6.671	2.300	9.855
1360	CG	LYS 101	B	-6.510	2.347	11.360
1361	CD	LYS 101	B	-5.813	3.616	11.889
1362	CE	LYS 101	B	-4.227	3.513	12.046
1363	NZ	LYS 101	B	-3.387	3.806	10.849
1364	N	PRO 102	B	-6.219	1.262	6.840
1365	CA	PRO 102	B	-5.321	0.916	5.769
1366	C	PRO 102	B	-3.970	1.433	6.090
1367	O	PRO 102	B	-3.853	2.346	6.892
1368	CB	PRO 102	B	-5.934	1.534	4.529
1369	CG	PRO 102	B	-6.576	2.741	5.074
1370	CD	PRO 102	B	-7.100	2.341	6.447
1371	N	CYS 103	B	-2.852	0.963	5.547
1372	CA	CYS 103	B	-1.653	1.665	5.922
1373	C	CYS 103	B	-0.993	2.294	4.757
1374	O	CYS 103	B	-1.354	2.179	3.600
1375	CB	CYS 103	B	-0.697	0.722	6.598
1376	SG	CYS 103	B	0.104	-0.064	5.231
1377	N	CYS 104	B	0.044	2.988	5.123
1378	CA	CYS 104	B	0.501	4.087	4.348
1379	C	CYS 104	B	1.771	3.676	3.652
1380	O	CYS 104	B	2.791	3.408	4.251
1381	CB	CYS 104	B	0.602	5.107	5.385
1382	SG	CYS 104	B	1.414	6.474	4.692
1383	N	ALA 105	B	1.749	3.613	2.347
1384	CA	ALA 105	B	2.637	2.781	1.614
1385	C	ALA 105	B	3.081	3.640	0.494
1386	O	ALA 105	B	2.412	4.644	0.296
1387	CB	ALA 105	B	1.841	1.604	1.154
1388	N	PRO 106	B	4.100	3.411	-0.258
1389	CA	PRO 106	B	4.400	4.197	-1.420
1390	C	PRO 106	B	3.597	3.811	-2.662
1391	O	PRO 106	B	3.311	2.672	-2.978
1392	CB	PRO 106	B	5.891	4.039	-1.557
1393	CG	PRO 106	B	6.123	2.662	-1.081
1394	CD	PRO 106	B	5.349	2.826	0.199
1395	N	THR 107	B	3.246	4.864	-3.375

54/98

Fig. 15.32

Atom	Residuc	Chain	X	Y	Z	
1396	CA	THR 107	B	2.321	4.885	-4.464
1397	C	THR 107	B	3.250	4.827	-5.644
1398	O	THR 107	B	3.206	3.871	-6.411
1399	CB	THR 107	B	1.582	6.175	-4.155
1400	OG1	THR 107	B	0.518	5.644	-3.377
1401	CG2	THR 107	B	1.134	7.066	-5.292
1402	N	GLN 108	B	4.151	5.763	-5.922
1403	CA	GLN 108	B	5.164	5.360	-6.846
1404	C	GLN 108	B	6.603	5.599	-6.478
1405	O	GLN 108	B	6.989	6.589	-5.880
1406	CB	GLN 108	B	4.712	6.001	-8.168
1407	CG	GLN 108	B	5.202	7.362	-8.639
1408	CD	GLN 108	B	4.487	8.380	-7.859
1409	OE1	GLN 108	B	3.911	8.140	-6.806
1410	NE2	GLN 108	B	4.552	9.578	-8.441
1411	N	LEU 109	B	7.351	4.579	-6.896
1412	CA	LEU 109	B	8.773	4.347	-6.622
1413	C	LEU 109	B	9.747	4.700	-7.774
1414	O	LEU 109	B	9.510	4.496	-8.963
1415	CB	LEU 109	B	8.945	2.848	-6.208
1416	CG	LEU 109	B	8.351	2.584	-4.813
1417	CD1	LEU 109	B	8.260	1.128	-4.489
1418	CD2	LEU 109	B	9.269	3.213	-3.792
1419	N	ASN 110	B	10.911	5.262	-7.494
1420	CA	ASN 110	B	11.823	5.551	-8.555
1421	C	ASN 110	B	13.077	4.792	-8.391
1422	O	ASN 110	B	13.238	4.036	-7.445
1423	CB	ASN 110	B	12.199	6.976	-8.586
1424	CG	ASN 110	B	11.025	7.786	-8.983
1425	OD1	ASN 110	B	11.064	9.002	-8.744
1426	ND2	ASN 110	B	9.942	7.206	-9.529
1427	N	ALA 111	B	14.021	4.976	-9.308
1428	CA	ALA 111	B	15.130	4.055	-9.447
1429	C	ALA 111	B	16.280	4.897	-9.088
1430	O	ALA 111	B	16.134	6.101	-9.021
1431	CB	ALA 111	B	15.249	3.543	-10.900
1432	N	ILE 112	B	17.436	4.324	-8.843
1433	CA	ILE 112	B	18.624	5.094	-8.502
1434	C	ILE 112	B	19.626	4.386	-9.369
1435	O	ILE 112	B	19.520	3.175	-9.473
1436	CB	ILE 112	B	18.834	4.979	-6.944
1437	CG1	ILE 112	B	18.727	6.452	-6.591
1438	CG2	ILE 112	B	20.119	4.396	-6.342
1439	CD1	ILE 112	B	20.014	7.252	-6.922
1440	N	SER 113	B	20.575	5.052	-9.999

55/98

Fig. 15.33

Atom	Residue	Chain	X	Y	Z
1441	CA	SER 113	B	21.646	4.330 -10.635
1442	C	SER 113	B	22.926	4.454 -9.879
1443	O	SER 113	B	23.220	5.579 -9.520
1444	CB	SER 113	B	21.834	4.850 -12.034
1445	OG	SER 113	B	21.085	3.985 -12.909
1446	N	VAL 114	B	23.726	3.408 -9.592
1447	CA	VAL 114	B	25.005	3.645 -8.929
1448	C	VAL 114	B	26.085	3.074 -9.841
1449	O	VAL 114	B	25.914	2.046 -10.484
1450	CB	VAL 114	B	25.150	2.977 -7.496
1451	CG1	VAL 114	B	23.888	3.143 -6.703
1452	CG2	VAL 114	B	25.446	1.508 -7.575
1453	N	LEU 115	B	27.215	3.773 -9.911
1454	CA	LEU 115	B	28.417	3.420 -10.663
1455	C	LEU 115	B	29.406	2.624 -9.859
1456	O	LEU 115	B	29.857	3.095 -8.814
1457	CB	LEU 115	B	29.096	4.718 -11.145
1458	CG	LEU 115	B	30.445	4.721 -11.850
1459	CD1	LEU 115	B	30.271	4.338 -13.289
1460	CD2	LEU 115	B	31.054	6.115 -11.750
1461	N	TYR 116	B	29.854	1.432 -10.212
1462	CA	TYR 116	B	30.750	0.783 -9.287
1463	C	TYR 116	B	31.728	-0.199 -9.868
1464	O	TYR 116	B	31.498	-0.667 -10.976
1465	CB	TYR 116	B	29.872	0.123 -8.278
1466	CG	TYR 116	B	29.144	-1.012 -8.907
1467	CD1	TYR 116	B	27.919	-0.788 -9.480
1468	CD2	TYR 116	B	29.728	-2.243 -8.856
1469	CE1	TYR 116	B	27.289	-1.824 -10.101
1470	CE2	TYR 116	B	29.105	-3.280 -9.474
1471	CZ	TYR 116	B	27.925	-3.035 -10.113
1472	OH	TYR 116	B	27.396	-4.037 -10.879
1473	N	PHE 117	B	32.820	-0.583 -9.201
1474	CA	PHE 117	B	33.733	-1.520 -9.828
1475	C	PHE 117	B	33.168	-2.861 -9.521
1476	O	PHE 117	B	33.029	-3.159 -8.347
1477	CB	PHE 117	B	35.137	-1.590 -9.242
1478	CG	PHE 117	B	35.701	-0.236 -8.937
1479	CD1	PHE 117	B	35.512	0.315 -7.691
1480	CD2	PHE 117	B	36.392	0.459 -9.903
1481	CE1	PHE 117	B	36.002	1.579 -7.433
1482	CE2	PHE 117	B	36.872	1.732 -9.656
1483	CZ	PHE 117	B	36.672	2.294 -8.412
1484	N	ASP 118	B	32.811	-3.756 -10.426
1485	CA	ASP 118	B	32.630	-5.124 -9.935

56/98

Fig. 15.34

Atom	Residue	Chain	X	Y	Z	
1486	C	ASP 118	B	34.049	-5.641	-9.774
1487	O	ASP 118	B	35.001	-4.878	-9.844
1488	CB	ASP 118	B	31.868	-6.043	-10.948
1489	CG	ASP 118	B	32.575	-6.053	-12.291
1490	OD1	ASP 118	B	32.115	-6.790	-13.188
1491	OD2	ASP 118	B	33.580	-5.313	-12.438
1492	N	ASP 119	B	34.193	-6.943	-9.569
1493	CA	ASP 119	B	35.472	-7.584	-9.692
1494	C	ASP 119	B	36.172	-7.231	-11.043
1495	O	ASP 119	B	35.617	-6.552	-11.918
1496	CB	ASP 119	B	35.288	-9.105	-9.596
1497	CG	ASP 119	B	34.198	-9.528	-10.579
1498	OD1	ASP 119	B	34.168	-8.947	-11.699
1499	OD2	ASP 119	B	33.370	-10.399	-10.216
1500	N	SER 120	B	37.418	-7.741	-11.190
1501	CA	SER 120	B	38.384	-7.188	-12.116
1502	C	SER 120	B	38.552	-5.769	-11.587
1503	O	SER 120	B	38.771	-5.557	-10.396
1504	CB	SER 120	B	37.765	-7.322	-13.505
1505	OG	SER 120	B	37.361	-6.068	-14.048
1506	N	SER 121	B	38.475	-4.737	-12.351
1507	CA	SER 121	B	37.795	-3.641	-11.746
1508	C	SER 121	B	36.964	-3.170	-12.910
1509	O	SER 121	B	37.218	-2.073	-13.421
1510	CB	SER 121	B	38.808	-2.596	-11.297
1511	OG	SER 121	B	39.930	-3.318	-10.767
1512	N	ASN 122	B	35.961	-3.865	-13.437
1513	CA	ASN 122	B	35.251	-3.163	-14.482
1514	C	ASN 122	B	34.567	-1.986	-13.818
1515	O	ASN 122	B	33.919	-2.205	-12.820
1516	CB	ASN 122	B	34.175	-4.002	-15.124
1517	CG	ASN 122	B	34.720	-4.905	-16.200
1518	OD1	ASN 122	B	35.807	-5.455	-16.161
1519	ND2	ASN 122	B	33.842	-5.095	-17.193
1520	N	VAL 123	B	34.623	-0.738	-14.229
1521	CA	VAL 123	B	33.707	0.247	-13.704
1522	C	VAL 123	B	32.323	0.092	-14.263
1523	O	VAL 123	B	32.121	0.390	-15.418
1524	CB	VAL 123	B	34.253	1.583	-14.033
1525	CG1	VAL 123	B	33.293	2.705	-13.768
1526	CG2	VAL 123	B	35.490	1.719	-13.184
1527	N	ILE 124	B	31.270	-0.356	-13.598
1528	CA	ILE 124	B	29.966	-0.307	-14.250
1529	C	ILE 124	B	28.844	0.618	-13.776
1530	O	ILE 124	B	28.948	1.374	-12.826

57/98
Fig. 15.35

Atom	Residue	Chain	X	Y	Z	
1531	CB	ILE 124	B	29.448	-1.693	-14.301
1532	CG1	ILE 124	B	29.346	-2.339	-12.994
1533	CG2	ILE 124	B	30.460	-2.470	-15.084
1534	CD1	ILE 124	B	28.941	-3.771	-13.335
1535	N	LEU 125	B	27.672	0.649	-14.394
1536	CA	LEU 125	B	26.633	1.575	-13.951
1537	C	LEU 125	B	25.470	0.617	-13.807
1538	O	LEU 125	B	24.992	0.201	-14.856
1539	CB	LEU 125	B	26.338	2.598	-15.029
1540	CG	LEU 125	B	25.522	3.838	-14.719
1541	CD1	LEU 125	B	26.188	4.811	-13.745
1542	CD2	LEU 125	B	25.371	4.560	-16.049
1543	N	LYS 126	B	25.012	0.237	-12.583
1544	CA	LYS 126	B	23.788	-0.531	-12.459
1545	C	LYS 126	B	22.674	0.399	-12.077
1546	O	LYS 126	B	22.853	1.502	-11.576
1547	CB	LYS 126	B	23.882	-1.627	-11.411
1548	CG	LYS 126	B	22.887	-2.671	-12.000
1549	CD	LYS 126	B	22.624	-3.980	-11.185
1550	CE	LYS 126	B	21.336	-4.756	-11.620
1551	NZ	LYS 126	B	20.100	-3.953	-11.514
1552	N	LYS 127	B	21.457	-0.039	-12.325
1553	CA	LYS 127	B	20.275	0.766	-12.115
1554	C	LYS 127	B	19.538	-0.097	-11.161
1555	O	LYS 127	B	19.442	-1.295	-11.349
1556	CB	LYS 127	B	19.357	0.902	-13.313
1557	CG	LYS 127	B	18.494	2.130	-13.043
1558	CD	LYS 127	B	17.595	2.495	-14.203
1559	CE	LYS 127	B	16.759	1.303	-14.621
1560	NZ	LYS 127	B	15.623	1.849	-15.326
1561	N	TYR 128	B	19.018	0.491	-10.119
1562	CA	TYR 128	B	18.374	-0.282	-9.120
1563	C	TYR 128	B	17.039	0.270	-9.294
1564	O	TYR 128	B	16.919	1.484	-9.458
1565	CB	TYR 128	B	18.944	0.051	-7.765
1566	CG	TYR 128	B	20.187	-0.723	-7.613
1567	CD1	TYR 128	B	21.334	-0.090	-7.202
1568	CD2	TYR 128	B	20.189	-2.046	-7.962
1569	CE1	TYR 128	B	22.524	-0.785	-7.189
1570	CE2	TYR 128	B	21.354	-2.752	-7.957
1571	CZ	TYR 128	B	22.522	-2.118	-7.590
1572	OH	TYR 128	B	23.726	-2.834	-7.651
1573	N	ARG 129	B	15.995	-0.539	-9.273
1574	CA	ARG 129	B	14.738	0.093	-9.471
1575	C	ARG 129	B	14.148	0.079	-8.127

58/98
Fig. 15.36

Atom	Residue	Chain	X	Y	Z	
1576	O	ARG 129	B	14.539	-0.696	-7.272
1577	CB	ARG 129	B	13.872	-0.699	-10.389
1578	CG	ARG 129	B	14.448	-1.877	-11.194
1579	CD	ARG 129	B	13.392	-2.283	-12.257
1580	NE	ARG 129	B	12.867	-1.068	-12.880
1581	CZ	ARG 129	B	13.523	-0.465	-13.872
1582	NH1	ARG 129	B	14.600	-1.048	-14.514
1583	NH2	ARG 129	B	13.110	0.783	-14.224
1584	N	ASN 130	B	13.187	0.959	-7.931
1585	CA	ASN 130	B	12.337	0.930	-6.773
1586	C	ASN 130	B	13.101	1.191	-5.580
1587	O	ASN 130	B	12.956	0.581	-4.555
1588	CB	ASN 130	B	11.644	-0.406	-6.596
1589	CG	ASN 130	B	10.513	-0.422	-7.589
1590	OD1	ASN 130	B	9.991	0.615	-7.968
1591	ND2	ASN 130	B	10.106	-1.571	-8.113
1592	N	MET 131	B	13.978	2.163	-5.682
1593	CA	MET 131	B	14.779	2.497	-4.552
1594	C	MET 131	B	14.403	3.805	-3.898
1595	O	MET 131	B	14.888	4.135	-2.829
1596	CB	MET 131	B	16.193	2.489	-5.056
1597	CG	MET 131	B	16.596	1.063	-5.233
1598	SD	MET 131	B	17.774	0.746	-3.920
1599	CE	MET 131	B	17.131	-0.880	-3.651
1600	N	VAL 132	B	13.540	4.602	-4.492
1601	CA	VAL 132	B	13.272	5.952	-4.016
1602	C	VAL 132	B	11.778	6.099	-3.895
1603	O	VAL 132	B	11.075	5.709	-4.791
1604	CB	VAL 132	B	13.804	6.928	-5.035
1605	CG1	VAL 132	B	13.726	8.358	-4.584
1606	CG2	VAL 132	B	15.252	6.549	-5.285
1607	N	VAL 133	B	11.136	6.617	-2.884
1608	CA	VAL 133	B	9.714	6.827	-2.893
1609	C	VAL 133	B	9.534	8.146	-3.591
1610	O	VAL 133	B	10.322	9.039	-3.333
1611	CB	VAL 133	B	9.257	6.869	-1.443
1612	CG1	VAL 133	B	8.030	7.727	-1.166
1613	CG2	VAL 133	B	9.043	5.425	-1.098
1614	N	ARG 134	B	8.526	8.283	-4.462
1615	CA	ARG 134	B	8.074	9.561	-4.979
1616	C	ARG 134	B	6.805	9.998	-4.253
1617	O	ARG 134	B	6.613	11.145	-3.844
1618	CB	ARG 134	B	7.767	9.438	-6.471
1619	CG	ARG 134	B	8.655	10.170	-7.503
1620	CD	ARG 134	B	7.931	10.074	-8.884

59/98

Fig. 15.37

Atom	Residue	Chain	X	Y	Z	
1621	NE	ARG 134	B	8.585	10.702	-10.068
1622	CZ	ARG 134	B	9.074	11.973	-10.169
1623	NH1	ARG 134	B	9.514	12.419	-11.395
1624	NH2	ARG 134	B	9.202	12.813	-9.089
1625	N	ALA 135	B	5.839	9.108	-4.056
1626	CA	ALA 135	B	4.734	9.536	-3.247
1627	C	ALA 135	B	4.163	8.375	-2.497
1628	O	ALA 135	B	4.422	7.228	-2.831
1629	CB	ALA 135	B	3.713	10.178	-4.165
1630	N	CYS 136	B	3.370	8.691	-1.459
1631	CA	CYS 136	B	2.737	7.740	-0.529
1632	C	CYS 136	B	1.234	7.935	-0.411
1633	O	CYS 136	B	0.778	9.062	-0.565
1634	CB	CYS 136	B	3.124	7.893	0.896
1635	SG	CYS 136	B	4.770	8.538	1.134
1636	N	GLY 137	B	0.452	6.910	-0.132
1637	CA	GLY 137	B	-0.882	7.202	0.246
1638	C	GLY 137	B	-1.562	5.941	0.556
1639	O	GLY 137	B	-0.925	4.911	0.422
1640	N	CYS 138	B	-2.836	5.929	0.957
1641	CA	CYS 138	B	-3.377	4.818	1.746
1642	C	CYS 138	B	-3.790	3.683	0.897
1643	O	CYS 138	B	-4.502	4.036	-0.016
1644	CB	CYS 138	B	-4.575	5.261	2.510
1645	SG	CYS 138	B	-4.204	6.910	3.144
1646	N	HIS 139	B	-3.491	2.408	1.017
1647	CA	HIS 139	B	-4.209	1.485	0.173
1648	C	HIS 139	B	-4.475	0.369	1.139
1649	O	HIS 139	B	-3.753	0.227	2.148
1650	CB	HIS 139	B	-3.407	0.969	-1.034
1651	CG	HIS 139	B	-2.881	2.085	-1.888
1652	ND1	HIS 139	B	-2.348	3.285	-1.424
1653	CD2	HIS 139	B	-3.320	2.198	-3.200
1654	CE1	HIS 139	B	-2.505	4.047	-2.523
1655	NE2	HIS 139	B	-3.077	3.480	-3.620
1656	OT	HIS 139	B	-5.079	-0.465	0.767

60/98
Fig. 16\1

Atom	Residue	Chain	X	Y	Z	δ	
1	CB	GLN 36	A	34.688	54.268	11.979	0.80
2	CG	GLN 36	A	34.454	55.622	11.327	0.98
3	CD	GLN 36	A	35.339	55.912	10.142	1.07
4	OE1	GLN 36	A	36.525	55.589	10.061	0.98
5	NE2	GLN 36	A	34.724	56.519	9.123	1.05
6	C	GLN 36	A	33.292	52.214	11.583	0.62
7	O	GLN 36	A	33.249	52.268	10.349	0.67
8	N	GLN 36	A	33.249	53.240	13.781	0.82
9	CA	GLN 36	A	33.430	53.511	12.360	0.72
10	N	ALA 37	A	33.283	51.110	12.314	0.65
11	CA	ALA 37	A	33.203	49.784	11.644	0.64
12	CB	ALA 37	A	33.628	48.665	12.576	0.58
13	C	ALA 37	A	31.758	49.525	11.201	0.62
14	O	ALA 37	A	30.825	49.957	11.868	0.67
15	N	CYS 38	A	31.645	48.593	10.285	0.64
16	CA	CYS 38	A	30.373	48.172	9.714	0.53
17	C	CYS 38	A	29.433	47.699	10.797	0.63
18	O	CYS 38	A	29.746	46.761	11.527	0.66
19	CB	CYS 38	A	30.559	47.212	8.582	0.58
20	SG	CYS 38	A	29.056	46.598	7.828	0.63
21	N	LYS 39	A	28.382	48.487	11.010	0.63
22	CA	LYS 39	A	27.261	48.136	11.860	0.50
23	CB	LYS 39	A	27.365	48.642	13.296	0.56
24	CG	LYS 39	A	28.183	49.898	13.549	0.69
25	CD	LYS 39	A	27.844	50.564	14.870	0.62
26	CE	LYS 39	A	26.701	49.872	15.592	0.92
27	NZ	LYS 39	A	26.052	50.747	16.607	0.92
28	C	LYS 39	A	25.926	48.637	11.262	0.50
29	O	LYS 39	A	25.872	49.334	10.242	0.49
30	N	LYS 40	A	24.890	48.324	11.988	0.59
31	CA	LYS 40	A	23.483	48.556	11.659	0.64
32	CB	LYS 40	A	22.606	47.362	12.132	0.61
33	CG	LYS 40	A	21.167	47.575	11.621	0.60
34	CD	LYS 40	A	20.283	46.452	12.085	0.48
35	CE	LYS 40	A	19.989	46.487	13.565	0.60
36	NZ	LYS 40	A	19.214	45.258	13.873	0.73
37	C	LYS 40	A	23.019	49.744	12.504	0.63
38	O	LYS 40	A	23.463	49.862	13.659	0.61
39	N	HIS 41	A	22.524	50.763	11.826	0.65
40	CA	HIS 41	A	22.150	51.977	12.590	0.58
41	CB	HIS 41	A	22.881	53.195	12.011	0.57
42	CG	HIS 41	A	24.359	52.907	11.899	0.60
43	CD2	HIS 41	A	25.094	52.514	10.838	0.63
44	ND1	HIS 41	A	25.270	53.346	12.835	0.61
45	CE1	HIS 41	A	26.482	53.207	12.338	0.63

61/98
Fig. 16|2

Atom	Residue	Chain	X	Y	Z	δ	
46	NE2	HIS 41	A	26.411	52.515	11.217	0.67
47	C	HIS 41	A	20.645	52.146	12.505	0.58
48	O	HIS 41	A	19.994	51.655	11.564	0.60
49	N	GLU 42	A	20.173	53.007	13.373	0.63
50	CA	GLU 42	A	18.747	53.323	13.460	0.65
51	CB	GLU 42	A	18.389	53.683	14.894	0.67
52	CG	GLU 42	A	17.998	52.443	15.724	0.61
53	CD	GLU 42	A	17.770	52.760	17.171	0.80
54	OE1	GLU 42	A	18.180	52.068	18.082	0.89
55	OE2	GLU 42	A	17.221	53.875	17.309	0.92
56	C	GLU 42	A	18.296	54.397	12.507	0.63
57	O	GLU 42	A	19.083	55.313	12.261	0.68
58	N	LEU 43	A	17.163	54.191	11.858	0.57
59	CA	LEU 43	A	16.444	55.149	11.012	0.57
60	CB	LEU 43	A	16.827	54.887	9.561	0.56
61	CG	LEU 43	A	16.795	55.850	8.455	0.65
62	CD1	LEU 43	A	16.623	55.408	7.043	0.59
63	CD2	LEU 43	A	16.638	57.296	8.715	0.64
64	C	LEU 43	A	14.940	54.753	11.210	0.63
65	O	LEU 43	A	14.568	53.682	10.710	0.62
66	N	TYR 44	A	14.119	55.703	11.580	0.58
67	CA	TYR 44	A	12.646	55.504	11.590	0.54
68	CB	TYR 44	A	12.117	56.253	12.845	0.52
69	CG	TYR 44	A	10.648	55.954	13.022	0.63
70	CD1	TYR 44	A	9.663	56.853	12.620	0.50
71	CE1	TYR 44	A	8.318	56.490	12.742	0.58
72	CD2	TYR 44	A	10.267	54.766	13.634	0.52
73	CE2	TYR 44	A	8.957	54.379	13.694	0.56
74	CZ	TYR 44	A	7.976	55.253	13.235	0.69
75	OH	TYR 44	A	6.668	54.894	13.434	0.81
76	C	TYR 44	A	12.114	56.295	10.395	0.51
77	O	TYR 44	A	12.538	57.447	10.316	0.63
78	N	VAL 45	A	11.336	55.713	9.534	0.53
79	CA	VAL 45	A	10.722	56.389	8.368	0.53
80	CB	VAL 45	A	10.910	55.465	7.148	0.56
81	CG1	VAL 45	A	10.355	56.000	5.844	0.47
82	CG2	VAL 45	A	12.346	54.969	7.035	0.52
83	C	VAL 45	A	9.220	56.590	8.673	0.55
84	O	VAL 45	A	8.585	55.655	9.193	0.53
85	N	SER 46	A	8.851	57.845	8.765	0.59
86	CA	SER 46	A	7.518	58.414	8.858	0.52
87	CB	SER 46	A	7.518	59.907	9.287	0.46
88	OG	SER 46	A	6.397	60.090	10.135	0.69
89	C	SER 46	A	6.883	58.405	7.482	0.52
90	O	SER 46	A	7.484	58.947	6.553	0.56

62/98

Fig. 16\3

Atom	Residue	Chain	X	Y	Z	δ	
91	N	PHE 47	A	5.713	57.819	7.380	0.59
92	CA	PHE 47	A	5.009	57.837	6.069	0.57
93	CB	PHE 47	A	3.844	56.851	6.088	0.53
94	CG	PHE 47	A	4.154	55.395	6.174	0.55
95	CD1	PHE 47	A	5.279	54.859	5.517	0.45
96	CD2	PHE 47	A	3.287	54.509	6.818	0.54
97	CE1	PHE 47	A	5.580	53.514	5.633	0.71
98	CE2	PHE 47	A	3.618	53.164	7.010	0.56
99	CZ	PHE 47	A	4.739	52.644	6.313	0.60
100	C	PHE 47	A	4.737	59.281	5.648	0.49
101	O	PHE 47	A	4.748	59.578	4.426	0.61
102	N	ARG 48	A	4.422	60.187	6.541	0.51
103	CA	ARG 48	A	4.356	61.641	6.314	0.57
104	CB	ARG 48	A	4.183	62.463	7.564	0.49
105	CG	ARG 48	A	3.277	62.040	8.662	0.74
106	CD	ARG 48	A	2.096	62.917	8.844	0.83
107	NE	ARG 48	A	2.148	63.816	9.964	0.71
108	CZ	ARG 48	A	1.255	64.003	10.923	0.66
109	NH1	ARG 48	A	-0.043	63.685	10.831	0.68
110	NH2	ARG 48	A	1.739	64.330	12.123	0.61
111	C	ARG 48	A	5.516	62.216	5.517	0.64
112	O	ARG 48	A	5.342	62.761	4.397	0.69
113	N	ASP 49	A	6.737	61.953	5.977	0.60
114	CA	ASP 49	A	7.962	62.298	5.274	0.57
115	CB	ASP 49	A	9.205	61.850	6.047	0.68
116	CG	ASP 49	A	9.188	62.342	7.481	0.80
117	OD1	ASP 49	A	10.062	61.939	8.272	0.89
118	OD2	ASP 49	A	8.235	63.099	7.801	0.83
119	C	ASP 49	A	7.998	61.913	3.814	0.66
120	O	ASP 49	A	8.811	62.490	3.062	0.72
121	N	LEU 50	A	7.456	60.746	3.465	0.70
122	CA	LEU 50	A	7.487	60.243	2.086	0.60
123	CB	LEU 50	A	7.446	58.703	2.187	0.62
124	CG	LEU 50	A	8.580	58.098	3.022	0.69
125	CD1	LEU 50	A	8.786	56.662	2.544	0.66
126	CD2	LEU 50	A	9.836	58.911	2.667	0.66
127	C	LEU 50	A	6.275	60.748	1.275	0.62
128	O	LEU 50	A	6.080	60.203	0.167	0.69
129	N	GLY 51	A	5.315	61.265	2.005	0.62
130	CA	GLY 51	A	3.950	61.540	1.563	0.59
131	C	GLY 51	A	3.215	60.231	1.303	0.68
132	O	GLY 51	A	2.960	59.916	0.130	0.73
133	N	TRP 52	A	3.076	59.385	2.325	0.60
134	CA	TRP 52	A	2.385	58.104	2.063	0.63
135	CB	TRP 52	A	3.281	56.916	2.022	0.76

63/98
*Fig. 16*4

Atom	Residuc	Chain	X	Y	Z	δ	
136	CG	TRP 52	A	4.282	56.698	0.951	0.80
137	CD2	TRP 52	A	5.364	55.744	1.004	0.74
138	CE2	TRP 52	A	5.991	55.750	-0.263	0.92
139	CE3	TRP 52	A	5.811	54.835	1.963	1.01
140	CD1	TRP 52	A	4.256	57.163	-0.333	0.82
141	NE1	TRP 52	A	5.315	56.649	-1.050	0.73
142	CZ2	TRP 52	A	7.038	54.881	-0.576	0.75
143	CZ3	TRP 52	A	6.886	54.012	1.684	0.68
144	CH2	TRP 52	A	7.495	54.041	0.428	0.80
145	C	TRP 52	A	1.218	57.994	3.046	0.72
146	O	TRP 52	A	0.425	57.045	2.965	0.78
147	N	GLN 53	A	1.009	59.076	3.769	0.64
148	CA	GLN 53	A	0.000	59.125	4.822	0.74
149	CB	GLN 53	A	0.235	60.290	5.798	0.68
150	CG	GLN 53	A	0.108	61.647	5.138	0.56
151	CD	GLN 53	A	1.210	62.152	4.268	0.78
152	OE1	GLN 53	A	1.734	61.501	3.363	0.75
153	NE2	GLN 53	A	1.476	63.465	4.447	0.80
154	C	GLN 53	A	-1.419	59.100	4.280	0.74
155	O	GLN 53	A	-2.376	59.017	5.080	0.83
156	N	ASP 54	A	-1.613	59.500	3.040	0.77
157	CA	ASP 54	A	-2.945	59.676	2.432	0.76
158	CB	ASP 54	A	-2.747	60.190	1.004	0.92
159	CG	ASP 54	A	-2.418	61.667	0.920	0.92
160	OD1	ASP 54	A	-2.672	62.439	1.863	0.96
161	OD2	ASP 54	A	-2.074	62.062	-0.217	0.75
162	C	ASP 54	A	-3.808	58.408	2.491	0.80
163	O	ASP 54	A	-4.988	58.426	2.898	0.82
164	N	TRP 55	A	-3.265	57.334	1.953	0.72
165	CA	TRP 55	A	-3.905	56.052	1.723	0.78
166	CB	TRP 55	A	-3.527	55.614	0.293	0.69
167	CG	TRP 55	A	-3.900	56.679	-0.691	0.80
168	CD2	TRP 55	A	-4.955	57.641	-0.465	0.76
169	CE2	TRP 55	A	-4.935	58.522	-1.569	0.85
170	CE3	TRP 55	A	-6.064	57.611	0.374	0.87
171	CD1	TRP 55	A	-3.230	57.113	-1.796	0.74
172	NE1	TRP 55	A	-3.835	58.229	-2.337	0.76
173	CZ2	TRP 55	A	-5.913	59.493	-1.745	0.78
174	CZ3	TRP 55	A	-7.074	58.520	0.151	0.88
175	CH2	TRP 55	A	-7.012	59.424	-0.905	0.82
176	C	TRP 55	A	-3.640	54.977	2.770	0.84
177	O	TRP 55	A	-4.364	53.952	2.758	0.77
178	N	ILE 56	A	-2.653	55.170	3.642	0.83
179	CA	ILE 56	A	-2.272	54.240	4.693	0.78
180	CB	ILE 56	A	-0.750	54.279	5.076	0.77

^{64/98}
Fig. 16\5

Atom	Residuc	Chain	X	Y	Z	δ	
181	CG2	ILE 56	A	-0.539	54.069	6.606	0.71
182	CG1	ILE 56	A	-0.007	53.147	4.316	0.82
183	CD	ILE 56	A	1.044	53.678	3.298	0.90
184	C	ILE 56	A	-3.122	54.432	5.944	0.74
185	O	ILE 56	A	-3.277	55.575	6.388	0.80
186	N	ILE 57	A	-3.481	53.314	6.567	0.68
187	CA	ILE 57	A	-4.235	53.305	7.822	0.64
188	CB	ILE 57	A	-5.462	52.319	7.725	0.65
189	CG2	ILE 57	A	-6.003	51.940	9.109	0.55
190	CG1	ILE 57	A	-6.532	52.829	6.736	0.64
191	CD	ILE 57	A	-7.736	51.860	6.540	0.62
192	C	ILE 57	A	-3.350	52.989	9.020	0.60
193	O	ILE 57	A	-3.634	53.435	10.145	0.62
194	N	ALA 58	A	-2.432	52.035	8.834	0.58
195	CA	ALA 58	A	-1.568	51.588	9.967	0.49
196	CB	ALA 58	A	-2.381	50.553	10.789	0.55
197	C	ALA 58	A	-0.372	50.826	9.376	0.46
198	O	ALA 58	A	-0.718	50.311	8.302	0.49
199	N	PRO 59	A	0.849	51.149	9.744	0.53
200	CD	PRO 59	A	2.056	50.542	9.192	0.46
201	CA	PRO 59	A	1.271	52.004	10.870	0.45
202	CB	PRO 59	A	2.504	51.340	11.429	0.47
203	CG	PRO 59	A	3.139	50.681	10.217	0.43
204	C	PRO 59	A	1.496	53.434	10.384	0.53
205	O	PRO 59	A	1.157	53.822	9.239	0.55
206	N	GLU 60	A	2.009	54.297	11.255	0.49
207	CA	GLU 60	A	2.315	55.678	10.776	0.58
208	CB	GLU 60	A	1.953	56.668	11.906	0.64
209	CG	GLU 60	A	0.536	57.221	11.886	0.77
210	CD	GLU 60	A	-0.327	57.374	13.082	0.79
211	OE1	GLU 60	A	-1.419	56.801	13.170	0.84
212	OE2	GLU 60	A	0.122	58.112	13.997	0.61
213	C	GLU 60	A	3.778	55.806	10.374	0.53
214	O	GLU 60	A	4.180	56.692	9.605	0.52
215	N	GLY 61	A	4.521	54.712	10.496	0.52
216	CA	GLY 61	A	5.866	54.512	9.965	0.54
217	C	GLY 61	A	6.549	53.304	10.585	0.58
218	O	GLY 61	A	5.879	52.548	11.298	0.55
219	N	TYR 62	A	7.807	53.047	10.231	0.55
220	CA	TYR 62	A	8.462	51.838	10.845	0.54
221	CB	TYR 62	A	8.178	50.656	9.853	0.38
222	CG	TYR 62	A	8.980	50.842	8.558	0.47
223	CD1	TYR 62	A	8.538	51.787	7.649	0.47
224	CE1	TYR 62	A	9.247	52.151	6.499	0.51
225	CD2	TYR 62	A	10.091	50.092	8.201	0.45

65/98
Fig. 16

Atom	Residue	Chain	X	Y	Z	δ	
226	CE2	TYR 62	A	10.662	50.225	6.912	0.35
227	CZ	TYR 62	A	10.294	51.305	6.115	0.51
228	OH	TYR 62	A	10.785	51.429	4.839	0.65
229	C	TYR 62	A	9.960	52.092	10.973	0.59
230	O	TYR 62	A	10.580	52.978	10.330	0.55
231	N	ALA 63	A	10.590	51.236	11.731	0.54
232	CA	ALA 63	A	12.067	51.311	11.984	0.49
233	CB	ALA 63	A	12.243	50.507	13.282	0.58
234	C	ALA 63	A	12.734	50.563	10.845	0.51
235	O	ALA 63	A	12.508	49.358	10.665	0.58
236	N	ALA 64	A	13.289	51.297	9.895	0.54
237	CA	ALA 64	A	13.878	50.657	8.735	0.53
238	CB	ALA 64	A	13.779	51.617	7.550	0.53
239	C	ALA 64	A	15.319	50.155	8.972	0.52
240	O	ALA 64	A	15.797	49.399	8.091	0.54
241	N	TYR 65	A	16.092	50.933	9.671	0.51
242	CA	TYR 65	A	17.512	50.804	9.978	0.54
243	CB	TYR 65	A	17.944	49.390	10.438	0.47
244	CG	TYR 65	A	17.297	49.063	11.777	0.47
245	CD1	TYR 65	A	17.763	49.564	12.974	0.44
246	CE1	TYR 65	A	17.164	49.219	14.190	0.47
247	CD2	TYR 65	A	15.978	48.600	11.756	0.61
248	CE2	TYR 65	A	15.250	48.484	12.922	0.44
249	CZ	TYR 65	A	15.849	48.733	14.139	0.67
250	OH	TYR 65	A	15.077	48.514	15.253	0.70
251	C	TYR 65	A	18.332	51.188	8.752	0.51
252	O	TYR 65	A	17.714	51.205	7.664	0.49
253	N	TYR 66	A	19.638	51.001	8.928	0.52
254	CA	TYR 66	A	20.521	51.078	7.729	0.50
255	CB	TYR 66	A	20.564	52.484	7.126	0.60
256	CG	TYR 66	A	21.276	53.507	7.983	0.59
257	CD1	TYR 66	A	20.563	54.255	8.917	0.59
258	CE1	TYR 66	A	21.155	55.260	9.659	0.53
259	CD2	TYR 66	A	22.592	53.893	7.721	0.53
260	CE2	TYR 66	A	23.149	55.001	8.335	0.47
261	CZ	TYR 66	A	22.511	55.532	9.448	0.58
262	OH	TYR 66	A	23.095	56.600	10.066	0.70
263	C	TYR 66	A	21.939	50.673	8.166	0.51
264	O	TYR 66	A	22.223	50.678	9.351	0.52
265	N	CYS 67	A	22.764	50.444	7.173	0.58
266	CA	CYS 67	A	24.127	49.897	7.372	0.57
267	C	CYS 67	A	25.134	50.997	7.066	0.52
268	O	CYS 67	A	25.035	51.589	5.968	0.48
269	CB	CYS 67	A	24.270	48.712	6.413	0.54
270	SG	CYS 67	A	23.285	47.244	6.831	0.54

66/98

Fig. 16\7

Atom	Residue	Chain	X	Y	Z	δ	
271	N	GLU 68	A	26.124	51.140	7.930	0.56
272	CA	GLU 68	A	27.251	52.049	7.617	0.47
273	CB	GLU 68	A	26.880	53.495	7.993	0.68
274	CG	GLU 68	A	27.575	54.628	7.264	0.53
275	CD	GLU 68	A	27.058	56.019	7.510	0.67
276	OE1	GLU 68	A	27.235	56.925	6.720	0.64
277	OE2	GLU 68	A	26.696	56.240	8.687	0.63
278	C	GLU 68	A	28.467	51.725	8.485	0.43
279	O	GLU 68	A	28.311	51.441	9.663	0.42
280	N	GLY 69	A	29.646	52.013	7.922	0.48
281	CA	GLY 69	A	30.920	51.860	8.669	0.44
282	C	GLY 69	A	31.899	51.218	7.670	0.51
283	O	GLY 69	A	31.525	50.790	6.563	0.53
284	N	GLU 70	A	33.165	51.150	8.048	0.57
285	CA	GLU 70	A	34.166	50.669	7.116	0.63
286	CB	GLU 70	A	35.428	51.440	6.924	0.74
287	CG	GLU 70	A	36.419	51.567	8.062	0.77
288	CD	GLU 70	A	37.808	51.898	7.554	1.02
289	OE1	GLU 70	A	38.022	52.115	6.372	0.84
290	OE2	GLU 70	A	38.665	51.734	8.441	0.95
291	C	GLU 70	A	34.384	49.172	7.133	0.44
292	O	GLU 70	A	34.224	48.564	8.186	0.42
293	N	CYS 71	A	34.793	48.707	5.962	0.50
294	CA	CYS 71	A	35.136	47.281	5.805	0.62
295	C	CYS 71	A	36.652	47.128	5.637	0.54
296	O	CYS 71	A	37.173	47.188	4.547	0.51
297	CB	CYS 71	A	34.283	46.609	4.771	0.62
298	SG	CYS 71	A	32.525	46.435	5.196	0.57
299	N	ALA 72	A	37.345	47.031	6.745	0.50
300	CA	ALA 72	A	38.781	47.033	6.862	0.61
301	CB	ALA 72	A	39.243	48.414	7.425	0.61
302	C	ALA 72	A	39.174	45.969	7.903	0.62
303	O	ALA 72	A	38.444	45.747	8.880	0.59
304	N	PHE 73	A	40.472	45.646	7.785	0.59
305	CA	PHE 73	A	41.088	44.789	8.805	0.46
306	CB	PHE 73	A	42.425	44.197	8.472	0.54
307	CG	PHE 73	A	42.433	43.361	7.198	0.45
308	CD1	PHE 73	A	41.659	42.210	7.129	0.43
309	CD2	PHE 73	A	43.175	43.752	6.117	0.45
310	CE1	PHE 73	A	41.516	41.496	5.935	0.34
311	CE2	PHE 73	A	43.202	42.987	4.959	0.37
312	CZ	PHE 73	A	42.340	41.869	4.895	0.33
313	C	PHE 73	A	40.895	45.531	10.083	0.52
314	O	PHE 73	A	41.336	46.672	9.925	0.54
315	N	PRO 74	A	40.737	44.872	11.197	0.57

67/98

Fig. 16\8

Atom	Residue		Chain	X	Y	Z	δ
316	CD	PRO 74	A	40.751	45.550	12.495	0.57
317	CA	PRO 74	A	40.171	43.560	11.370	0.56
318	CB	PRO 74	A	40.235	43.204	12.832	0.52
319	CG	PRO 74	A	40.362	44.522	13.519	0.54
320	C	PRO 74	A	38.776	43.425	10.805	0.66
321	O	PRO 74	A	37.919	44.259	11.112	0.74
322	N	LEU 75	A	38.607	42.334	10.090	0.58
323	CA	LEU 75	A	37.302	41.889	9.627	0.61
324	CB	LEU 75	A	37.400	41.397	8.209	0.63
325	CG	LEU 75	A	37.038	42.169	7.006	0.60
326	CD1	LEU 75	A	36.742	43.625	7.025	0.57
327	CD2	LEU 75	A	37.596	41.719	5.703	0.49
328	C	LEU 75	A	36.805	40.946	10.726	0.68
329	O	LEU 75	A	36.983	39.722	10.655	0.72
330	N	ASN 76	A	36.381	41.589	11.802	0.68
331	CA	ASN 76	A	35.736	40.919	12.939	0.66
332	CB	ASN 76	A	35.320	41.901	14.028	0.62
333	CG	ASN 76	A	36.481	42.611	14.697	0.83
334	OD1	ASN 76	A	37.623	42.110	14.677	0.97
335	ND2	ASN 76	A	36.246	43.740	15.370	0.81
336	C	ASN 76	A	34.658	39.972	12.449	0.67
337	O	ASN 76	A	34.194	39.990	11.281	0.73
338	N	SER 77	A	34.415	38.958	13.260	0.73
339	CA	SER 77	A	33.432	37.909	12.886	0.74
340	CB	SER 77	A	33.434	36.747	13.894	0.54
341	OG	SER 77	A	33.117	35.573	13.128	0.96
342	C	SER 77	A	32.033	38.539	12.767	0.62
343	O	SER 77	A	31.287	38.245	11.835	0.66
344	N	TYR 78	A	31.705	39.321	13.767	0.63
345	CA	TYR 78	A	30.579	40.220	13.850	0.75
346	CB	TYR 78	A	30.615	41.057	15.137	0.63
347	CG	TYR 78	A	30.922	42.527	14.976	0.91
348	CD1	TYR 78	A	32.115	43.118	15.447	0.81
349	CE1	TYR 78	A	32.383	44.471	15.228	0.88
350	CD2	TYR 78	A	29.929	43.382	14.494	0.98
351	CE2	TYR 78	A	30.178	44.734	14.265	0.99
352	CZ	TYR 78	A	31.402	45.277	14.640	0.94
353	OH	TYR 78	A	31.541	46.629	14.483	0.97
354	C	TYR 78	A	30.340	41.045	12.592	0.78
355	O	TYR 78	A	29.244	41.627	12.487	0.79
356	N	MET 79	A	31.322	41.194	11.719	0.74
357	CA	MET 79	A	31.228	41.956	10.475	0.62
358	CB	MET 79	A	32.506	42.736	10.191	0.68
359	CG	MET 79	A	32.600	43.844	11.207	0.51
360	SD	MET 79	A	34.243	44.508	11.149	0.67

68/98

Fig. 16|9

Atom	Residue		Chain	X	Y	Z	δ
361	CE	MET 79	A	34.390	45.358	9.625	0.53
362	C	MET 79	A	30.807	41.104	9.309	0.58
363	O	MET 79	A	30.601	41.571	8.186	0.61
364	N	ASN 80	A	30.496	39.834	9.575	0.55
365	CA	ASN 80	A	30.017	39.029	8.431	0.54
366	CB	ASN 80	A	28.501	38.847	8.469	0.86
367	CG	ASN 80	A	27.937	38.129	7.244	1.05
368	OD1	ASN 80	A	28.023	38.584	6.079	0.95
369	ND2	ASN 80	A	27.180	37.052	7.518	0.86
370	C	ASN 80	A	30.573	39.451	7.091	0.57
371	O	ASN 80	A	29.821	39.382	6.054	0.67
372	N	ALA 81	A	31.852	39.123	6.890	0.60
373	CA	ALA 81	A	32.527	39.331	5.577	0.52
374	CB	ALA 81	A	34.012	39.611	5.894	0.53
375	C	ALA 81	A	32.276	38.077	4.802	0.53
376	O	ALA 81	A	32.376	37.027	5.445	0.60
377	N	THR 82	A	32.057	38.098	3.500	0.54
378	CA	THR 82	A	32.162	36.889	2.680	0.44
379	CB	THR 82	A	31.426	37.170	1.326	0.45
380	OG1	THR 82	A	32.057	38.409	0.821	0.49
381	CG2	THR 82	A	29.937	37.606	1.601	0.58
382	C	THR 82	A	33.652	36.578	2.443	0.67
383	O	THR 82	A	34.576	37.324	2.852	0.55
384	N	ASN 83	A	33.946	35.396	1.913	0.61
385	CA	ASN 83	A	35.352	35.123	1.529	0.56
386	CB	ASN 83	A	35.532	33.739	0.901	0.53
387	CG	ASN 83	A	35.404	32.602	1.900	0.48
388	OD1	ASN 83	A	35.542	32.820	3.108	0.55
389	ND2	ASN 83	A	34.880	31.483	1.410	0.55
390	C	ASN 83	A	35.756	36.228	0.535	0.55
391	O	ASN 83	A	36.971	36.442	0.428	0.54
392	N	HIS 84	A	35.021	36.349	-0.569	0.55
393	CA	HIS 84	A	35.225	37.440	-1.518	0.41
394	CB	HIS 84	A	34.045	37.721	-2.462	0.58
395	CG	HIS 84	A	34.351	38.672	-3.572	0.40
396	CD2	HIS 84	A	34.775	38.378	-4.809	0.29
397	ND1	HIS 84	A	34.471	40.064	-3.426	0.22
398	CE1	HIS 84	A	34.778	40.549	-4.612	0.18
399	NE2	HIS 84	A	34.912	39.565	-5.513	0.31
400	C	HIS 84	A	35.639	38.754	-0.878	0.47
401	O	HIS 84	A	36.412	39.493	-1.484	0.50
402	N	ALA 85	A	35.014	39.218	0.190	0.52
403	CA	ALA 85	A	35.340	40.514	0.773	0.53
404	CB	ALA 85	A	34.341	40.834	1.868	0.48
405	C	ALA 85	A	36.756	40.456	1.388	0.57

69/98

Fig. 16\10

Atom	Residue	Chain	X	Y	Z	δ	
406	O	ALA 85	A	37.222	41.564	1.781	0.55
407	N	ILE 86	A	37.011	39.347	2.095	0.50
408	CA	ILE 86	A	38.305	39.171	2.783	0.52
409	CB	ILE 86	A	38.390	37.738	3.425	0.57
410	CG2	ILE 86	A	39.877	37.505	3.912	0.42
411	CG1	ILE 86	A	37.435	37.747	4.652	0.40
412	CD	ILE 86	A	37.270	36.306	5.234	0.49
413	C	ILE 86	A	39.432	39.316	1.751	0.45
414	O	ILE 86	A	40.345	40.113	1.894	0.58
415	N	VAL 87	A	39.232	38.678	0.618	0.49
416	CA	VAL 87	A	40.135	38.769	-0.517	0.42
417	CB	VAL 87	A	39.663	37.786	-1.570	0.43
418	CG1	VAL 87	A	40.259	38.002	-2.950	0.39
419	CG2	VAL 87	A	39.952	36.365	-1.097	0.50
420	C	VAL 87	A	40.276	40.212	-0.984	0.52
421	O	VAL 87	A	41.217	40.442	-1.765	0.57
422	N	GLN 88	A	39.086	40.754	-1.301	0.64
423	CA	GLN 88	A	38.956	42.130	-1.773	0.50
424	CB	GLN 88	A	37.685	42.583	-2.372	0.54
425	CG	GLN 88	A	37.381	44.072	-2.355	0.50
426	CD	GLN 88	A	36.114	44.340	-3.126	0.53
427	OE1	GLN 88	A	35.371	43.417	-3.448	0.63
428	NE2	GLN 88	A	35.759	45.595	-3.183	0.42
429	C	GLN 88	A	39.679	43.085	-0.867	0.42
430	O	GLN 88	A	40.292	44.052	-1.314	0.52
431	N	THR 89	A	39.512	42.927	0.420	0.52
432	CA	THR 89	A	40.198	43.702	1.434	0.51
433	CB	THR 89	A	39.557	43.573	2.830	0.44
434	OG1	THR 89	A	38.120	43.824	2.666	0.58
435	CG2	THR 89	A	40.094	44.473	3.906	0.28
436	C	THR 89	A	41.709	43.490	1.437	0.60
437	O	THR 89	A	42.412	44.454	1.772	0.60
438	N	LEU 90	A	42.218	42.388	0.944	0.60
439	CA	LEU 90	A	43.647	42.052	0.965	0.54
440	CB	LEU 90	A	43.888	40.582	1.267	0.64
441	CG	LEU 90	A	45.291	40.015	1.078	0.62
442	CD1	LEU 90	A	46.152	40.465	2.272	0.49
443	CD2	LEU 90	A	45.199	38.484	1.213	0.62
444	C	LEU 90	A	44.288	42.626	-0.279	0.47
445	O	LEU 90	A	45.284	43.352	-0.185	0.63
446	N	VAL 91	A	43.579	42.594	-1.376	0.43
447	CA	VAL 91	A	43.901	43.242	-2.618	0.40
448	CB	VAL 91	A	42.933	42.872	-3.726	0.35
449	CG1	VAL 91	A	43.378	43.386	-5.070	0.33
450	CG2	VAL 91	A	42.891	41.354	-3.941	0.47

70/98

Fig. 16\11

Atom	Residue	Chain	X	Y	Z	δ	
451	C	VAL 91	A	44.178	44.727	-2.505	0.49
452	O	VAL 91	A	44.650	45.311	-3.477	0.59
453	N	HIS 92	A	43.441	45.341	-1.635	0.51
454	CA	HIS 92	A	43.261	46.741	-1.409	0.56
455	CB	HIS 92	A	41.870	47.044	-0.770	0.55
456	CG	HIS 92	A	41.747	48.505	-0.469	0.52
457	CD2	HIS 92	A	42.030	49.193	0.668	0.36
458	ND1	HIS 92	A	41.469	49.429	-1.483	0.42
459	CE1	HIS 92	A	41.437	50.616	-0.876	0.49
460	NE2	HIS 92	A	41.799	50.525	0.379	0.43
461	C	HIS 92	A	44.394	47.223	-0.492	0.50
462	O	HIS 92	A	45.068	48.210	-0.766	0.62
463	N	PHE 93	A	44.696	46.435	0.494	0.52
464	CA	PHE 93	A	45.853	46.579	1.335	0.62
465	CB	PHE 93	A	45.848	45.644	2.519	0.52
466	CG	PHE 93	A	47.210	45.573	3.167	0.65
467	CD1	PHE 93	A	48.069	44.545	2.795	0.60
468	CD2	PHE 93	A	47.593	46.500	4.135	0.66
469	CE1	PHE 93	A	49.393	44.536	3.237	0.64
470	CE2	PHE 93	A	48.928	46.509	4.580	0.66
471	CZ	PHE 93	A	49.711	45.369	4.295	0.51
472	C	PHE 93	A	47.144	46.550	0.506	0.66
473	O	PHE 93	A	47.943	47.489	0.617	0.73
474	N	ILE 94	A	47.331	45.557	-0.315	0.64
475	CA	ILE 94	A	48.423	45.367	-1.252	0.61
476	CB	ILE 94	A	48.402	43.957	-1.897	0.63
477	CG2	ILE 94	A	49.466	43.645	-2.972	0.63
478	CG1	ILE 94	A	48.404	42.831	-0.814	0.51
479	CD	ILE 94	A	48.147	41.476	-1.602	0.63
480	C	ILE 94	A	48.550	46.477	-2.289	0.70
481	O	ILE 94	A	49.659	46.609	-2.824	0.75
482	N	ASN 95	A	47.465	46.873	-2.923	0.74
483	CA	ASN 95	A	47.356	47.921	-3.913	0.64
484	CB	ASN 95	A	47.834	47.543	-5.288	0.70
485	CG	ASN 95	A	48.029	48.754	-6.197	0.86
486	OD1	ASN 95	A	47.797	49.907	-5.782	0.79
487	ND2	ASN 95	A	48.255	48.499	-7.487	0.87
488	C	ASN 95	A	46.014	48.658	-3.867	0.65
489	O	ASN 95	A	45.144	48.489	-4.746	0.65
490	N	PRO 96	A	46.039	49.731	-3.096	0.64
491	CD	PRO 96	A	47.201	50.067	-2.228	0.68
492	CA	PRO 96	A	44.914	50.589	-2.805	0.65
493	CB	PRO 96	A	45.500	51.713	-1.943	0.69
494	CG	PRO 96	A	46.582	50.992	-1.180	0.72
495	C	PRO 96	A	44.106	51.115	-3.961	0.72

71/98

Fig. 16|12

Atom	Residue	Chain	X	Y	Z	δ	
496	O	PRO 96	A	42.930	51.520	-3.812	0.73
497	N	GLU 97	A	44.662	50.992	-5.140	0.72
498	CA	GLU 97	A	44.135	51.677	-6.338	0.70
499	CB	GLU 97	A	45.301	52.282	-7.121	0.84
500	CG	GLU 97	A	45.685	53.733	-7.069	0.80
501	CD	GLU 97	A	45.275	54.630	-5.952	1.11
502	OE1	GLU 97	A	44.477	55.564	-6.044	1.13
503	OE2	GLU 97	A	46.026	54.515	-4.948	1.11
504	C	GLU 97	A	43.468	50.625	-7.224	0.72
505	O	GLU 97	A	42.759	50.951	-8.192	0.74
506	N	THR 98	A	43.863	49.373	-7.015	0.69
507	CA	THR 98	A	43.305	48.280	-7.830	0.67
508	CB	THR 98	A	44.071	46.915	-7.570	0.65
509	OG1	THR 98	A	45.398	47.345	-7.123	0.77
510	CG2	THR 98	A	44.197	46.102	-8.854	0.78
511	C	THR 98	A	41.817	48.097	-7.527	0.63
512	O	THR 98	A	41.046	47.740	-8.443	0.67
513	N	VAL 99	A	41.488	48.234	-6.249	0.57
514	CA	VAL 99	A	40.083	47.984	-5.846	0.52
515	CB	VAL 99	A	39.921	46.444	-5.871	0.56
516	CG1	VAL 99	A	40.467	45.868	-4.573	0.44
517	CG2	VAL 99	A	38.532	45.932	-6.138	0.62
518	C	VAL 99	A	39.813	48.669	-4.532	0.52
519	O	VAL 99	A	40.679	48.871	-3.666	0.53
520	N	PRO 100	A	38.557	49.141	-4.403	0.53
521	CD	PRO 100	A	37.460	48.923	-5.352	0.54
522	CA	PRO 100	A	38.083	49.661	-3.125	0.49
523	CB	PRO 100	A	36.659	50.149	-3.411	0.50
524	CG	PRO 100	A	36.231	49.470	-4.666	0.47
525	C	PRO 100	A	38.085	48.608	-2.041	0.41
526	O	PRO 100	A	37.913	47.407	-2.250	0.58
527	N	LYS 101	A	37.739	49.029	-0.836	0.48
528	CA	LYS 101	A	37.365	48.136	0.241	0.40
529	CB	LYS 101	A	37.549	48.844	1.590	0.58
530	CG	LYS 101	A	38.756	48.257	2.367	0.69
531	CD	LYS 101	A	39.462	49.214	3.271	0.65
532	CE	LYS 101	A	38.839	50.616	3.297	0.61
533	NZ	LYS 101	A	39.223	51.248	4.589	0.76
534	C	LYS 101	A	35.850	47.825	0.048	0.56
535	O	LYS 101	A	35.301	48.383	-0.922	0.54
536	N	PRO 102	A	35.434	46.638	0.475	0.57
537	CD	PRO 102	A	36.255	45.605	1.164	0.48
538	CA	PRO 102	A	34.017	46.290	0.428	0.59
539	CB	PRO 102	A	33.921	44.912	1.028	0.43
540	CG	PRO 102	A	35.209	44.663	1.736	0.54

72/98

Fig. 16\13

Atom	Residuc	Chain	X	Y	Z	δ	
541	C	PRO 102	A	33.146	47.342	1.076	0.56
542	O	PRO 102	A	33.527	48.083	2.006	0.63
543	N	CYS 103	A	31.861	47.298	0.744	0.58
544	CA	CYS 103	A	30.857	48.177	1.400	0.48
545	CB	CYS 103	A	30.152	49.052	0.384	0.75
546	SG	CYS 103	A	28.673	48.328	-0.344	0.81
547	C	CYS 103	A	29.943	47.435	2.355	0.55
548	O	CYS 103	A	29.809	46.208	2.353	0.53
549	N	CYS 104	A	29.644	48.101	3.461	0.32
550	CA	CYS 104	A	28.717	47.709	4.468	0.48
551	C	CYS 104	A	27.265	47.729	3.963	0.60
552	O	CYS 104	A	26.729	48.760	3.570	0.60
553	CB	CYS 104	A	28.932	48.683	5.594	0.32
554	SG	CYS 104	A	28.146	48.223	7.124	0.54
555	N	ALA 105	A	26.605	46.583	4.024	0.51
556	CA	ALA 105	A	25.412	46.300	3.222	0.51
557	CB	ALA 105	A	25.801	45.889	1.838	0.51
558	C	ALA 105	A	24.633	45.240	4.017	0.51
559	O	ALA 105	A	25.189	44.843	5.068	0.57
560	N	PRO 106	A	23.328	45.294	3.866	0.59
561	CD	PRO 106	A	22.564	46.026	2.822	0.51
562	CA	PRO 106	A	22.416	44.526	4.739	0.59
563	CB	PRO 106	A	21.014	45.021	4.321	0.54
564	CG	PRO 106	A	21.235	46.282	3.559	0.50
565	C	PRO 106	A	22.577	43.053	4.332	0.58
566	O	PRO 106	A	22.779	42.755	3.154	0.54
567	N	THR 107	A	22.616	42.182	5.306	0.60
568	CA	THR 107	A	22.791	40.733	4.913	0.65
569	CB	THR 107	A	23.831	40.113	5.928	0.45
570	OG1	THR 107	A	23.141	40.103	7.208	0.68
571	CG2	THR 107	A	25.153	40.852	6.091	0.56
572	C	THR 107	A	21.417	40.091	5.133	0.68
573	O	THR 107	A	21.037	39.069	4.543	0.68
574	N	GLN 108	A	20.597	40.806	5.902	0.67
575	CA	GLN 108	A	19.208	40.371	6.067	0.64
576	CB	GLN 108	A	19.029	39.316	7.105	0.58
577	CG	GLN 108	A	19.016	39.718	8.547	0.70
578	CD	GLN 108	A	18.119	38.791	9.358	0.99
579	OE1	GLN 108	A	17.035	38.401	8.916	0.83
580	NE2	GLN 108	A	18.624	38.356	10.508	0.94
581	C	GLN 108	A	18.217	41.519	6.135	0.63
582	O	GLN 108	A	18.323	42.360	7.047	0.64
583	N	LEU 109	A	17.213	41.401	5.288	0.60
584	CA	LEU 109	A	15.994	42.229	5.307	0.65
585	CB	LEU 109	A	15.882	42.694	3.831	0.55

73/98

*Fig. 16*14

Atom	Residue	Chain	X	Y	Z	δ	
586	CG	LEU 109	A	17.101	43.501	3.407	0.59
587	CD1	LEU 109	A	17.359	43.341	1.936	0.68
588	CD2	LEU 109	A	16.896	44.975	3.746	0.56
589	C	LEU 109	A	14.740	41.492	5.774	0.70
590	O	LEU 109	A	14.393	40.460	5.142	0.75
591	N	ASN 110	A	13.876	42.190	6.507	0.67
592	CA	ASN 110	A	12.512	41.767	6.849	0.65
593	CB	ASN 110	A	12.192	41.773	8.350	0.59
594	CG	ASN 110	A	13.043	40.783	9.115	0.79
595	OD1	ASN 110	A	13.197	40.850	10.340	0.90
596	ND2	ASN 110	A	13.591	39.872	8.302	0.79
597	C	ASN 110	A	11.441	42.589	6.116	0.65
598	O	ASN 110	A	11.655	43.703	5.641	0.61
599	N	ALA 111	A	10.255	41.992	6.127	0.68
600	CA	ALA 111	A	9.062	42.537	5.489	0.64
601	CB	ALA 111	A	8.158	41.466	4.906	0.59
602	C	ALA 111	A	8.315	43.435	6.467	0.57
603	O	ALA 111	A	8.500	43.354	7.697	0.54
604	N	ILE 112	A	7.548	44.344	5.879	0.54
605	CA	ILE 112	A	6.518	45.030	6.743	0.47
606	CB	ILE 112	A	6.821	46.531	6.985	0.63
607	CG2	ILE 112	A	7.990	46.879	7.923	0.68
608	CG1	ILE 112	A	6.825	47.379	5.712	0.56
609	CD	ILE 112	A	6.369	48.858	5.888	0.64
610	C	ILE 112	A	5.183	44.777	5.997	0.45
611	O	ILE 112	A	5.166	44.766	4.747	0.54
612	N	SER 113	A	4.150	44.802	6.770	0.44
613	CA	SER 113	A	2.734	44.868	6.330	0.61
614	CB	SER 113	A	1.946	43.791	7.102	0.63
615	OG	SER 113	A	2.163	42.503	6.515	0.62
616	C	SER 113	A	2.172	46.244	6.694	0.53
617	O	SER 113	A	2.299	46.730	7.828	0.52
618	N	VAL 114	A	1.400	46.804	5.778	0.55
619	CA	VAL 114	A	0.651	48.025	6.121	0.58
620	CB	VAL 114	A	1.371	49.244	5.514	0.64
621	CG1	VAL 114	A	2.885	49.190	5.667	0.49
622	CG2	VAL 114	A	0.949	49.424	4.094	0.55
623	C	VAL 114	A	-0.813	47.879	5.686	0.60
624	O	VAL 114	A	-1.189	47.327	4.637	0.51
625	N	LEU 115	A	-1.652	48.484	6.510	0.49
626	CA	LEU 115	A	-3.116	48.419	6.182	0.58
627	CB	LEU 115	A	-3.789	48.328	7.603	0.55
628	CG	LEU 115	A	-5.284	47.999	7.568	0.57
629	CD1	LEU 115	A	-5.523	46.603	7.036	0.59
630	CD2	LEU 115	A	-5.843	48.190	8.975	0.52

74/98

Fig. 16\15

Atom	Residue	Chain	X	Y	Z	δ	
631	C	LEU 115	A	-3.421	49.685	5.403	0.47
632	O	LEU 115	A	-3.244	50.761	5.955	0.57
633	N	TYR 116	A	-3.971	49.616	4.196	0.57
634	CA	TYR 116	A	-4.302	50.847	3.479	0.69
635	CB	TYR 116	A	-3.192	51.211	2.480	0.70
636	CG	TYR 116	A	-3.119	50.316	1.269	0.46
637	CD1	TYR 116	A	-2.487	49.097	1.272	0.57
638	CE1	TYR 116	A	-2.542	48.256	0.154	0.66
639	CD2	TYR 116	A	-3.584	50.775	0.050	0.57
640	CE2	TYR 116	A	-3.689	49.971	-1.067	0.52
641	CZ	TYR 116	A	-3.151	48.697	-1.016	0.76
642	OH	TYR 116	A	-3.119	47.979	-2.184	0.78
643	C	TYR 116	A	-5.692	50.766	2.849	0.69
644	O	TYR 116	A	-6.250	49.665	2.752	0.69
645	N	PHE 117	A	-6.209	51.919	2.485	0.56
646	CA	PHE 117	A	-7.443	52.158	1.793	0.60
647	CB	PHE 117	A	-8.462	53.064	2.413	0.64
648	CG	PHE 117	A	-8.074	54.254	3.206	0.82
649	CD1	PHE 117	A	-8.221	54.256	4.599	0.91
650	CD2	PHE 117	A	-7.813	55.472	2.561	1.11
651	CE1	PHE 117	A	-7.964	55.401	5.343	1.12
652	CE2	PHE 117	A	-7.547	56.638	3.292	1.03
653	CZ	PHE 117	A	-7.638	56.601	4.697	1.02
654	C	PHE 117	A	-7.240	52.362	0.315	0.62
655	O	PHE 117	A	-6.666	53.381	-0.059	0.72
656	N	ASP 118	A	-7.607	51.365	-0.474	0.67
657	CA	ASP 118	A	-7.388	51.415	-1.922	0.67
658	CB	ASP 118	A	-7.342	50.074	-2.610	0.73
659	CG	ASP 118	A	-8.694	49.558	-3.079	0.82
660	OD1	ASP 118	A	-8.748	48.554	-3.798	0.72
661	OD2	ASP 118	A	-9.705	50.249	-2.831	0.60
662	C	ASP 118	A	-8.337	52.433	-2.526	0.68
663	O	ASP 118	A	-9.108	53.096	-1.826	0.75
664	N	ASP 119	A	-8.344	52.429	-3.851	0.76
665	CA	ASP 119	A	-9.035	53.486	-4.614	0.83
666	CB	ASP 119	A	-8.356	53.711	-5.961	0.94
667	CG	ASP 119	A	-7.989	52.393	-6.634	1.08
668	OD1	ASP 119	A	-8.883	51.605	-6.991	1.04
669	OD2	ASP 119	A	-6.773	52.103	-6.627	1.06
670	C	ASP 119	A	-10.531	53.204	-4.673	0.87
671	O	ASP 119	A	-11.339	54.138	-4.854	0.90
672	N	SER 120	A	-10.886	51.946	-4.448	0.85
673	CA	SER 120	A	-12.314	51.574	-4.389	0.80
674	CB	SER 120	A	-12.618	50.258	-5.030	0.83
675	OG	SER 120	A	-11.534	49.729	-5.781	1.02

75/98

Fig. 16\16

Atom	Residue	Chain	X	Y	Z	δ	
676	C	SER 120	A	-12.796	51.719	-2.957	0.80
677	O	SER 120	A	-14.013	51.770	-2.686	0.83
678	N	SER 121	A	-11.823	51.902	-2.064	0.73
679	CA	SER 121	A	-12.197	52.163	-0.653	0.69
680	CB	SER 121	A	-13.578	52.816	-0.607	0.66
681	OG	SER 121	A	-13.479	54.163	-1.016	0.85
682	C	SER 121	A	-12.186	50.879	0.163	0.63
683	O	SER 121	A	-12.564	50.874	1.332	0.69
684	N	ASN 122	A	-11.745	49.825	-0.477	0.65
685	CA	ASN 122	A	-11.388	48.578	0.218	0.63
686	CB	ASN 122	A	-11.103	47.558	-0.897	0.69
687	CG	ASN 122	A	-12.211	47.680	-1.955	0.73
688	OD1	ASN 122	A	-13.370	47.437	-1.597	0.73
689	ND2	ASN 122	A	-11.867	47.636	-3.240	0.75
690	C	ASN 122	A	-10.237	48.837	1.176	0.67
691	O	ASN 122	A	-9.323	49.604	0.852	0.63
692	N	VAL 123	A	-10.305	48.227	2.346	0.66
693	CA	VAL 123	A	-9.252	48.270	3.360	0.61
694	CB	VAL 123	A	-9.927	48.279	4.744	0.58
695	CG1	VAL 123	A	-8.929	47.747	5.758	0.59
696	CG2	VAL 123	A	-10.405	49.677	5.089	0.56
697	C	VAL 123	A	-8.351	47.044	3.220	0.63
698	O	VAL 123	A	-8.901	45.942	3.098	0.65
699	N	ILE 124	A	-7.133	47.256	2.758	0.70
700	CA	ILE 124	A	-6.224	46.189	2.302	0.65
701	CB	ILE 124	A	-5.819	46.559	0.813	0.67
702	CG2	ILE 124	A	-4.839	45.531	0.198	0.67
703	CG1	ILE 124	A	-7.136	46.616	0.011	0.66
704	CD	ILE 124	A	-7.044	46.191	-1.468	0.84
705	C	ILE 124	A	-4.998	46.057	3.208	0.67
706	O	ILE 124	A	-4.326	47.034	3.582	0.63
707	N	LEU 125	A	-4.598	44.815	3.448	0.63
708	CA	LEU 125	A	-3.323	44.541	4.140	0.69
709	CB	LEU 125	A	-3.602	43.477	5.187	0.67
710	CG	LEU 125	A	-2.441	43.036	6.067	0.64
711	CD1	LEU 125	A	-2.163	44.105	7.130	0.49
712	CD2	LEU 125	A	-2.977	41.760	6.750	0.48
713	C	LEU 125	A	-2.249	44.132	3.135	0.67
714	O	LEU 125	A	-2.431	43.103	2.470	0.66
715	N	LYS 126	A	-1.263	44.993	2.961	0.66
716	CA	LYS 126	A	-0.174	44.777	2.001	0.59
717	CB	LYS 126	A	-0.157	45.640	0.788	0.61
718	CG	LYS 126	A	1.066	45.500	-0.128	0.78
719	CD	LYS 126	A	0.687	45.478	-1.601	0.91
720	CE	LYS 126	A	1.833	45.164	-2.543	0.82

76/98

Fig. 16\17

Atom	Residue	Chain	X	Y	Z	δ	
721	NZ	LYS 126	A	1.346	44.434	-3.746	0.86
722	C	LYS 126	A	1.173	44.619	2.669	0.50
723	O	LYS 126	A	1.398	44.978	3.845	0.57
724	N	LYS 127	A	1.836	43.565	2.204	0.61
725	CA	LYS 127	A	3.124	43.060	2.664	0.60
726	CB	LYS 127	A	3.188	41.534	2.620	0.58
727	CG	LYS 127	A	4.330	40.965	3.471	0.57
728	CD	LYS 127	A	4.828	39.671	2.833	0.62
729	CE	LYS 127	A	5.639	38.881	3.845	0.76
730	NZ	LYS 127	A	5.283	39.360	5.208	0.83
731	C	LYS 127	A	4.245	43.589	1.755	0.60
732	O	LYS 127	A	4.058	43.699	0.521	0.49
733	N	TYR 128	A	5.136	44.302	2.424	0.56
734	CA	TYR 128	A	6.222	44.975	1.669	0.50
735	CB	TYR 128	A	6.302	46.457	2.089	0.77
736	CG	TYR 128	A	5.310	47.284	1.298	0.64
737	CD1	TYR 128	A	4.269	47.954	1.935	0.68
738	CE1	TYR 128	A	3.320	48.646	1.174	0.68
739	CD2	TYR 128	A	5.280	47.163	-0.085	0.72
740	CE2	TYR 128	A	4.318	47.820	-0.857	0.77
741	CZ	TYR 128	A	3.268	48.459	-0.200	0.89
742	OH	TYR 128	A	2.279	49.021	-0.961	0.90
743	C	TYR 128	A	7.527	44.284	2.110	0.46
744	O	TYR 128	A	7.781	44.139	3.308	0.59
745	N	ARG 129	A	7.978	43.511	1.150	0.56
746	CA	ARG 129	A	9.194	42.698	1.396	0.61
747	CB	ARG 129	A	9.175	41.646	0.273	0.74
748	CG	ARG 129	A	8.115	40.550	0.447	0.74
749	CD	ARG 129	A	8.486	39.314	-0.337	0.61
750	NE	ARG 129	A	9.295	38.414	0.488	1.06
751	CZ	ARG 129	A	10.599	38.183	0.310	1.07
752	NH1	ARG 129	A	11.211	38.363	-0.861	0.91
753	NH2	ARG 129	A	11.370	37.849	1.349	0.87
754	C	ARG 129	A	10.405	43.636	1.302	0.54
755	O	ARG 129	A	10.406	44.605	0.522	0.55
756	N	ASN 130	A	11.403	43.372	2.095	0.60
757	CA	ASN 130	A	12.792	43.856	2.013	0.59
758	CB	ASN 130	A	13.359	43.611	0.610	0.52
759	CG	ASN 130	A	13.760	42.157	0.392	0.63
760	OD1	ASN 130	A	14.053	41.398	1.318	0.58
761	ND2	ASN 130	A	13.517	41.609	-0.782	0.55
762	C	ASN 130	A	12.807	45.343	2.380	0.55
763	O	ASN 130	A	13.426	46.094	1.648	0.48
764	N	MET 131	A	12.267	45.671	3.520	0.54
765	CA	MET 131	A	11.990	47.026	3.974	0.51

77/98

Fig. 16\18

Atom	Residue	Chain	X	Y	Z	δ	
766	CB	MET 131	A	10.470	47.189	4.327	0.54
767	CG	MET 131	A	9.590	47.290	3.137	0.44
768	SD	MET 131	A	9.927	48.720	2.052	0.61
769	CE	MET 131	A	8.873	49.928	2.915	0.60
770	C	MET 131	A	12.790	47.266	5.256	0.52
771	O	MET 131	A	13.048	48.429	5.570	0.55
772	N	VAL 132	A	12.997	46.230	6.055	0.59
773	CA	VAL 132	A	13.670	46.399	7.359	0.60
774	CB	VAL 132	A	12.807	46.001	8.517	0.52
775	CG1	VAL 132	A	13.412	46.036	9.898	0.48
776	CG2	VAL 132	A	11.331	46.318	8.485	0.66
777	C	VAL 132	A	15.035	45.704	7.308	0.69
778	O	VAL 132	A	15.148	44.495	7.012	0.63
779	N	VAL 133	A	16.039	46.488	7.668	0.65
780	CA	VAL 133	A	17.424	45.971	7.709	0.62
781	CB	VAL 133	A	18.486	47.081	7.591	0.56
782	CG1	VAL 133	A	19.842	46.568	8.105	0.53
783	CG2	VAL 133	A	18.577	47.505	6.128	0.39
784	C	VAL 133	A	17.542	45.241	9.052	0.62
785	O	VAL 133	A	17.392	45.958	10.044	0.54
786	N	ARG 134	A	17.821	43.930	8.941	0.53
787	CA	ARG 134	A	17.924	43.188	10.208	0.56
788	CB	ARG 134	A	17.114	41.877	10.158	0.68
789	CG	ARG 134	A	15.779	41.942	10.908	0.80
790	CD	ARG 134	A	15.932	41.805	12.379	0.79
791	NE	ARG 134	A	15.072	40.814	12.976	1.02
792	CZ	ARG 134	A	15.265	39.514	13.159	1.10
793	NH1	ARG 134	A	16.185	38.783	12.533	1.08
794	NH2	ARG 134	A	14.403	38.858	13.952	1.09
795	C	ARG 134	A	19.370	42.957	10.622	0.57
796	O	ARG 134	A	19.660	42.742	11.813	0.61
797	N	ALA 135	A	20.282	42.931	9.659	0.53
798	CA	ALA 135	A	21.722	42.782	9.986	0.49
799	CB	ALA 135	A	22.059	41.272	9.934	0.64
800	C	ALA 135	A	22.466	43.400	8.774	0.42
801	O	ALA 135	A	21.937	43.282	7.673	0.49
802	N	CYS 136	A	23.726	43.635	9.022	0.59
803	CA	CYS 136	A	24.682	44.274	8.089	0.67
804	C	CYS 136	A	25.954	43.443	7.969	0.61
805	O	CYS 136	A	26.383	42.850	8.985	0.59
806	CB	CYS 136	A	25.073	45.641	8.701	0.61
807	SG	CYS 136	A	23.632	46.772	8.673	0.61
808	N	GLY 137	A	26.579	43.476	6.808	0.59
809	CA	GLY 137	A	27.915	42.790	6.772	0.59
810	C	GLY 137	A	28.804	43.523	5.787	0.52

78/98

Fig. 16\19

Atom	Residue	Chain	X	Y	Z	δ	
811	O	GLY 137	A	28.249	44.112	4.845	0.49
812	N	CYS 138	A	30.019	43.036	5.634	0.57
813	CA	CYS 138	A	30.912	43.537	4.545	0.48
814	C	CYS 138	A	30.845	42.606	3.369	0.56
815	O	CYS 138	A	31.335	41.457	3.446	0.55
816	CB	CYS 138	A	32.348	43.567	5.123	0.38
817	SG	CYS 138	A	32.401	44.877	6.428	0.57
818	N	HIS 139	A	30.522	43.184	2.210	0.63
819	CA	HIS 139	A	30.557	42.395	0.970	0.54
820	CB	HIS 139	A	29.665	41.188	0.842	0.88
821	CG	HIS 139	A	28.437	41.289	1.707	0.94
822	CD2	HIS 139	A	27.571	42.337	1.811	0.79
823	ND1	HIS 139	A	28.113	40.406	2.706	0.91
824	CE1	HIS 139	A	27.001	40.825	3.288	0.97
825	NE2	HIS 139	A	26.665	41.993	2.774	0.96
826	C	HIS 139	A	30.936	43.135	-0.264	0.58
827	OT1	HIS 139	A	31.164	42.426	-1.268	0.75
828	OT2	HIS 139	A	31.123	44.371	-0.256	0.79
829	CB	GLN 36	B	29.653	57.175	-11.979	0.80
830	CG	GLN 36	B	30.943	57.649	-11.327	0.98
831	CD	GLN 36	B	30.751	58.560	-10.142	1.07
832	OE1	GLN 36	B	29.879	59.426	-10.061	0.98
833	NE2	GLN 36	B	31.585	58.331	-9.123	1.05
834	C	GLN 36	B	28.572	54.939	-11.583	0.62
835	O	GLN 36	B	28.641	54.928	-10.349	0.67
836	N	GLN 36	B	29.482	55.414	-13.781	0.82
837	CA	GLN 36	B	29.627	55.707	-12.360	0.72
838	N	ALA 37	B	27.621	54.379	-12.314	0.65
839	CA	ALA 37	B	26.512	53.646	-11.644	0.64
840	CB	ALA 37	B	25.331	53.455	-12.576	0.58
841	C	ALA 37	B	27.011	52.266	-11.201	0.62
842	O	ALA 37	B	27.851	51.674	-11.868	0.67
843	N	CYS 38	B	26.260	51.702	-10.285	0.64
844	CA	CYS 38	B	26.531	50.390	-9.714	0.53
845	C	CYS 38	B	26.592	49.339	-10.797	0.63
846	O	CYS 38	B	25.623	49.141	-11.527	0.66
847	CB	CYS 38	B	25.607	50.071	-8.582	0.58
848	SG	CYS 38	B	25.827	48.462	-7.828	0.63
849	N	LYS 39	B	27.800	48.823	-11.010	0.63
850	CA	LYS 39	B	28.056	47.677	-11.860	0.50
851	CB	LYS 39	B	28.442	48.020	-13.296	0.56
852	CG	LYS 39	B	29.121	49.356	-13.549	0.69
853	CD	LYS 39	B	29.867	49.395	-14.870	0.62
854	CE	LYS 39	B	29.840	48.060	-15.592	0.92
855	NZ	LYS 39	B	30.922	47.935	-16.607	0.92

79/98

Fig. 16\20

Atom	Residue	Chain	X	Y	Z	δ	
856	C	LYS 39	B	29.158	46.771	-11.262	0.50
857	O	LYS 39	B	29.788	47.073	-10.242	0.49
858	N	LYS 40	B	29.405	45.717	-11.988	0.59
859	CA	LYS 40	B	30.309	44.615	-11.659	0.64
860	CB	LYS 40	B	29.713	43.258	-12.132	0.61
861	CG	LYS 40	B	30.617	42.119	-11.621	0.60
862	CD	LYS 40	B	30.087	40.791	-12.085	0.48
863	CE	LYS 40	B	30.264	40.554	-13.565	0.60
864	NZ	LYS 40	B	29.587	39.269	-13.873	0.73
865	C	LYS 40	B	31.570	44.807	-12.504	0.63
866	O	LYS 40	B	31.450	45.250	-13.659	0.61
867	N	HIS 41	B	32.700	44.888	-11.826	0.65
868	CA	HIS 41	B	33.938	45.171	-12.590	0.58
869	CB	HIS 41	B	34.627	46.413	-12.011	0.57
870	CG	HIS 41	B	33.639	47.549	-11.899	0.60
871	CD2	HIS 41	B	32.931	47.989	-10.838	0.63
872	ND1	HIS 41	B	33.564	48.557	-12.835	0.61
873	CE1	HIS 41	B	32.837	49.537	-12.338	0.63
874	NE2	HIS 41	B	32.274	49.130	-11.217	0.67
875	C	HIS 41	B	34.837	43.952	-12.505	0.58
876	O	HIS 41	B	34.737	43.143	-11.564	0.60
877	N	GLU 42	B	35.819	43.974	-13.373	0.63
878	CA	GLU 42	B	36.805	42.897	-13.460	0.65
879	CB	GLU 42	B	37.296	42.767	-14.894	0.67
880	CG	GLU 42	B	36.418	41.808	-15.724	0.61
881	CD	GLU 42	B	36.806	41.769	-17.171	0.80
882	OE1	GLU 42	B	36.002	41.778	-18.082	0.89
883	OE2	GLU 42	B	38.046	41.851	-17.309	0.92
884	C	GLU 42	B	37.961	43.043	-12.507	0.63
885	O	GLU 42	B	38.361	44.183	-12.261	0.68
886	N	LEU 43	B	38.349	41.959	-11.858	0.57
887	CA	LEU 43	B	39.538	41.815	-11.012	0.57
888	CB	LEU 43	B	39.120	42.016	-9.561	0.56
889	CG	LEU 43	B	39.970	42.470	-8.455	0.65
890	CD1	LEU 43	B	39.673	42.100	-7.043	0.59
891	CD2	LEU 43	B	41.300	43.057	-8.715	0.64
892	C	LEU 43	B	39.947	40.315	-11.210	0.63
893	O	LEU 43	B	39.206	39.457	-10.710	0.62
894	N	TYR 44	B	41.180	40.079	-11.580	0.58
895	CA	TYR 44	B	41.745	38.704	-11.590	0.54
896	CB	TYR 44	B	42.658	38.620	-12.845	0.52
897	CG	TYR 44	B	43.133	37.198	-13.022	0.63
898	CD1	TYR 44	B	44.404	36.795	-12.620	0.50
899	CE1	TYR 44	B	44.762	35.449	-12.742	0.58
900	CD2	TYR 44	B	42.295	36.274	-13.634	0.52

80/98

Fig. 16\21

Atom	Residue	Chain	X	Y	Z	δ	
901	CE2	TYR 44	B	42.615	34.946	-13.694	0.56
902	CZ	TYR 44	B	43.862	34.534	-13.235	0.69
903	OH	TYR 44	B	44.205	33.222	-13.434	0.81
904	C	TYR 44	B	42.696	38.638	-10.395	0.51
905	O	TYR 44	B	43.481	39.582	-10.316	0.63
906	N	VAL 45	B	42.581	37.674	-9.534	0.53
907	CA	VAL 45	B	43.473	37.480	-8.368	0.53
908	CB	VAL 45	B	42.579	37.181	-7.148	0.56
909	CG1	VAL 45	B	43.320	36.968	-5.844	0.47
910	CG2	VAL 45	B	41.431	38.176	-7.035	0.52
911	C	VAL 45	B	44.398	36.280	-8.673	0.55
912	O	VAL 45	B	43.906	35.262	-9.193	0.53
913	N	SER 46	B	45.669	36.588	-8.765	0.59
914	CA	SER 46	B	46.829	35.718	-8.858	0.52
915	CB	SER 46	B	48.122	36.464	-9.287	0.46
916	OG	SER 46	B	48.841	35.585	-10.135	0.69
917	C	SER 46	B	47.138	35.163	-7.482	0.52
918	O	SER 46	B	47.307	35.955	-6.553	0.56
919	N	PHE 47	B	47.216	33.857	-7.380	0.59
920	CA	PHE 47	B	47.583	33.256	-6.069	0.57
921	CB	PHE 47	B	47.312	31.754	-6.088	0.53
922	CG	PHE 47	B	45.896	31.295	-6.174	0.55
923	CD1	PHE 47	B	44.869	32.001	-5.517	0.45
924	CD2	PHE 47	B	45.562	30.101	-6.818	0.54
925	CE1	PHE 47	B	43.554	31.589	-5.633	0.71
926	CE2	PHE 47	B	44.232	29.715	-7.010	0.56
927	CZ	PHE 47	B	43.221	30.426	-6.313	0.60
928	C	PHE 47	B	48.970	33.743	-5.648	0.49
929	O	PHE 47	B	49.222	33.901	-4.426	0.61
930	N	ARG 48	B	49.912	33.923	-6.541	0.51
931	CA	ARG 48	B	51.204	34.593	-6.314	0.57
932	CB	ARG 48	B	52.003	34.854	-7.564	0.49
933	CG	ARG 48	B	52.089	33.858	-8.662	0.74
934	CD	ARG 48	B	53.439	33.274	-8.844	0.83
935	NE	ARG 48	B	54.192	33.768	-9.964	0.71
936	CZ	ARG 48	B	54.800	33.088	-10.923	0.66
937	NH1	ARG 48	B	55.174	31.805	-10.831	0.68
938	NH2	ARG 48	B	54.842	33.671	-12.123	0.61
939	C	ARG 48	B	51.122	35.885	-5.517	0.64
940	O	ARG 48	B	51.681	36.007	-4.397	0.69
941	N	ASP 49	B	50.284	36.811	-5.977	0.60
942	CA	ASP 49	B	49.970	38.044	-5.274	0.57
943	CB	ASP 49	B	48.961	38.897	-6.047	0.68
944	CG	ASP 49	B	49.395	39.128	-7.481	0.80
945	OD1	ASP 49	B	48.609	39.683	-8.272	0.89

81/98

*Fig. 16*22

Atom	Residue	Chain	X	Y	Z	δ	
946	OD2	ASP 49	B	50.527	38.681	-7.801	0.83
947	C	ASP 49	B	49.619	37.883	-3.814	0.66
948	O	ASP 49	B	49.712	38.876	-3.062	0.72
949	N	LEU 50	B	48.879	36.830	-3.465	0.70
950	CA	LEU 50	B	48.428	36.605	-2.086	0.60
951	CB	LEU 50	B	47.115	35.800	-2.187	0.62
952	CG	LEU 50	B	46.024	36.479	-3.022	0.69
953	CD1	LEU 50	B	44.677	35.940	-2.544	0.66
954	CD2	LEU 50	B	46.100	37.974	-2.667	0.66
955	C	LEU 50	B	49.471	35.808	-1.275	0.62
956	O	LEU 50	B	49.097	35.367	-0.167	0.69
957	N	GLY 51	B	50.399	35.235	-2.005	0.62
958	CA	GLY 51	B	51.320	34.191	-1.563	0.59
959	C	GLY 51	B	50.554	32.900	-1.303	0.68
960	O	GLY 51	B	50.408	32.521	-0.130	0.73
961	N	TRP 52	B	49.891	32.356	-2.325	0.60
962	CA	TRP 52	B	49.127	31.117	-2.063	0.63
963	CB	TRP 52	B	47.650	31.299	-2.022	0.76
964	CG	TRP 52	B	46.961	32.057	-0.951	0.80
965	CD2	TRP 52	B	45.593	32.517	-1.004	0.74
966	CE2	TRP 52	B	45.285	33.063	0.263	0.92
967	CE3	TRP 52	B	44.583	32.450	-1.963	1.01
968	CD1	TRP 52	B	47.376	32.267	0.333	0.82
969	NE1	TRP 52	B	46.402	32.927	1.050	0.73
970	CZ2	TRP 52	B	44.009	33.536	0.576	0.75
971	CZ3	TRP 52	B	43.332	32.969	-1.684	0.68
972	CH2	TRP 52	B	43.053	33.511	-0.428	0.80
973	C	TRP 52	B	49.615	30.052	-3.046	0.72
974	O	TRP 52	B	49.190	28.891	-2.965	0.78
975	N	GLN 53	B	50.656	30.412	-3.769	0.64
976	CA	GLN 53	B	51.203	29.563	-4.822	0.74
977	CB	GLN 53	B	52.095	30.349	-5.798	0.68
978	CG	GLN 53	B	53.334	30.917	-5.138	0.56
979	CD	GLN 53	B	53.220	32.124	-4.268	0.78
980	OE1	GLN 53	B	52.394	32.252	-3.363	0.75
981	NE2	GLN 53	B	54.224	33.011	-4.447	0.80
982	C	GLN 53	B	51.891	28.321	-4.280	0.74
983	O	GLN 53	B	52.298	27.451	-5.080	0.83
984	N	ASP 54	B	52.335	28.353	-3.040	0.77
985	CA	ASP 54	B	53.153	27.288	-2.432	0.76
986	CB	ASP 54	B	53.499	27.716	-1.004	0.92
987	CG	ASP 54	B	54.614	28.739	-0.920	0.92
988	OD1	ASP 54	B	55.409	28.905	-1.863	0.96
989	OD2	ASP 54	B	54.784	29.235	0.217	0.75
990	C	ASP 54	B	52.486	25.906	-2.491	0.80

82/98

Fig. 16\23

Atom	Residue	Chain	X	Y	Z	δ
991 O	ASP 54	B	53.092	24.893	-2.898	0.82
992 N	TRP 55	B	51.285	25.839	-1.953	0.72
993 CA	TRP 55	B	50.495	24.644	-1.723	0.78
994 CB	TRP 55	B	49.926	24.753	-0.293	0.69
995 CG	TRP 55	B	51.035	24.962	0.691	0.80
996 CD2	TRP 55	B	52.396	24.529	0.465	0.76
997 CE2	TRP 55	B	53.149	24.987	1.569	0.85
998 CE3	TRP 55	B	52.924	23.554	-0.374	0.87
999 CD1	TRP 55	B	51.076	25.759	1.796	0.74
1000 NE1	TRP 55	B	52.345	25.793	2.337	0.76
1001 CZ2	TRP 55	B	54.479	24.626	1.745	0.78
1002 CZ3	TRP 55	B	54.216	23.134	-0.151	0.88
1003 CH2	TRP 55	B	54.968	23.639	0.905	0.82
1004 C	TRP 55	B	49.431	24.336	-2.770	0.84
1005 O	TRP 55	B	48.906	23.197	-2.758	0.77
1006 N	ILE 56	B	49.105	25.287	-3.642	0.83
1007 CA	ILE 56	B	48.109	25.152	-4.693	0.78
1008 CB	ILE 56	B	47.382	26.490	-5.076	0.77
1009 CG2	ILE 56	B	47.094	26.568	-6.606	0.71
1010 CG1	ILE 56	B	46.030	26.567	-4.316	0.82
1011 CD	ILE 56	B	45.964	27.743	-3.298	0.90
1012 C	ILE 56	B	48.700	24.512	-5.944	0.74
1013 O	ILE 56	B	49.768	24.950	-6.388	0.80
1014 N	ILE 57	B	47.911	23.642	-6.567	0.68
1015 CA	ILE 57	B	48.281	22.985	-7.822	0.64
1016 CB	ILE 57	B	48.040	21.429	-7.725	0.65
1017 CG2	ILE 57	B	47.983	20.771	-9.109	0.55
1018 CG1	ILE 57	B	49.017	20.758	-6.736	0.64
1019 CD	ILE 57	B	48.780	19.230	-6.540	0.62
1020 C	ILE 57	B	47.565	23.593	-9.020	0.60
1021 O	ILE 57	B	48.093	23.570	-10.145	0.62
1022 N	ALA 58	B	46.279	23.911	-8.834	0.58
1023 CA	ALA 58	B	45.460	24.436	-9.967	0.49
1024 CB	ALA 58	B	44.970	23.215	-10.789	0.55
1025 C	ALA 58	B	44.202	25.091	-9.376	0.46
1026 O	ALA 58	B	43.929	24.534	-8.302	0.49
1027 N	PRO 59	B	43.872	26.310	-9.744	0.53
1028 CD	PRO 59	B	42.742	27.052	-9.192	0.46
1029 CA	PRO 59	B	44.401	27.103	-10.870	0.45
1030 CB	PRO 59	B	43.209	27.839	-11.429	0.47
1031 CG	PRO 59	B	42.321	28.059	-10.217	0.43
1032 C	PRO 59	B	45.527	28.013	-10.384	0.53
1033 O	PRO 59	B	46.032	27.913	-9.239	0.55
1034 N	GLU 60	B	46.018	28.888	-11.255	0.49
1035 CA	GLU 60	B	47.061	29.844	-10.776	0.55

83/98

Fig. 16\24

Atom	Residue	Chain	X	Y	Z	δ	
1036	CB	GLU 60	B	48.099	30.025	-11.906	0.64
1037	CG	GLU 60	B	49.287	29.075	-11.886	0.77
1038	CD	GLU 60	B	49.851	28.404	-13.082	0.79
1039	OE1	GLU 60	B	49.900	27.172	-13.170	0.84
1040	OE2	GLU 60	B	50.265	29.162	-13.997	0.61
1041	C	GLU 60	B	46.440	31.175	-10.374	0.53
1042	O	GLU 60	B	47.006	31.966	-9.605	0.52
1043	N	GLY 61	B	45.121	31.271	-10.496	0.52
1044	CA	GLY 61	B	44.275	32.336	-9.965	0.54
1045	C	GLY 61	B	42.888	32.324	-10.585	0.58
1046	O	GLY 61	B	42.568	31.365	-11.298	0.55
1047	N	TYR 62	B	42.036	33.285	-10.231	0.55
1048	CA	TYR 62	B	40.662	33.247	-10.845	0.54
1049	CB	TYR 62	B	39.780	32.410	-9.853	0.38
1050	CG	TYR 62	B	39.540	33.198	-8.558	0.47
1051	CD1	TYR 62	B	40.580	33.288	-7.649	0.47
1052	CE1	TYR 62	B	40.540	34.084	-6.499	0.51
1053	CD2	TYR 62	B	38.335	33.785	-8.201	0.45
1054	CE2	TYR 62	B	38.165	34.346	-6.912	0.35
1055	CZ	TYR 62	B	39.284	34.567	-6.115	0.51
1056	OH	TYR 62	B	39.146	35.055	-4.839	0.65
1057	C	TYR 62	B	40.133	34.672	-10.973	0.59
1058	O	TYR 62	B	40.590	35.651	-10.330	0.55
1059	N	ALA 63	B	39.076	34.789	-11.731	0.54
1060	CA	ALA 63	B	38.403	36.106	-11.984	0.49
1061	CB	ALA 63	B	37.619	35.856	-13.282	0.58
1062	C	ALA 63	B	37.422	36.309	-10.845	0.51
1063	O	ALA 63	B	36.491	35.511	-10.665	0.58
1064	N	ALA 64	B	37.780	37.157	-9.895	0.54
1065	CA	ALA 64	B	36.931	37.347	-8.735	0.53
1066	CB	ALA 64	B	37.812	37.741	-7.550	0.53
1067	C	ALA 64	B	35.776	38.344	-8.972	0.52
1068	O	ALA 64	B	34.882	38.380	-8.091	0.54
1069	N	TYR 65	B	36.063	39.402	-9.671	0.51
1070	CA	TYR 65	B	35.241	40.568	-9.978	0.54
1071	CB	TYR 65	B	33.801	40.235	-10.438	0.47
1072	CG	TYR 65	B	33.841	39.511	-11.777	0.47
1073	CD1	TYR 65	B	34.042	40.165	-12.974	0.44
1074	CE1	TYR 65	B	34.043	39.474	-14.190	0.47
1075	CD2	TYR 65	B	34.100	38.137	-11.756	0.61
1076	CE2	TYR 65	B	34.363	37.449	-12.922	0.44
1077	CZ	TYR 65	B	34.279	38.092	-14.139	0.67
1078	OH	TYR 65	B	34.476	37.314	-15.253	0.70
1079	C	TYR 65	B	35.164	41.470	-8.752	0.51
1080	O	TYR 65	B	35.488	40.943	-7.664	0.49

84/98

Fig. 16\25

Atom		Residuc		Chain		X	Y	Z	δ
1081	N	TYR	66	B		34.349	42.507	-8.928	0.52
1082	CA	TYR	66	B		33.974	43.311	-7.729	0.50
1083	CB	TYR	66	B		35.170	44.051	-7.126	0.60
1084	CG	TYR	66	B		35.700	45.179	-7.983	0.59
1085	CD1	TYR	66	B		36.704	44.935	-8.917	0.59
1086	CE1	TYR	66	B		37.279	45.951	-9.659	0.53
1087	CD2	TYR	66	B		35.376	46.512	-7.721	0.53
1088	CE2	TYR	66	B		36.057	47.548	-8.335	0.47
1089	CZ	TYR	66	B		36.836	47.261	-9.448	0.58
1090	OH	TYR	66	B		37.469	48.301	-10.066	0.70
1091	C	TYR	66	B		32.914	44.336	-8.166	0.51
1092	O	TYR	66	B		32.777	44.585	-9.351	0.52
1093	N	CYS	67	B		32.304	44.936	-7.173	0.58
1094	CA	CYS	67	B		31.148	45.843	-7.372	0.57
1095	C	CYS	67	B		31.597	47.265	-7.066	0.52
1096	O	CYS	67	B		32.160	47.475	-5.968	0.48
1097	CB	CYS	67	B		30.051	45.374	-6.413	0.54
1098	SG	CYS	67	B		29.272	43.787	-6.831	0.54
1099	N	GLU	68	B		31.226	48.194	-7.930	0.56
1100	CA	GLU	68	B		31.450	49.624	-7.617	0.47
1101	CB	GLU	68	B		32.888	50.026	-7.993	0.68
1102	CG	GLU	68	B		33.521	51.195	-7.264	0.53
1103	CD	GLU	68	B		34.985	51.442	-7.510	0.67
1104	OE1	GLU	68	B		35.681	52.049	-6.720	0.64
1105	OE2	GLU	68	B		35.357	51.239	-8.687	0.63
1106	C	GLU	68	B		30.561	50.515	-8.485	0.43
1107	O	GLU	68	B		30.393	50.238	-9.663	0.42
1108	N	GLY	69	B		30.221	51.681	-7.922	0.48
1109	CA	GLY	69	B		29.452	52.707	-8.669	0.44
1110	C	GLY	69	B		28.406	53.234	-7.670	0.51
1111	O	GLY	69	B		28.223	52.696	-6.563	0.53
1112	N	GLU	70	B		27.714	54.297	-8.048	0.57
1113	CA	GLU	70	B		26.797	54.923	-7.116	0.63
1114	CB	GLU	70	B		26.834	56.401	-6.924	0.74
1115	CG	GLU	70	B		26.449	57.323	-8.062	0.77
1116	CD	GLU	70	B		26.041	58.691	-7.554	1.02
1117	OE1	GLU	70	B		26.122	58.985	-6.372	0.84
1118	OE2	GLU	70	B		25.470	59.352	-8.441	0.95
1119	C	GLU	70	B		25.392	54.363	-7.133	0.44
1120	O	GLU	70	B		24.945	53.921	-8.186	0.42
1121	N	CYS	71	B		24.785	54.485	-5.962	0.50
1122	CA	CYS	71	B		23.378	54.069	-5.805	0.62
1123	C	CYS	71	B		22.488	55.305	-5.637	0.54
1124	O	CYS	71	B		22.279	55.787	-4.547	0.51
1125	CB	CYS	71	B		23.223	52.994	-4.771	0.62

85/98

Fig. 16|26

Atom	Residue	Chain	X	Y	Z	δ	
1126	SG	CYS 71	B	23.951	51.385	-5.196	0.57
1127	N	ALA 72	B	22.057	55.857	-6.745	0.50
1128	CA	ALA 72	B	21.341	57.102	-6.862	0.61
1129	CB	ALA 72	B	22.306	58.192	-7.425	0.61
1130	C	ALA 72	B	20.223	56.910	-7.903	0.62
1131	O	ALA 72	B	20.396	56.167	-8.880	0.59
1132	N	PHE 73	B	19.294	57.873	-7.785	0.59
1133	CA	PHE 73	B	18.244	57.978	-8.805	0.46
1134	CB	PHE 73	B	17.063	58.839	-8.472	0.54
1135	CG	PHE 73	B	16.335	58.428	-7.198	0.45
1136	CD1	PHE 73	B	15.725	57.183	-7.129	0.43
1137	CD2	PHE 73	B	16.303	59.266	-6.117	0.45
1138	CE1	PHE 73	B	15.178	56.702	-5.935	0.34
1139	CE2	PHE 73	B	15.627	58.907	-4.959	0.37
1140	CZ	PHE 73	B	15.089	57.602	-4.895	0.33
1141	C	PHE 73	B	18.983	58.181	-10.083	0.52
1142	O	PHE 73	B	19.751	59.134	-9.925	0.54
1143	N	PRO 74	B	18.492	57.715	-11.197	0.57
1144	CD	PRO 74	B	19.072	58.066	-12.495	0.57
1145	CA	PRO 74	B	17.638	56.569	-11.370	0.56
1146	CB	PRO 74	B	17.298	56.446	-12.832	0.52
1147	CG	PRO 74	B	18.376	57.215	-13.519	0.54
1148	C	PRO 74	B	18.219	55.293	-10.805	0.66
1149	O	PRO 74	B	19.370	54.968	-11.112	0.74
1150	N	LEU 75	B	17.359	54.601	-10.090	0.58
1151	CA	LEU 75	B	17.626	53.249	-9.627	0.61
1152	CB	LEU 75	B	17.151	53.088	-8.209	0.63
1153	CG	LEU 75	B	18.000	53.160	-7.006	0.60
1154	CD1	LEU 75	B	19.409	53.632	-7.025	0.57
1155	CD2	LEU 75	B	17.331	53.418	-5.703	0.49
1156	C	LEU 75	B	17.058	52.347	-10.726	0.68
1157	O	LEU 75	B	15.909	51.889	-10.655	0.72
1158	N	ASN 76	B	17.826	52.301	-11.802	0.68
1159	CA	ASN 76	B	17.569	51.408	-12.939	0.66
1160	CB	ASN 76	B	18.627	51.538	-14.028	0.62
1161	CG	ASN 76	B	18.661	52.899	-14.697	0.83
1162	OD1	ASN 76	B	17.657	53.637	-14.677	0.97
1163	ND2	ASN 76	B	19.757	53.260	-15.370	0.81
1164	C	ASN 76	B	17.288	50.001	-12.449	0.67
1165	O	ASN 76	B	17.535	49.608	-11.281	0.73
1166	N	SER 77	B	16.531	49.283	-13.260	0.73
1167	CA	SER 77	B	16.114	47.907	-12.886	0.74
1168	CB	SER 77	B	15.107	47.328	-13.894	0.54
1169	OG	SER 77	B	14.248	46.466	-13.128	0.96
1170	C	SER 77	B	17.359	47.011	-12.767	0.62

86/98

Fig. 16\27

Atom	Residuc	Chain	X	Y	Z	δ	
1171	O	SER 77	B	17.477	46.218	-11.835	0.66
1172	N	TYR 78	B	18.200	47.118	-13.767	0.63
1173	CA	TYR 78	B	19.542	46.592	-13.850	0.75
1174	CB	TYR 78	B	20.249	47.042	-15.137	0.63
1175	CG	TYR 78	B	21.368	48.043	-14.976	0.91
1176	CD1	TYR 78	B	21.284	49.371	-15.447	0.81
1177	CE1	TYR 78	B	22.321	50.280	-15.228	0.88
1178	CD2	TYR 78	B	22.605	47.610	-14.494	0.98
1179	CE2	TYR 78	B	23.652	48.502	-14.265	0.99
1180	CZ	TYR 78	B	23.510	49.833	-14.640	0.94
1181	OH	TYR 78	B	24.611	50.630	-14.483	0.97
1182	C	TYR 78	B	20.376	46.798	-12.592	0.78
1183	O	TYR 78	B	21.428	46.139	-12.487	0.79
1184	N	MET 79	B	20.014	47.722	-11.719	0.74
1185	CA	MET 79	B	20.721	48.022	-10.475	0.62
1186	CB	MET 79	B	20.757	49.519	-10.191	0.68
1187	CG	MET 79	B	21.670	50.154	-11.207	0.51
1188	SD	MET 79	B	21.423	51.909	-11.149	0.67
1189	CE	MET 79	B	22.086	52.461	-9.625	0.53
1190	C	MET 79	B	20.193	47.231	-9.309	0.58
1191	O	MET 79	B	20.701	47.287	-8.186	0.61
1192	N	ASN 80	B	19.249	46.327	-9.575	0.55
1193	CA	ASN 80	B	18.791	45.510	-8.431	0.54
1194	CB	ASN 80	B	19.392	44.106	-8.469	0.86
1195	CG	ASN 80	B	19.052	43.259	-7.244	1.05
1196	OD1	ASN 80	B	19.403	43.560	-6.079	0.95
1197	ND2	ASN 80	B	18.498	42.064	-7.518	0.86
1198	C	ASN 80	B	18.879	46.202	-7.091	0.57
1199	O	ASN 80	B	19.195	45.517	-6.054	0.67
1200	N	ALA 81	B	17.955	47.146	-6.890	0.60
1201	CA	ALA 81	B	17.798	47.835	-5.577	0.52
1202	CB	ALA 81	B	17.298	49.261	-5.894	0.53
1203	C	ALA 81	B	16.837	46.990	-4.802	0.53
1204	O	ALA 81	B	15.878	46.552	-5.445	0.60
1205	N	THR 82	B	16.965	46.811	-3.500	0.54
1206	CA	THR 82	B	15.866	46.297	-2.680	0.44
1207	CB	THR 82	B	16.477	45.801	-1.326	0.45
1208	OG1	THR 82	B	17.234	46.967	-0.821	0.49
1209	CG2	THR 82	B	17.599	44.729	-1.601	0.58
1210	C	THR 82	B	14.851	47.432	-2.443	0.67
1211	O	THR 82	B	15.035	48.606	-2.852	0.55
1212	N	ASN 83	B	13.681	47.096	-1.913	0.61
1213	CA	ASN 83	B	12.741	48.177	-1.529	0.56
1214	CB	ASN 83	B	11.453	47.641	-0.901	0.53
1215	CG	ASN 83	B	10.532	46.962	-1.900	0.48

87/98
Fig. 16\28

Atom		Residue	Chain	X	Y	Z	δ
1216	OD1	ASN 83	B	10.652	47.190	-3.108	0.55
1217	ND2	ASN 83	B	9.825	45.948	-1.410	0.55
1218	C	ASN 83	B	13.496	49.079	-0.535	0.55
1219	O	ASN 83	B	13.074	50.239	-0.428	0.54
1220	N	HIS 84	B	13.968	48.503	0.569	0.55
1221	CA	HIS 84	B	14.811	49.226	1.518	0.41
1222	CB	HIS 84	B	15.645	48.344	2.462	0.58
1223	CG	HIS 84	B	16.315	49.085	3.572	0.40
1224	CD2	HIS 84	B	15.849	49.305	4.809	0.29
1225	ND1	HIS 84	B	17.461	49.885	3.426	0.22
1226	CE1	HIS 84	B	17.727	50.393	4.612	0.18
1227	NE2	HIS 84	B	16.808	50.017	5.513	0.31
1228	C	HIS 84	B	15.742	50.241	0.878	0.47
1229	O	HIS 84	B	15.996	51.280	1.484	0.50
1230	N	ALA 85	B	16.457	49.932	-0.190	0.52
1231	CA	ALA 85	B	17.416	50.862	-0.773	0.53
1232	CB	ALA 85	B	18.193	50.157	-1.868	0.48
1233	C	ALA 85	B	16.658	52.059	-1.388	0.57
1234	O	ALA 85	B	17.384	53.017	-1.781	0.55
1235	N	ILE 86	B	15.570	51.726	-2.095	0.50
1236	CA	ILE 86	B	14.770	52.758	-2.783	0.52
1237	CB	ILE 86	B	13.487	52.116	-3.425	0.57
1238	CG2	ILE 86	B	12.542	53.287	-3.912	0.42
1239	CG1	ILE 86	B	13.972	51.293	-4.652	0.40
1240	CD	ILE 86	B	12.807	50.430	-5.234	0.49
1241	C	ILE 86	B	14.332	53.807	-1.751	0.45
1242	O	ILE 86	B	14.566	54.996	-1.894	0.58
1243	N	VAL 87	B	13.880	53.315	-0.618	0.49
1244	CA	VAL 87	B	13.507	54.142	0.517	0.42
1245	CB	VAL 87	B	12.892	53.242	1.570	0.43
1246	CG1	VAL 87	B	12.781	53.866	2.950	0.39
1247	CG2	VAL 87	B	11.517	52.782	1.097	0.50
1248	C	VAL 87	B	14.686	54.986	0.984	0.52
1249	O	VAL 87	B	14.415	55.916	1.765	0.57
1250	N	GLN 88	B	15.751	54.226	1.301	0.64
1251	CA	GLN 88	B	17.007	54.802	1.773	0.50
1252	CB	GLN 88	B	18.035	53.927	2.372	0.54
1253	CG	GLN 88	B	19.477	54.409	2.355	0.50
1254	CD	GLN 88	B	20.342	53.445	3.126	0.53
1255	OE1	GLN 88	B	19.914	52.340	3.448	0.63
1256	NE2	GLN 88	B	21.607	53.766	3.183	0.42
1257	C	GLN 88	B	17.473	55.905	0.867	0.42
1258	O	GLN 88	B	18.004	56.920	1.314	0.52
1259	N	THR 89	B	17.420	55.682	-0.420	0.52
1260	CA	THR 89	B	17.748	56.663	-1.434	0.51

88/98

Fig. 16|29

Atom	Residuc	Chain	X	Y	Z	δ	
1261	CB	THR 89	B	17.957	56.044	-2.830	0.44
1262	OG1	THR 89	B	18.892	54.925	-2.666	0.58
1263	CG2	THR 89	B	18.468	56.959	-3.906	0.28
1264	C	THR 89	B	16.809	57.866	-1.437	0.60
1265	O	THR 89	B	17.292	58.957	-1.772	0.60
1266	N	LEU 90	B	15.600	57.756	-0.944	0.60
1267	CA	LEU 90	B	14.594	58.825	-0.965	0.54
1268	CB	LEU 90	B	13.201	58.299	-1.267	0.64
1269	CG	LEU 90	B	12.008	59.230	-1.078	0.62
1270	CD1	LEU 90	B	11.967	60.201	-2.272	0.49
1271	CD2	LEU 90	B	10.728	58.385	-1.213	0.62
1272	C	LEU 90	B	14.771	59.667	0.279	0.47
1273	O	LEU 90	B	14.902	60.893	0.185	0.63
1274	N	VAL 91	B	15.098	59.037	1.376	0.43
1275	CA	VAL 91	B	15.498	59.640	2.618	0.40
1276	CB	VAL 91	B	15.662	58.617	3.726	0.35
1277	CG1	VAL 91	B	15.884	59.259	5.070	0.33
1278	CG2	VAL 91	B	14.368	57.821	3.941	0.47
1279	C	VAL 91	B	16.645	60.623	2.505	0.49
1280	O	VAL 91	B	16.915	61.323	3.477	0.59
1281	N	HIS 92	B	17.546	60.291	1.635	0.51
1282	CA	HIS 92	B	18.848	60.835	1.409	0.56
1283	CB	HIS 92	B	19.806	59.782	0.770	0.55
1284	CG	HIS 92	B	21.133	60.406	0.469	0.52
1285	CD2	HIS 92	B	21.587	60.995	-0.668	0.36
1286	ND1	HIS 92	B	22.072	60.627	1.483	0.42
1287	CE1	HIS 92	B	23.116	61.193	0.876	0.49
1288	NE2	HIS 92	B	22.856	61.461	-0.379	0.43
1289	C	HIS 92	B	18.699	62.058	0.492	0.50
1290	O	HIS 92	B	19.217	63.135	0.766	0.62
1291	N	PHE 93	B	17.866	61.925	-0.494	0.52
1292	CA	PHE 93	B	17.412	62.999	-1.335	0.62
1293	CB	PHE 93	B	16.605	62.527	-2.519	0.52
1294	CG	PHE 93	B	15.862	63.671	-3.167	0.65
1295	CD1	PHE 93	B	14.542	63.901	-2.795	0.60
1296	CD2	PHE 93	B	16.473	64.466	-4.135	0.66
1297	CE1	PHE 93	B	13.873	65.043	-3.237	0.64
1298	CE2	PHE 93	B	15.814	65.627	-4.580	0.66
1299	CZ	PHE 93	B	14.435	65.735	-4.295	0.51
1300	C	PHE 93	B	16.741	64.103	-0.506	0.66
1301	O	PHE 93	B	17.155	65.264	-0.617	0.73
1302	N	ILE 94	B	15.788	63.768	0.315	0.64
1303	CA	ILE 94	B	15.077	64.619	1.252	0.61
1304	CB	ILE 94	B	13.867	63.896	1.897	0.63
1305	CG2	ILE 94	B	13.064	64.661	2.972	0.63

89/98

*Fig. 16*30

Atom	Residue	Chain	X	Y	Z	δ	
1306	CG1	ILE 94	B	12.891	63.334	0.814	0.51
1307	CD	ILE 94	B	11.846	62.434	1.602	0.63
1308	C	ILE 94	B	15.975	65.284	2.289	0.70
1309	O	ILE 94	B	15.535	66.310	2.824	0.75
1310	N	ASN 95	B	16.860	64.542	2.923	0.74
1311	CA	ASN 95	B	17.823	64.972	3.913	0.64
1312	CB	ASN 95	B	17.256	65.197	5.288	0.70
1313	CG	ASN 95	B	18.207	65.971	6.197	0.86
1314	OD1	ASN 95	B	19.322	66.347	5.782	0.79
1315	ND2	ASN 95	B	17.874	66.039	7.487	0.87
1316	C	ASN 95	B	19.132	64.178	3.867	0.65
1317	O	ASN 95	B	19.420	63.340	4.746	0.65
1318	N	PRO 96	B	20.049	64.736	3.096	0.64
1319	CD	PRO 96	B	19.759	65.911	2.228	0.68
1320	CA	PRO 96	B	21.354	64.191	2.805	0.65
1321	CB	PRO 96	B	22.034	65.260	1.943	0.69
1322	CG	PRO 96	B	20.869	65.837	1.180	0.72
1323	C	PRO 96	B	22.214	63.754	3.961	0.72
1324	O	PRO 96	B	23.152	62.938	3.812	0.73
1325	N	GLU 97	B	21.829	64.174	5.140	0.72
1326	CA	GLU 97	B	22.686	64.060	6.338	0.70
1327	CB	GLU 97	B	22.627	65.373	7.121	0.84
1328	CG	GLU 97	B	23.691	66.431	7.069	0.80
1329	CD	GLU 97	B	24.673	66.524	5.952	1.11
1330	OE1	GLU 97	B	25.881	66.300	6.044	1.13
1331	OE2	GLU 97	B	24.198	67.117	4.948	1.11
1332	C	GLU 97	B	22.108	62.957	7.224	0.72
1333	O	GLU 97	B	22.745	62.506	8.192	0.74
1334	N	THR 98	B	20.827	62.673	7.015	0.69
1335	CA	THR 98	B	20.159	61.643	7.830	0.67
1336	CB	THR 98	B	18.594	61.624	7.570	0.65
1337	OG1	THR 98	B	18.303	62.988	7.123	0.77
1338	CG2	THR 98	B	17.827	61.326	8.854	0.78
1339	C	THR 98	B	20.744	60.263	7.527	0.63
1340	O	THR 98	B	20.821	59.417	8.443	0.67
1341	N	VAL 99	B	21.028	60.046	6.249	0.57
1342	CA	VAL 99	B	21.514	58.705	5.846	0.52
1343	CB	VAL 99	B	20.261	57.794	5.871	0.56
1344	CG1	VAL 99	B	19.489	57.979	4.573	0.44
1345	CG2	VAL 99	B	20.512	56.335	6.138	0.62
1346	C	VAL 99	B	22.242	58.813	4.532	0.52
1347	O	VAL 99	B	21.984	59.664	3.666	0.53
1348	N	PRO 100	B	23.279	57.962	4.403	0.53
1349	CD	PRO 100	B	23.638	56.903	5.352	0.54
1350	CA	PRO 100	B	23.966	57.811	3.125	0.49

90/98

*Fig. 16*31

Atom	Residuc	Chain	X	Y	Z	δ	
1351	CB	PRO 100	B	25.101	56.822	3.411	0.50
1352	CG	PRO 100	B	24.727	56.112	4.666	0.47
1353	C	PRO 100	B	23.053	57.286	2.041	0.41
1354	O	PRO 100	B	22.099	56.537	2.250	0.58
1355	N	LYS 101	B	23.591	57.197	0.836	0.48
1356	CA	LYS 101	B	23.004	56.427	-0.241	0.40
1357	CB	LYS 101	B	23.525	56.940	-1.590	0.58
1358	CG	LYS 101	B	22.414	57.692	-2.367	0.69
1359	CD	LYS 101	B	22.889	58.782	-3.271	0.65
1360	CE	LYS 101	B	24.415	58.943	-3.297	0.61
1361	NZ	LYS 101	B	24.770	59.592	-4.589	0.76
1362	C	LYS 101	B	23.492	54.959	-0.048	0.56
1363	O	LYS 101	B	24.250	54.763	0.922	0.54
1364	N	PRO 102	B	22.672	54.006	-0.475	0.57
1365	CD	PRO 102	B	21.367	54.200	-1.164	0.48
1366	CA	PRO 102	B	23.080	52.604	-0.428	0.59
1367	CB	PRO 102	B	21.934	51.832	-1.028	0.43
1368	CG	PRO 102	B	21.075	52.823	-1.736	0.54
1369	C	PRO 102	B	24.426	52.376	-1.076	0.56
1370	O	PRO 102	B	24.877	53.077	-2.006	0.63
1371	N	CYS 103	B	25.031	51.241	-0.744	0.58
1372	CA	CYS 103	B	26.294	50.811	-1.400	0.48
1373	CB	CYS 103	B	27.404	50.638	-0.384	0.75
1374	SG	CYS 103	B	27.517	48.995	0.344	0.81
1375	C	CYS 103	B	26.108	49.649	-2.355	0.55
1376	O	CYS 103	B	25.113	48.919	-2.353	0.53
1377	N	CYS 104	B	26.834	49.723	-3.461	0.32
1378	CA	CYS 104	B	26.958	48.724	-4.468	0.48
1379	C	CYS 104	B	27.702	47.477	-3.963	0.60
1380	O	CYS 104	B	28.863	47.528	-3.570	0.60
1381	CB	CYS 104	B	27.694	49.397	-5.594	0.32
1382	SG	CYS 104	B	27.689	48.486	-7.124	0.54
1383	N	ALA 105	B	27.039	46.332	-4.024	0.51
1384	CA	ALA 105	B	27.391	45.157	-3.222	0.51
1385	CB	ALA 105	B	26.840	45.289	-1.838	0.51
1386	C	ALA 105	B	26.862	43.953	-4.017	0.51
1387	O	ALA 105	B	26.240	44.236	-5.068	0.57
1388	N	PRO 106	B	27.562	42.850	-3.866	0.59
1389	CD	PRO 106	B	28.577	42.554	-2.822	0.51
1390	CA	PRO 106	B	27.352	41.676	-4.739	0.59
1391	CB	PRO 106	B	28.482	40.709	-4.321	0.54
1392	CG	PRO 106	B	29.464	41.531	-3.559	0.50
1393	C	PRO 106	B	25.996	41.079	-4.332	0.58
1394	O	PRO 106	B	25.637	41.105	-3.154	0.54
1395	N	THR 107	B	25.222	40.677	-5.306	0.60

91/98

Fig. 16|32

Atom	Residue	Chain	X	Y	Z	δ	
1396	CA	THR 107	B	23.880	40.104	-4.913	0.65
1397	CB	THR 107	B	22.823	40.695	-5.928	0.45
1398	OG1	THR 107	B	23.160	40.092	-7.208	0.68
1399	CG2	THR 107	B	22.802	42.209	-6.091	0.56
1400	C	THR 107	B	24.011	38.593	-5.133	0.68
1401	O	THR 107	B	23.316	37.753	-4.543	0.68
1402	N	GLN 108	B	25.040	38.240	-5.902	0.67
1403	CA	GLN 108	B	25.358	36.820	-6.067	0.64
1404	CB	GLN 108	B	24.534	36.137	-7.105	0.58
1405	CG	GLN 108	B	24.889	36.327	-8.547	0.70
1406	CD	GLN 108	B	24.534	35.087	-9.358	0.99
1407	OE1	GLN 108	B	24.739	33.953	-8.916	0.83
1408	NE2	GLN 108	B	23.905	35.307	-10.508	0.94
1409	C	GLN 108	B	26.848	36.536	-6.135	0.63
1410	O	GLN 108	B	27.523	37.048	-7.047	0.64
1411	N	LEU 109	B	27.248	35.607	-5.288	0.60
1412	CA	LEU 109	B	28.574	34.966	-5.307	0.65
1413	CB	LEU 109	B	29.033	35.101	-3.831	0.55
1414	CG	LEU 109	B	29.122	36.560	-3.407	0.59
1415	CD1	LEU 109	B	28.855	36.704	-1.936	0.68
1416	CD2	LEU 109	B	30.501	37.120	-3.746	0.56
1417	C	LEU 109	B	28.563	33.511	-5.774	0.70
1418	O	LEU 109	B	27.843	32.695	-5.142	0.75
1419	N	ASN 110	B	29.599	33.112	-6.507	0.67
1420	CA	ASN 110	B	29.915	31.719	-6.849	0.65
1421	CB	ASN 110	B	30.080	31.445	-8.350	0.59
1422	CG	ASN 110	B	28.797	31.687	-9.115	0.79
1423	OD1	ASN 110	B	28.778	31.854	-10.340	0.90
1424	ND2	ASN 110	B	27.734	31.706	-8.302	0.79
1425	C	ASN 110	B	31.162	31.203	-6.116	0.65
1426	O	ASN 110	B	32.020	31.945	-5.641	0.61
1427	N	ALA 111	B	31.238	29.877	-6.127	0.68
1428	CA	ALA 111	B	32.307	29.116	-5.489	0.64
1429	CB	ALA 111	B	31.831	27.798	-4.906	0.59
1430	C	ALA 111	B	33.458	28.918	-6.467	0.57
1431	O	ALA 111	B	33.295	29.038	-7.697	0.54
1432	N	ILE 112	B	34.629	28.709	-5.879	0.54
1433	CA	ILE 112	B	35.738	28.160	-6.743	0.47
1434	CB	ILE 112	B	36.886	29.173	-6.985	0.63
1435	CG2	ILE 112	B	36.603	30.359	-7.923	0.68
1436	CG1	ILE 112	B	37.619	29.600	-5.712	0.56
1437	CD	ILE 112	B	39.128	29.945	-5.888	0.64
1438	C	ILE 112	B	36.186	26.877	-5.997	0.45
1439	O	ILE 112	B	36.185	26.857	-4.747	0.54
1440	N	SER 113	B	36.724	25.995	-6.770	0.44

92/98

*Fig. 16*33

Atom	Residue	Chain	X	Y	Z	δ	
1441	CA	SER 113	B	37.490	24.802	-6.330	0.61
1442	CB	SER 113	B	36.951	23.581	-7.102	0.63
1443	OG	SER 113	B	35.727	23.125	-6.515	0.62
1444	C	SER 113	B	38.962	25.003	-6.694	0.53
1445	O	SER 113	B	39.320	25.356	-7.828	0.52
1446	N	VAL 114	B	39.833	24.614	-5.778	0.55
1447	CA	VAL 114	B	41.265	24.576	-6.121	0.58
1448	CB	VAL 114	B	41.961	25.809	-5.514	0.64
1449	CG1	VAL 114	B	41.157	27.093	-5.667	0.49
1450	CG2	VAL 114	B	42.328	25.534	-4.094	0.55
1451	C	VAL 114	B	41.871	23.235	-5.686	0.60
1452	O	VAL 114	B	41.581	22.634	-4.637	0.51
1453	N	LEU 115	B	42.814	22.811	-6.510	0.49
1454	CA	LEU 115	B	43.490	21.511	-6.182	0.58
1455	CB	LEU 115	B	43.748	20.883	-7.603	0.55
1456	CG	LEU 115	B	44.210	19.423	-7.568	0.57
1457	CD1	LEU 115	B	43.121	18.518	-7.036	0.59
1458	CD2	LEU 115	B	44.655	19.035	-8.975	0.52
1459	C	LEU 115	B	44.739	21.880	-5.403	0.47
1460	O	LEU 115	B	45.582	22.571	-5.955	0.57
1461	N	TYR 116	B	44.954	21.369	-4.196	0.57
1462	CA	TYR 116	B	46.186	21.698	-3.479	0.69
1463	CB	TYR 116	B	45.946	22.841	-2.480	0.70
1464	CG	TYR 116	B	45.134	22.457	-1.269	0.46
1465	CD1	TYR 116	B	43.762	22.395	-1.272	0.57
1466	CE1	TYR 116	B	43.062	21.927	-0.154	0.66
1467	CD2	TYR 116	B	45.764	22.284	-0.050	0.57
1468	CE2	TYR 116	B	45.120	21.791	1.067	0.52
1469	CZ	TYR 116	B	43.748	21.620	1.016	0.76
1470	OH	TYR 116	B	43.110	21.288	2.184	0.78
1471	C	TYR 116	B	46.810	20.454	-2.849	0.69
1472	O	TYR 116	B	46.136	19.420	-2.752	0.69
1473	N	PHE 117	B	48.067	20.582	-2.485	0.56
1474	CA	PHE 117	B	48.891	19.633	-1.793	0.60
1475	CB	PHE 117	B	50.185	19.204	-2.413	0.64
1476	CG	PHE 117	B	51.022	20.135	-3.206	0.82
1477	CD1	PHE 117	B	51.097	20.008	-4.599	0.91
1478	CD2	PHE 117	B	51.946	20.970	-2.561	1.11
1479	CE1	PHE 117	B	51.960	20.804	-5.343	1.12
1480	CE2	PHE 117	B	52.823	21.783	-3.292	1.03
1481	CZ	PHE 117	B	52.837	21.686	-4.697	1.02
1482	C	PHE 117	B	48.967	19.911	-0.315	0.62
1483	O	PHE 117	B	49.562	20.918	0.059	0.72
1484	N	ASP 118	B	48.287	19.095	0.474	0.67
1485	CA	ASP 118	B	48.220	19.309	1.922	0.67

93/98

*Fig. 16*34

Atom	Residue	Chain	X	Y	Z	δ	
1486	CB	ASP 118	B	47.036	18.679	2.610	0.73
1487	CG	ASP 118	B	47.265	17.250	3.079	0.82
1488	OD1	ASP 118	B	46.423	16.701	3.798	0.72
1489	OD2	ASP 118	B	48.369	16.720	2.831	0.60
1490	C	ASP 118	B	49.577	18.996	2.526	0.68
1491	O	ASP 118	B	50.536	18.660	1.826	0.75
1492	N	ASP 119	B	49.577	18.988	3.851	0.76
1493	CA	ASP 119	B	50.837	18.919	4.614	0.83
1494	CB	ASP 119	B	50.693	19.619	5.961	0.94
1495	CG	ASP 119	B	49.368	19.278	6.634	1.08
1496	OD1	ASP 119	B	49.132	18.110	6.991	1.04
1497	OD2	ASP 119	B	48.509	20.186	6.627	1.06
1498	C	ASP 119	B	51.341	17.482	4.673	0.87
1499	O	ASP 119	B	52.554	17.249	4.854	0.90
1500	N	SER 120	B	50.429	16.546	4.448	0.85
1501	CA	SER 120	B	50.821	15.123	4.389	0.80
1502	CB	SER 120	B	49.833	14.202	5.030	0.83
1503	OG	SER 120	B	48.833	14.876	5.781	1.02
1504	C	SER 120	B	51.188	14.778	2.957	0.80
1505	O	SER 120	B	51.840	13.749	2.686	0.83
1506	N	SER 121	B	50.860	15.712	2.064	0.73
1507	CA	SER 121	B	51.273	15.519	0.653	0.69
1508	CB	SER 121	B	52.529	14.649	0.607	0.66
1509	OG	SER 121	B	53.646	15.408	1.016	0.85
1510	C	SER 121	B	50.155	14.886	-0.163	0.63
1511	O	SER 121	B	50.340	14.556	-1.332	0.69
1512	N	ASN 122	B	49.022	14.741	0.477	0.65
1513	CA	ASN 122	B	47.764	14.427	-0.218	0.63
1514	CB	ASN 122	B	46.738	14.164	0.897	0.69
1515	CG	ASN 122	B	47.397	13.265	1.955	0.73
1516	OD1	ASN 122	B	47.766	12.140	1.597	0.73
1517	ND2	ASN 122	B	47.187	13.541	3.240	0.75
1518	C	ASN 122	B	47.412	15.553	-1.176	0.67
1519	O	ASN 122	B	47.620	16.728	-0.852	0.63
1520	N	VAL 123	B	46.918	15.189	-2.346	0.66
1521	CA	VAL 123	B	46.429	16.123	-3.360	0.61
1522	CB	VAL 123	B	46.774	15.543	-4.744	0.58
1523	CG1	VAL 123	B	45.814	16.141	-5.758	0.59
1524	CG2	VAL 123	B	48.224	15.828	-5.089	0.56
1525	C	VAL 123	B	44.917	16.290	-3.220	0.63
1526	O	VAL 123	B	44.237	15.263	-3.098	0.65
1527	N	ILE 124	B	44.491	17.451	-2.758	0.70
1528	CA	ILE 124	B	43.113	17.704	-2.302	0.65
1529	CB	ILE 124	B	43.231	18.240	-0.813	0.67
1530	CG2	ILE 124	B	41.850	18.575	-0.198	0.67

94/98

Fig. 16|35

Atom	Residue	Chain	X	Y	Z	δ	
1531	CG1	ILE 124	B	43.938	17.128	-0.011	0.66
1532	CD	ILE 124	B	43.524	16.995	1.468	0.84
1533	C	ILE 124	B	42.385	18.700	-3.208	0.67
1534	O	ILE 124	B	42.895	19.771	-3.582	0.63
1535	N	LEU 125	B	41.110	18.426	-3.448	0.63
1536	CA	LEU 125	B	40.235	19.393	-4.140	0.69
1537	CB	LEU 125	B	39.453	18.619	-5.187	0.67
1538	CG	LEU 125	B	38.491	19.404	-6.067	0.64
1539	CD1	LEU 125	B	39.277	20.179	-7.130	0.49
1540	CD2	LEU 125	B	37.653	18.302	-6.750	0.48
1541	C	LEU 125	B	39.344	20.118	-3.135	0.67
1542	O	LEU 125	B	38.544	19.446	-2.470	0.66
1543	N	LYS 126	B	39.596	21.403	-2.961	0.66
1544	CA	LYS 126	B	38.865	22.238	-2.001	0.59
1545	CB	LYS 126	B	39.604	22.684	-0.788	0.61
1546	CG	LYS 126	B	38.871	23.673	0.128	0.78
1547	CD	LYS 126	B	39.041	23.334	1.601	0.91
1548	CE	LYS 126	B	38.196	24.169	2.543	0.82
1549	NZ	LYS 126	B	37.808	23.383	3.746	0.86
1550	C	LYS 126	B	38.054	23.325	-2.669	0.50
1551	O	LYS 126	B	38.253	23.700	-3.845	0.57
1552	N	LYS 127	B	36.810	23.373	-2.204	0.61
1553	CA	LYS 127	B	35.729	24.235	-2.664	0.60
1554	CB	LYS 127	B	34.375	23.528	-2.620	0.58
1555	CG	LYS 127	B	33.312	24.232	-3.471	0.57
1556	CD	LYS 127	B	31.942	24.017	-2.833	0.62
1557	CE	LYS 127	B	30.852	24.324	-3.845	0.76
1558	NZ	LYS 127	B	31.445	24.255	-5.208	0.83
1559	C	LYS 127	B	35.626	25.471	-1.755	0.60
1560	O	LYS 127	B	35.815	25.364	-0.521	0.49
1561	N	TYR 128	B	35.798	26.599	-2.424	0.56
1562	CA	TYR 128	B	35.838	27.876	-1.669	0.50
1563	CB	TYR 128	B	37.082	28.686	-2.089	0.77
1564	CG	TYR 128	B	38.294	28.241	-1.298	0.64
1565	CD1	TYR 128	B	39.395	27.674	-1.935	0.68
1566	CE1	TYR 128	B	40.468	27.198	-1.174	0.68
1567	CD2	TYR 128	B	38.204	28.154	0.085	0.72
1568	CE2	TYR 128	B	39.254	27.649	0.857	0.77
1569	CZ	TYR 128	B	40.332	27.060	0.200	0.89
1570	OH	TYR 128	B	41.314	26.484	0.961	0.90
1571	C	TYR 128	B	34.587	28.661	-2.110	0.46
1572	O	TYR 128	B	34.335	28.808	-3.308	0.59
1573	N	ARG 129	B	33.692	28.665	-1.150	0.56
1574	CA	ARG 129	B	32.380	29.311	-1.396	0.61
1575	CB	ARG 129	B	31.479	28.769	-0.273	0.74

95/98

Fig. 16|36

Atom	Residue	Chain	X	Y	Z	δ	
1576	CG	ARG 129	B	31.060	27.303	-0.447	0.74
1577	CD	ARG 129	B	29.804	27.006	0.337	0.61
1578	NE	ARG 129	B	28.620	27.257	-0.488	1.06
1579	CZ	ARG 129	B	27.768	28.270	-0.310	1.07
1580	NH1	ARG 129	B	27.618	28.890	0.861	0.91
1581	NH2	ARG 129	B	27.093	28.771	-1.349	0.87
1582	C	ARG 129	B	32.587	30.829	-1.302	0.54
1583	O	ARG 129	B	33.426	31.314	-0.522	0.55
1584	N	ASN 130	B	31.860	31.561	-2.095	0.60
1585	CA	ASN 130	B	31.584	33.006	-2.013	0.59
1586	CB	ASN 130	B	31.088	33.375	-0.610	0.52
1587	CG	ASN 130	B	29.629	32.995	-0.392	0.63
1588	OD1	ASN 130	B	28.825	32.869	-1.318	0.58
1589	ND2	ASN 130	B	29.276	32.510	0.782	0.55
1590	C	ASN 130	B	32.864	33.763	-2.380	0.55
1591	O	ASN 130	B	33.205	34.674	-1.648	0.48
1592	N	MET 131	B	33.418	33.459	-3.520	0.54
1593	CA	MET 131	B	34.730	33.897	-3.974	0.51
1594	CB	MET 131	B	35.632	32.662	-4.327	0.54
1595	CG	MET 131	B	36.159	31.950	-3.137	0.44
1596	SD	MET 131	B	37.229	32.957	-2.052	0.61
1597	CE	MET 131	B	38.802	32.648	-2.915	0.60
1598	C	MET 131	B	34.538	34.709	-5.256	0.52
1599	O	MET 131	B	35.416	35.514	-5.570	0.59
1600	N	VAL 132	B	33.538	34.371	-6.055	0.59
1601	CA	VAL 132	B	33.347	35.038	-7.359	0.60
1602	CB	VAL 132	B	33.434	34.092	-8.517	0.52
1603	CG1	VAL 132	B	33.162	34.633	-9.898	0.48
1604	CG2	VAL 132	B	34.447	32.972	-8.485	0.66
1605	C	VAL 132	B	32.063	35.873	-7.308	0.69
1606	O	VAL 132	B	30.960	35.366	-7.012	0.63
1607	N	VAL 133	B	32.240	37.134	-7.668	0.65
1608	CA	VAL 133	B	31.100	38.075	-7.709	0.62
1609	CB	VAL 133	B	31.530	39.550	-7.591	0.56
1610	CG1	VAL 133	B	30.408	40.468	-8.105	0.53
1611	CG2	VAL 133	B	31.852	39.841	-6.128	0.39
1612	C	VAL 133	B	30.409	37.812	-9.052	0.62
1613	O	VAL 133	B	31.105	38.041	-10.044	0.54
1614	N	ARG 134	B	29.134	37.398	-8.941	0.53
1615	CA	ARG 134	B	28.440	37.117	-10.208	0.56
1616	CB	ARG 134	B	27.709	35.760	-10.158	0.68
1617	CG	ARG 134	B	28.433	34.636	-10.908	0.80
1618	CD	ARG 134	B	28.238	34.700	-12.379	0.79
1619	NE	ARG 134	B	27.810	33.460	-12.976	1.02
1620	CZ	ARG 134	B	26.587	32.977	-13.159	1.10

96/98

Fig. 16\37

Atom	Residue	Chain	X	Y	Z	δ	
1621	NH1	ARG 134	B	25.494	33.408	-12.533	1.08
1622	NH2	ARG 134	B	26.450	31.902	-13.952	1.09
1623	C	ARG 134	B	27.517	38.253	-10.622	0.57
1624	O	ARG 134	B	27.185	38.397	-11.813	0.61
1625	N	ALA 135	B	27.038	39.030	-9.659	0.53
1626	CA	ALA 135	B	26.189	40.203	-9.986	0.44
1627	CB	ALA 135	B	24.713	39.740	-9.934	0.64
1628	C	ALA 135	B	26.352	41.156	-8.774	0.42
1629	O	ALA 135	B	26.515	40.639	-7.673	0.49
1630	N	CYS 136	B	25.926	42.365	-9.022	0.59
1631	CA	CYS 136	B	26.001	43.512	-8.089	0.67
1632	C	CYS 136	B	24.646	44.198	-7.969	0.61
1633	O	CYS 136	B	23.917	44.273	-8.985	0.59
1634	CB	CYS 136	B	26.990	44.534	-8.701	0.61
1635	SG	CYS 136	B	28.689	43.852	-8.673	0.61
1636	N	GLY 137	B	24.362	44.756	-6.808	0.59
1637	CA	GLY 137	B	23.099	45.570	-6.772	0.59
1638	C	GLY 137	B	23.290	46.706	-5.787	0.52
1639	O	GLY 137	B	24.077	46.520	-4.845	0.49
1640	N	CYS 138	B	22.261	47.515	-5.634	0.57
1641	CA	CYS 138	B	22.248	48.539	-4.545	0.48
1642	C	CYS 138	B	21.475	48.015	-3.369	0.56
1643	O	CYS 138	B	20.235	47.865	-3.446	0.55
1644	CB	CYS 138	B	21.556	49.798	-5.123	0.38
1645	SG	CYS 138	B	22.664	50.498	-6.428	0.57
1646	N	HIS 139	B	22.137	48.025	-2.210	0.63
1647	CA	HIS 139	B	21.436	47.660	-0.970	0.54
1648	CB	HIS 139	B	20.837	46.284	-0.842	0.88
1649	CG	HIS 139	B	21.539	45.272	-1.707	0.94
1650	CD2	HIS 139	B	22.879	45.046	-1.811	0.79
1651	ND1	HIS 139	B	20.936	44.549	-2.706	0.91
1652	CE1	HIS 139	B	21.855	43.796	-3.288	0.97
1653	NE2	HIS 139	B	23.034	44.089	-2.774	0.96
1654	C	HIS 139	B	21.888	48.359	0.264	0.58
1655	OT1	HIS 139	B	21.160	48.202	1.268	0.75
1656	OT2	HIS 139	B	22.865	49.139	0.256	0.79
1657	OT	WAT 201	A	31.351	45.516	-2.695	0.57
1658	OT	WAT 202	A	10.574	42.304	-3.269	0.77
1659	OT	WAT 203	A	41.094	47.385	-10.715	0.74
1660	OT	WAT 204	A	-6.527	46.271	-4.416	0.83
1661	OT	WAT 205	A	-7.389	42.963	-2.480	0.75
1662	OT	WAT 206	A	-5.998	42.514	2.104	0.60
1663	OT	WAT 207	A	25.154	37.549	3.436	0.76
1664	OT	WAT 208	A	31.925	33.286	2.732	0.58
1665	OT	WAT 209	A	32.701	43.734	-4.779	0.52

97/98
Fig. 16\38

Atom	Residue	Chain	X	Y	Z	δ	
1666	OT	WAT 210	A	15.485	51.948	13.181	0.60
1667	OT	WAT 211	A	9.829	38.538	6.111	0.71
1668	OT	WAT 212	A	11.550	40.302	3.050	0.58
1669	OT	WAT 213	A	42.134	46.885	5.506	0.73
1670	OT	WAT 214	A	37.738	52.318	-0.352	0.72
1671	OT	WAT 215	A	40.582	52.333	-4.324	0.66
1672	OT	WAT 216	A	22.375	54.496	15.975	0.63
1673	OT	WAT 217	A	49.983	39.399	2.310	0.69
1674	OT	WAT 218	A	5.369	58.756	12.045	0.61
1675	OT	WAT 219	A	0.867	40.439	5.311	0.67
1676	OT	WAT 220	A	25.522	37.902	-0.824	0.80
1677	OT	WAT 221	A	12.228	59.495	7.513	0.81
1678	OT	WAT 222	A	10.798	47.556	11.898	0.67
1679	OT	WAT 223	A	0.494	40.963	0.254	0.75
1680	OT	WAT 224	A	33.591	41.614	-1.644	0.50
1681	OT	WAT 225	A	24.730	59.387	8.279	0.74
1682	OT	WAT 226	A	17.020	38.835	3.348	0.73
1683	OT	WAT 227	A	34.395	49.780	3.674	0.61
1684	OT	WAT 228	A	6.972	43.390	-2.094	0.59
1685	OT	WAT 229	A	25.493	43.453	12.008	0.68
1686	OT	WAT 230	A	31.349	49.756	-3.421	0.70
1687	OT	WAT 231	A	2.519	49.133	-4.219	0.80
1688	OT	WAT 232	A	24.405	52.256	3.441	0.61
1689	OT	WAT 233	A	-0.363	65.457	3.148	0.79
1690	OT	WAT 201	B	23.742	49.909	2.695	0.57
1691	OT	WAT 202	B	31.349	30.309	3.269	0.77
1692	OT	WAT 203	B	20.489	59.281	10.715	0.74
1693	OT	WAT 204	B	43.335	17.483	4.416	0.83
1694	OT	WAT 205	B	40.901	15.082	2.480	0.75
1695	OT	WAT 206	B	39.817	16.063	-2.104	0.60
1696	OT	WAT 207	B	19.941	40.558	-3.436	0.76
1697	OT	WAT 208	B	12.864	44.291	-2.732	0.58
1698	OT	WAT 209	B	21.524	50.187	4.779	0.52
1699	OT	WAT 210	B	37.246	39.384	-13.181	0.60
1700	OT	WAT 211	B	28.460	27.781	-6.111	0.71
1701	OT	WAT 212	B	29.127	30.154	-3.050	0.58
1702	OT	WAT 213	B	19.536	59.931	-5.506	0.73
1703	OT	WAT 214	B	26.439	58.841	0.352	0.72
1704	OT	WAT 215	B	25.030	61.311	4.324	0.66
1705	OT	WAT 216	B	36.007	46.625	-15.975	0.63
1706	OT	WAT 217	B	9.129	62.986	-2.310	0.69
1707	OT	WAT 218	B	48.199	34.028	-12.045	0.61
1708	OT	WAT 219	B	34.587	20.970	-5.311	0.67
1709	OT	WAT 220	B	20.063	41.054	0.824	0.80
1710	OT	WAT 221	B	45.410	40.337	-7.513	0.81

98/98

Fig. 16|39

Atom		Residue	Chain	X	Y	Z	δ
1711	OT	WAT 222	B	35.785	33.129	-11.898	0.67
1712	OT	WAT 223	B	35.228	20.909	-0.254	0.75
1713	OT	WAT 224	B	19.243	49.897	1.644	0.50
1714	OT	WAT 225	B	39.065	51.110	-8.279	0.74
1715	OT	WAT 226	B	25.122	34.157	-3.348	0.73
1716	OT	WAT 227	B	25.913	54.677	-3.674	0.61
1717	OT	WAT 228	B	34.091	27.733	2.094	0.59
1718	OT	WAT 229	B	24.885	43.804	-12.008	0.68
1719	OT	WAT 230	B	27.415	52.027	3.421	0.70
1720	OT	WAT 231	B	41.291	26.748	4.219	0.80
1721	OT	WAT 232	B	33.052	47.263	-3.441	0.61
1722	OT	WAT 233	B	56.869	32.414	-3.148	0.79

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/51, G06F 17/50, C07K 1/00	A3	(11) International Publication Number: WO 97/26277 (43) International Publication Date: 24 July 1997 (24.07.97)
(21) International Application Number: PCT/US97/01071 (22) International Filing Date: 22 January 1997 (22.01.97) (30) Priority Data: 08/589,552 22 January 1996 (22.01.96) US (71) Applicants: CREATIVE BIOMOLECULES, INC. [US/US]; 45 South Street, Hopkinton, MA 01748 (US). BRANDEIS UNIVERSITY [US/US]; 45 South Street, Waltham, MA 02254-9110 (US). (72) Inventors: KECK, Peter; 50 Dolan Road, Millbury, MA 01527 (US). GRIFFITH, Diana, L.; 15 Woodridge Circle, Weston, MA 02193 (US). CARLSON, William, D.; 40 Black Oak Road, Weston, MA 02193 (US). RUEGER, David, C.; 19 Downey Street, Hopkinton, MA 01748 (US). SAMPATH, Kuber, T.; 6 Spring Street, Medway, MA 02053 (US). (74) Agent: GREENHALGH, Duncan, A.; Testa, Hurwitz & Thibault, L.L.P., High Street Tower, 125 High Street, Boston, MA 02110 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> (88) Date of publication of the international search report: 25 September 1997 (25.09.97)
(54) Title: MORPHOGEN ANALOGS AND METHODS FOR PRODUCING THEM (57) Abstract The invention disclosed herein provides methods and compositions for the computer-assisted design of morphogen analogs. Practice of the invention is enabled by the use of at least a portion of the atomic co-ordinates defining the three-dimensional structure of human osteogenic protein-1 (hOP-1) as a starting point in the design of the morphogen analogs. In addition, the invention provides methods for producing morphogen analogs of interest, and methods for testing whether the resulting analogs mimic or agonize human OP-1-like biological activity. The invention also provides a family of morphogen analogs produced by such methods.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 97/01071

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K14/51 G06F17/50 C07K1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C07K G06F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 92 07073 A (CREATIVE BIOMOLECULES) 30 April 1992 see page 8, line 8 - page 9, line 9; claim 21	22,23
X	WO 89 09788 A (CREATIVE BIOMOLECULES) 19 October 1989 see claims 21-42	22,23

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

14 July 1997

Date of mailing of the international search report

23.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (-31-70) 340-3016

Authorized officer

Masturzo, P

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/US 97/01071

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 93, no. 2, 23 January 1996, WASHINGTON US, pages 878-883, XP002035127 D L GRIFFITH ET AL.: "Three-dimensional structure of recombinant human osteogenic proteins 1; structural paradigm for the transforming growth factor beta superfamily" see the whole document -----</p>	1-21

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/01071

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9207073 A	30-04-92	AU 663689 B	19-10-95
		AU 8900091 A	20-05-92
		CA 2094027 A	19-04-92
		EP 0643767 A	22-03-95
		JP 7500487 T	19-01-95
		US 5468845 A	21-11-95
		US 5344654 A	06-09-94
		US 5354557 A	11-10-94
<hr/>			
WO 8909788 A	19-10-89	US 4968590 A	06-11-90
		US 5011691 A	30-04-91
		AT 142644 T	15-09-96
		AT 134647 T	15-03-96
		AU 628050 B	10-09-92
		AU 3444989 A	03-11-89
		AU 618357 B	19-12-91
		AU 3530589 A	03-11-89
		CA 1338663 A	22-10-96
		DE 68925773 D	04-04-96
		DE 68925773 T	17-10-96
		DE 68927153 D	17-10-96
		DE 68927153 T	06-03-97
		EP 0372031 A	13-06-90
		EP 0362367 A	11-04-90
		EP 0714665 A	05-06-96
		EP 0723013 A	24-07-96
		JP 8336390 A	24-12-96
		JP 8322570 A	10-12-96
		JP 8187084 A	23-07-96
		JP 2522568 B	07-08-96
		JP 3500655 T	14-02-91
		JP 2548414 B	30-10-96
		JP 3502579 T	13-06-91
		WO 8909787 A	19-10-89
		US 5468845 A	21-11-95
		US 5496552 A	05-03-96
		US 5108753 A	28-04-92
		US 5182365 A	26-01-93
		US 5250302 A	05-10-93
		US 5266683 A	30-11-93

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/01071

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 8909788 A		US 5344654 A	06-09-94
		US 5324819 A	28-06-94
		US 5354557 A	11-10-94
		US 5258494 A	02-11-93
		AT 124051 T	15-07-95
		AU 627850 B	03-09-92
		AU 5174790 A	26-09-90
		DE 69020254 D	27-07-95
		DE 69020254 T	25-01-96
		EP 0411105 A	06-02-91
		ES 2074158 T	01-09-95
		JP 8024710 B	13-03-96
		JP 3504736 T	17-10-91
		WO 9010018 A	07-09-90
		US 4975526 A	04-12-90
		US 5171574 A	15-12-92
		US 5162114 A	10-11-92
